

Diversity of Ceftazidime-Avibactam Resistance Mechanism in KPC2-Producing *Klebsiella pneumoniae* Under Antibiotic Selection Pressure

Min Jiang*, Bin Sun*, Yong Huang, Chengyang Liu, Yan Wang, Yanli Ren, Yuhong Zhang, Yuning Wang, Di Mu

Department of Laboratory Medicine, The Second Affiliated Hospital of Chongqing Medical University, Chongqing, People's Republic of China

*These authors contributed equally to this work

Correspondence: Di Mu; Yuning Wang, Department of Laboratory Medicine, The Second Affiliated Hospital of Chongqing Medical University, No. 74 Linjiang Road, Yuzhong District, Chongqing, People's Republic of China, Tel/Fax +86 23-62887386, Email 303921@hospital.cqmu.edu.cn; Wangyuning@hospital.cqmu.edu.cn

Purpose: The aim of this study was to understand the resistance mechanism of ceftazidime/avibactam (CZA) in carbapenem-resistant *Klebsiella pneumoniae* under antibiotic selection pressure.

Patients and Methods: Four CZA-resistant *Klebsiella pneumoniae* strains were isolated from two patients, and six CZA-resistant strains that were produced in vitro were screened from 25 carbapenem-resistant *Klebsiella pneumoniae* strains. The mechanisms of resistance to CZA of these strains were characterized by PCR and Sanger sequencing.

Results: CZA-resistant *Klebsiella pneumoniae* with different resistance mechanisms (including upregulation of the expression of efflux pumps and KPC variants (KPC-14, KPC-44)) were isolated from the same patient (patient 1). In patient 2, the resistance mechanism of CZA-resistant *Klebsiella pneumoniae* was the mutation of KPC-2 to KPC-33. In addition, among the CZA-resistant *Klebsiella pneumoniae* that were produced in vitro, we found 3 new KPC variants: KPC-86 (D179G), KPC-87 (GT241A) and KPC-88 (G523T).

Conclusion: In this study, although the CZA-resistant bacteria originated from only two clinical patients, four different mechanisms of CZA resistance were detected. In the in vitro induction experiment, the mechanisms of resistance to CZA in strains from different patients were also different. The above result implies that the mechanisms of resistance to CZA are generally random and diverse. Therefore, elucidating the mechanism of resistance to CZA can provide a certain theoretical basis for the effective response of CZA-resistant strains and the selection of antibiotics.

Keywords: ceftazidime/avibactam-resistance, KPC mutants, antibiotic pressure, *Klebsiella pneumoniae*, resistance mechanisms

Introduction

Due to restrictions on antibiotic options, carbapenem-resistant *Enterobacteriales* (CRE) has become a serious threat to human health. Ceftazidime/avibactam (CZA) is a new β -lactam antibacterial combination (2 g ceftazidime and 0.5 g avibactam) that was approved by the United States Food and Drug Administration in 2015 for the treatment of infections caused by CRE.¹ CZA could effectively inhibit the *enterobacteria* that produce A-, C-, and D-type β -lactamases (include KPCs, AmpC and OXA-48).²

According to clinical research reports, the application of CZA has significantly improved the clinical cure rate and survival rate of patients with CRE infection, and CZA is superior to other drug treatment options.³ According to a survey of 210 hospitals released by the United States in 2020, the number of CZA hospital prescriptions increased nearly 18-fold from 2015 to 2017. However, the first case of CZA-resistant bacteria was reported in the United States in 2016. From 2016 to 2021, multiple emergent cases of strains resistant to CZA were reported worldwide. Among them, *Klebsiella pneumoniae* is one of the main resistant bacteria.⁴⁻⁶ *Klebsiella pneumoniae* was considered an opportunistic nosocomial

pathogen and had attracted widespread attention due to its disease severity and antibiotic resistance. *Klebsiella pneumoniae* was highly virulent and multidrug-resistant and could cause life-threatening infections.^{7,8}

In China, carbapenem-resistant *Klebsiella pneumoniae* increased nearly 10-fold during the decade from 2010 to 2021. CZA was approved by the China National Medical Products Administration (NMPA) to enter the Chinese clinic on May 21, 2019. In the first half of 2020, the drug resistance rate of CZA reached 14.9% according to the analysis of the drug data from 4395 cases of carbapenem-resistant *Klebsiella pneumoniae* obtained by the China Bacterial Resistance Surveillance Network (CHINET) from more than 50 hospitals in China.^{9–11} The emergence of CZA-resistant strains indicates that human health could be seriously threatened, and it also brings new challenges for clinicians.

Recent studies have reported that the resistance mechanisms of carbapenem-resistant *Klebsiella pneumoniae* against CZA mainly include mutations in KPCs (KPC-2, KPC-3), mutations in porins, the effects of efflux pumps, and the variation of the copy number of the KPC-encoding plasmid.^{12–14} However, due to the relatively short time that CZA has been used in clinical practice in China, the number of collections and studies of clinically resistant strains in China is small. Therefore, the resistance mechanism of *Klebsiella pneumoniae* against CZA and new medication regimens in China urgently need to be studied.

In this study, CZA-resistant strains from 2 patients were collected, and the resistance mechanism was analysed. The result showed the same specimen of the same patient could produce CZA resistant strains with multiple resistance mechanism. And the resistance mechanisms of the CZA-resistance *Klebsiella pneumoniae* isolated from the two patients included KPC mutations (KPC-2 was mutated to KPC-44, KPC-14 and KPC-33) and upregulated efflux pumps. However, we discovered that inducing CZA-resistant *Klebsiella pneumoniae* in vitro not only showed the same KPC mutations (KPC-2 was mutated to KPC-44) as the in-vivo CZA-resistant strain, but also three new KPC variants (KPC-86, KPC-87 and KPC-88) were discovered. The above results indicate the diversity of the resistance mechanisms of *Klebsiella pneumoniae* against CZA.

Materials and Methods

Bacterial Strains

Four CZA-resistant *Klebsiella pneumoniae* strains were collected, which were derived from different body fluid samples (ascites, blood and sputum) of two clinical patients in 2020. There was no significant difference in the growth rate and morphology of these strains. The patients were admitted to the Second Affiliated Hospital of Chongqing Medical University of China.

Induction of CZA-Resistant Strains

Twenty-five *Klebsiella pneumoniae* strains, used to induce CZA resistance in vitro, also came from clinical patients. There was no significant difference in the growth rate and morphology of these strains. The initial *Klebsiella pneumoniae* isolates were cultured in 5 mL nutrient broth medium respectively. Fifty microliters of the bacterial solution and CZA (ceftazidime: avibactam 3:2, 50 µg) were added to 1 mL broth medium and cultured for 24 hours in a 35 °C bacterial incubator to induce CZA-resistant strains. Bacterial strains were passaged every day for 10 generations.

Antimicrobial Susceptibilities

The antibiotics imipenem (IMP), meropenem (MER), tigecycline (TGC) and CZA were used to test the sensitivity to drugs of the bacterial strains. Antimicrobial susceptibilities were determined with the broth microdilution method, as previously described by CLSI.¹⁵

Detection of Carbapenemase Type

The detection of the carbapenemase type used the APB(3-aminophenylboronic acid)/EDTA method.¹⁶ This method is based on disk diffusion method in the CLSI guidelines.¹⁵ Briefly, a 0.5 McFarland turbidity of *Klebsiella pneumoniae* was spread on a Mueller-Hinton agar plate, and then 4 tablets of imipenem (10 µg/tablet) were placed on the agar plate. APB (300 µg) and EDTA (292 µg) are KPC-type carbapenemase and metallo β-lactamases inhibitors, respectively, and

they were dropped on two pieces of imipenem tablets. A mixture of APC and EDTA was dropped on another imipenem tablet, and nothing was dropped on the last imipenem tablet. If the difference in the inhibition zone diameter between the enzyme inhibitor mixture and the single drug was more than 5 mm, it could be determined that the tested strain produces class A carbapenemase (mainly KPC enzyme), class B metallo β -lactamases, or both types of carbapenemase.

The detection of the carbapenemase type was also determined with the method of modified carbapenem inactivation method (mCIM) and EDTA-CIM (eCIM), as previously described by CLSI.¹⁵ Briefly, 1 μ L loopful of isolates were mixed in 2mL of tryptone soya broth (TSB). One meropenem disc (10 μ g) was immersed in the suspension and incubated in a 35°C incubator for 4 hours. The *E. coli* ATCC 25922 with 0.5 McFarland turbidity was inoculated on the MHA plate, and the meropenem tablet was taken out of the TSB broth and stuck it on the MHA plate. The plate was incubated at 35 degrees Celsius for 18–24 hours. If the meropenem inhibition zone was 6–15mm, or the diameter of the inhibition zone was 16–18mm and there are scattered colonies, the strain was tested produces carbapenemase. If the mCIM test was positive, the eCIM test was performed. Another 2mL TSB broth tube was added 20 μ L 0.5M EDTA. The rest of the operation steps were the same as mCIM. If the diameter of the inhibition zone of meropenem was \geq 5mm compared with mCIM, the metallo-beta lactamase was considered positive.

Extraction of Bacterial DNA and Screening of Drug Resistance Genes

Bacterial genomic DNA was extracted using a bacterial genome extraction kit (Thermo Scientific, China). The KPC resistance genes of drug-resistant strains were detected by PCR according to the research of Dallenne et al.¹⁷ The positive products amplified by PCR were sent to Shanghai Sangon Biological Engineering Technology Co., Ltd. for Sanger sequencing. The sequences were compared with those deposited in GenBank database (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>).

Multi-Locus Sequence Typing (MLST)

MLST typing could provide a valuable method and clear data for epidemiological studies of *Klebsiella pneumoniae* isolates and could represent different spatial and temporal dynamics of *Klebsiella pneumoniae*. Seven housekeeping genes (*tonB* (plasma perimeter energy sensor), *gapA* (glyceraldehyde 3-phosphate dehydrogenase), *mdh* (malate dehydrogenase), *infB* (translation initiation factor 2), *rpoB* (beta subunit of RNA polymerase), *pgi* (phosphoglucose isomerase) and *phoE* (phospho E)) were used for MLST PCR amplification, and the PCR products were sequenced and analyzed according to Pasteur MLST site (<https://bigsd.b.pasteur.fr/klebsiella/klebsiella.html>).¹⁸

MIC Determination of the Efflux Pump Inhibitors (CCCP)

Carbonyl cyanochlorophenylhydrazone (CCCP) is an efflux pump inhibitor that was purchased from Sigma–Aldrich (St Louis, USA). MIC determination of CCCP was performed with the broth microdilution method as previously described.¹⁹ Briefly, CCCP solutions with concentrations of 256, 128, 64, 32, 16, 8, 4, 2, 1, 0.5, 0.25, and 0 μ g/mL were added to 96-well plates in sequence. Then, 50 μ L of bacterial suspension with optical density equivalent to 0.5 McFarland was added dropwise to each well of 96-well plates containing different concentrations of CCCP. The 96-well plate was placed in a 37 °C CO₂ incubator and incubated for 16–20 hours. The highest CCCP concentration (8 μ g/mL) with obvious bacterial growth was selected as the optimal concentration for the efflux pump inhibition experiment.

CZA-CCCP Disc Synergy Test

Then, CCCP (32 μ g/mL, 50 μ L/well) was mixed with different concentrations of CZA (64, 32, 16, 8, 4, 2, 1, 0.5, 0.25, and 0 μ g/mL, 50 μ L/well) and configured as a mixed solution. At this time, the final concentration of CCCP was 16 μ g/mL. Then, a mixed solution (50 μ L) was added to the bacterial suspension (0.5 McFarland, 50 μ L/well) and mixed thoroughly. At this time, the final concentration of CCCP was 8 μ g/mL. Compared with the control group (CZA), the MIC value of the experimental group (CZA and CCCP) decreased by more than two gradients, and thus, the result was determined to be positive.

Immune-Enzymatic Test

KPC was detected with a colloidal gold immunochromatographic assay using the NG-test CARBA5 kit (NG-biotech, France), and the experimental protocol was performed according to the product manual.

Conjugation Experiment

The CZA-resistant strains with KPC mutants (KPC-86, KPC-87 and KPC-88) were used as donors. And rifampicin-resistant *E. coli* (EC600) was used as the recipient. The CZA-resistant strains (KPC-86, KPC-87 and KPC-88) also exhibited ampicillin resistance. Transconjugants were selected on MH agar plates supplemented with ampicillin (128 µg/mL) and rifampicin (256 µg/mL). *E. coli* EC600 transconjugants were identified using the mass spectrum system, and KPC mutations were further confirmed by PCR and Sanger sequencing. The conjugation efficiency was measured and calculated following the protocol in <https://openwetware.org/wiki/conjugation>.

Results

Clinical Case Report

A 40-year-old male patient was admitted to the Second Affiliated Hospital of Chongqing Medical University on April 27, 2020 due to upper abdominal pain for more than one month and fever for three days. He was diagnosed with acute severe pancreatitis and sepsis. Carbapenem-resistant and CZA-sensitive *Klebsiella pneumoniae* were isolated from the peritoneal ascites, blood and sputum (strain number: ascites LY0B1, blood LY0B2 and sputum LY0B3, Table 1). The three primary isolates all produced carbapenemase and the enzyme type was KPC-2. Tigecycline plus meropenem was used to fight infection for 14 days from April 27 to May 10, but the patient's condition did not improve. On May 10, the patient was treated with CZA (ceftazidime/avibactam, 2 g/0.5 g, tid). On May 20, the ascites and sputum were taken again for culture (Figure 1, left panel).

Another case was a 75-year-old male patient who was admitted to the Second Affiliated Hospital of Chongqing Medical University on July 13, 2020 due to severe pneumonia. The patient's alveolar lavage fluid was cultured with carbapenem-resistant *Klebsiella pneumoniae* (strain number: HM0B1), and the type of carbapenemase was KPC-2. From July 15 to August 5, the patient was treated with CZA (ceftazidime/avibactam, 2 g/0.5 g, tid) for anti-infection. On July 25, CZA-resistant *Klebsiella pneumoniae* (strain number: HM0F1) was cultured from sputum (Figure 1, right panel).

The Mechanisms of Resistance to CZA of the Clinical Isolates

Interestingly, three CZA-resistant strains (LY0F1-1, LY0F1-2 and LY0F2) with different resistance mechanisms were isolated from patient 1. LY0F1-1 and LY0F1-2 were isolated from the same drainage sample (Figure 2A). The antimicrobial susceptibility

Table 1 Antimicrobial Susceptibility and Resistance Mechanism of *Klebsiella pneumoniae* from Patients

Isolate	MIC (µg/mL)				Carbapenemases	MLST	ESBL	AmpC	Efflux Pump
	IMP	MEM	TGC	CZA					
LY0B1	≥64	≥64	0.5	0.5	KPC-2	ST11	–	–	/
LY0B2	≥64	≥64	0.5	0.5	KPC-2	ST11	–	–	/
LY0B3	≥64	≥64	0.5	0.5	KPC-2	ST11	–	–	/
LY0F1-1	≥64	≥64	1	128	KPC-2	ST11	–	–	↑
LY0F1-2	0.25	2	0.5	128	KPC -14	ST11	TEM	–	/
LY0F2	8	32	0.5	128	KPC-44	ST11	–	–	/
HM0B1	≥64	≥64	1	0.5	KPC-2	ST11	–	–	/
HM0F1	0.25	2	1	256	KPC-33	ST11	TEM	–	/

Abbreviations: IMP, imipenem; MEM, meropenem; TGC, tigecycline; CZA, ceftazidime/avibactam; MLST, Multiple-locus sequence typing; ESBL, extended-spectrum beta-lactamase; AmpC, AmpC beta lactamase.




Figure 1 Schedule of antibacterial treatment and isolation of *Klebsiella pneumoniae* strains. The red arrows represent the time period of medication. The blue arrow represents the specimen collection at the corresponding time.

Abbreviations: IMP, imipenem; MEM, meropenem; CZA, ceftazidime/avibactam.




Figure 2 Isolation of CZA-resistant *Klebsiella pneumoniae* and detection of the enzyme type of the strain. (A) The top panel shows that CZA-resistant *Klebsiella pneumoniae* was isolated from the puncture fluid sample of patient 1 after 10 days of CZA administration, and the sensitivity of CZA, MEM, and IMP was tested by the K-B method. The bottom panel in (A) and (B) shows that the enzyme types of isolates LY0F1-1, LY0F1-2 and HM0F1 were detected by enzyme inhibitor enhancement experiments and mCIM and eCIM experiments.

Abbreviations: MEM, meropenem; TGC, tigecycline; CZA, ceftazidime/avibactam; mCIM, modified carbapenem inactivation method; eCIM, EDTA-modified carbapenem inactivation method.

results are shown in Table 1. LY0F1-1 exhibited IMP and MER resistance (MIC, 64 µg/mL), while LY0F1-2 was susceptible to IMP and intermediate to MER (MIC, 0.25 µg/mL and 2 µg/mL, respectively) (Table 1). The strain cultured from the sputum specimens of patient 1 (LY0F2) was also resistant to CZA, but the MICs of IMP and MER were 8 µg/mL and 32 µg/mL, respectively (Table 1). Therefore, the three CZA-resistant strains isolated from the same patient have different MICs for IMP and MER. The antimicrobial susceptibility results indicated that the HM0F1 isolate from patient 2 was resistant to CZA, and the MIC values of IMP and MER were 0.25 µg/mL and 2 µg/mL, respectively (Table 1).

To research the mechanism of CZA resistance in the above four KPC-producing *Klebsiella pneumoniae* strains, the activity and genotype of carbapenemase were detected. The results showed that the carbapenemase type of the CZA-resistant strain LY0F1-1 was KPC-2, and the expression of KPC-2 did not increase (Figure 2A). However, the efflux pump was upregulated in the LY0F1-1 strain (Table 1). We speculated that LY0F1-1 may also have other resistance mechanisms, which need to be further studied. The carbapenemase types in the CZA-resistant strains LY0F1-2, LY0F2 and HM0F1 were mutated from the original KPC-2 to KPC-14, KPC-44 and KPC-33, respectively, which was the mechanism of CZA resistance in the three isolates (Table 1). The mCIM and eCIM tests did not find that LY0F1-2, LY0F2 and HM0F1 produced serine carbapenemase and metallo β -lactamases (Figure 2A and B), but the immune-enzymatic test for the KPC enzyme was positive. KPC gene was detected to be positive in both primary strains and CZA-resistant strains by PCR (Supplementary Figure 1A).

The above results indicate that the same specimen (LY0F1-1 and LY0F1-2) from the same patient could produce CZA-resistant strains with multiple resistance mechanisms (upregulation of the efflux pump, KPC-14), which shows the diversity of CZA resistance mechanisms.

The Resistance Mechanism of CZA-Resistant *Klebsiella pneumoniae* That Were Produced in vitro

To verify the diversity of KPC mutations in CZA-resistant *Klebsiella pneumoniae* under CZA drug selection. Twenty-five strains of carbapenem-resistant *Klebsiella pneumoniae* were induced with CZA resistance in vitro. The results showed that 6 strains developed CZA resistance, and the resistance rate was as high as 24%. And *blaKPC* was detected by PCR (Supplementary Figure 1B). As shown in Table 2, the carbapenemase genotypes of the primary strains were all *blaKPC*-

Table 2 Antimicrobial Susceptibility and Resistance Mechanism of CZA-Resistant *Klebsiella pneumoniae* Induced in vitro

Isolate	MIC (µg/mL)				Carbapenemases	MLST	ESBL	AMP _c	Efflux Pump
	IMP	MEM	TGC	CZA					
LY0B1	≥64	≥64	0.5	0.5	KPC-2	ST11		–	/
LY0F1Y	1	4	1	128	KPC-86	ST11	TEM	–	/
LY0B2	≥64	≥64	0.5	0.5	KPC-2	ST11		–	/
LY0F2Y	≥16	≥64	1	128	KPC-44	ST11		–	/
LY0B3	≥64	≥64	0.5	0.5	KPC-2	ST11		–	/
LY0F3Y	8	32	0.5	128	KPC-44	ST11		–	/
10B	≥16	≥64	0.5	≤2	KPC-2	ST11		–	/
10F	≥16	16	1	64	KPC-87	ST11		–	/
207B	≥16	≥64	0.25	4	KPC-2	ST11		–	/
207F	2	2	0.5	64	KPC-88	ST11	TEM	–	↑
14B	≥16	≥64	2	≤2	KPC-2	ST11		–	/
14F	≥16	32	0.5	256	KPC-2	ST11		–	↑

Abbreviations: IMP, imipenem; MEM, meropenem; TGC, tigecycline; CZA, ceftazidime/avibactam; MLST, Multiple-locus sequence typing; ESBL, extended-spectrum beta-lactamase; AmpC, AmpC beta lactamase.

2, and the main mechanism for the induced strains to acquire CZA resistance was mutation of the KPC-2 protein. Among them, KPC-86, KPC-87 and KPC-88 were first discovered by us and have been submitted to PubMed (Accession numbers: MZ067229, MZ067230, MZ067231). The variant KPC-86 was derived from the LY0F1 strain of patient 1, which was completely different from the KPC variant (KPC-14) that formed the CZA-resistant strain (LY0F1-1, LY0F1-2) in vivo. KPC-86 presented the nucleotide A533G mutation, resulting in the substitution of aspartic acid by a glycine at position 179 (D179G). In addition, the mutant KPC-87 of the 10B isolate showed deletion of the nucleotides GCA after position 721, which causes the amino acid substitution of GT241A. Isolate 207B acquired CZA resistance through variant KPC-88, which was associated with a nucleotide substitution at position 523 (G523T). This substitution results in the amino acid change D176Y in the KPC-2 protein. In order to demonstrate the association between CZA resistance and mutated enzymes, the conjugation experiments was performed. The results showed that the receptor bacteria (*E. coli* EC600 strain) developed CZA resistance ([Supplementary Figure 2](#)) and the gene sequencing results showed that there were *KPC* genes (KPC-86, KPC-87, KPC-88) in the corresponding bacteria. The above results further illustrate the diversity of KPC variants in CZA-resistant *Klebsiella pneumoniae*.

To further explore the relationship between the changes in drug susceptibility and the characteristics of the resistant phenotypes for the different KPC mutant types of CZA-resistant strains, we analyzed the results of drug susceptibility tests and detected other drug-resistant phenotypes of the strains. The results showed that the carbapenemase (KPC-2) variants KPC-14, KPC-33, KPC-86, and KPC-88 of the CZA-resistant strains were all sensitive to IMP and MEM and were positive for ESBL ([Table 3](#), [Figure 3](#)). However, the variants KPC-44 and KPC-87 of the CZA-resistant strains showed resistance to IMP and MEM and were negative for ESBL ([Table 3](#)). The above results indicate that the KPC-14, KPC-33, KPC-86 and KPC-88 variants tend to have similar IMP and MEM drug resistance characteristics and resistance phenotypes. The KPC-44 and KPC-87 variants tended to have similar IMP and MEM resistance characteristics and resistance phenotypes.

Discussion

In this study, we isolated CZA-resistant *Klebsiella pneumoniae* with different resistance mechanisms from clinical patients treated with CZA. We also found that CZA-resistant strains with different resistance mechanisms could appear in the same clinical patient sample. In addition, we discovered three new CZA resistance mechanisms of new KPC variants (KPC-86, KPC-87 and KPC-88).

In patient 1, we found carbapenem-resistant *Klebsiella pneumoniae* in ascites, sputum and blood. However, after 10 days of CZA medication, two CZA-resistant strains with different resistance mechanisms were isolated from the same strain of ascites. CZA-resistant strains with different resistance mechanisms also appeared in sputum specimens. Shields et al also reported that three different KPC-3 mutant CZA strains appeared in the same patient.²⁰ The widespread

Table 3 Antimicrobial Susceptibility Changes and the Characteristics of Resistant Phenotypes of Different KPC Mutant Types of CZA Resistant Strains

Carbapenemases	Isolate	Interpretive Categories				ESBL	AMP _c	Efflux Pump
		IMP	MEM	TGC	CZA			
KPC-2	LY0B1	R	R	S	S	–	–	–
KPC-33/86/88	LYOF1 HMOF1 207F	S/I	S/I	S	R	+	–	–
KPC-44/87	LYOF2 LYOF2Y LYOF3Y	R	R	S	R	–	–	–

Abbreviations: IMP, imipenem; MEM, meropenem; TGC, tigecycline; CZA, ceftazidime/avibactam; MLST, Multiple-locus sequence typing; ESBL, extended-spectrum beta-lactamase; AmpC, AmpC beta lactamase.




Figure 3 ESBL confirmation test of the CZA-resistant *Klebsiella pneumoniae* (LY0F1Y and 207F) that were produced in vitro.

Abbreviations: CAZ, ceftazidime; CTX, cefotaxime; CAZ/CLA, ceftazidime/clavulanic acid; CTX/CLA, cefotaxime/ clavulanic acid.

appearance of different resistant strains in the same patient indicates that CZA-resistant mutations are diverse. However, in our research, KPC mutant CZA-resistant strains were isolated from ascites, and strains with upregulated pump protein expression as the mechanism of resistance were also found. These findings fully demonstrate the diversity of drug resistance. In addition, although CZA-resistant *Klebsiella pneumoniae* was isolated in the ascites and sputum of patient 1, the blood isolate did not develop CZA resistance. This result may be related to the highest drug concentration in the blood, which was higher than the mutation concentration, leading to the direct killing of all bacteria. The abdominal cavity, which has a large amount of bacteria, was the primary site of infection. In the abdominal cavity, the drug concentration was lower than that in the blood, and the drug concentration in the lung was lower. Therefore, bacteria are more likely to acquire drug-resistant mutations in the abdominal cavity and sputum.

In this study, we found that the resistance mechanism of CZA-resistant strains isolated from patients was mainly due to KPC variants. The mutants are KPC-14, KPC-44 and KPC-33. The MIC values of IMP and MEM of bacteria with these KPC variants (KPC-14, KPC-44 and KPC-33) were consistent with the results of other research teams.^{21–24} In addition, we were surprised to find that the mechanism of resistance to CZA that was induced in vitro by the bacterial isolates in the ascites specimen in patient 1 was different from the mechanism of resistance of the drug-resistant strains in vivo. After induction of CZA resistance in vitro, a new KPC variant, KPC-86 (D179G), appeared. The MIC values of IMP and MEM of this variant were significantly reduced (Table 2), which may be due to the location of the mutation site on the Ω -loop.²⁵ However, the change in MIC value was not as significant as that of KPC-33 (D179Y),²¹ which also had a mutation at position 179. Compared with the KPC-14 variant (2-amino-acid 242Gly and 243Thr deletion) that occurred in resistant strains in vivo, the MIC values IMP and MEM of KPC-86 were not significantly lower than that of KPC-14.²⁴ Another new KPC variant, KPC-87 (GT242A), appeared in isolate 10B, and the MIC values of IMP and MEM were not significantly reduced. Although the mutation site of KPC-87 was the same as that of KPC-14, the difference in the mutation mode resulted in a large difference in the MIC values of IMP and MEM between KPC-87 and KPC-14 cells. The MIC values of IMP and MEM of another new KPC variant, KPC-88 (D176Y), found in this study were similar to those of KPC-33, which may also be related to the location of the mutation site on the Ω -loop.²⁵ The above results fully indicate that the CZA resistance mechanism of these strains are random, diverse, and uncertain.

Conclusion

In this study, we detected different resistance mechanisms (KPC-14, KPC-44 and KPC-33) of CZA-resistant *Klebsiella pneumoniae* from clinical patients treated with CZA for approximately 10 days. We discovered three new KPC variants (KPC-86, KPC-87 and KPC-88) of these CZA-resistant strains that were induced by CZA in vitro. These findings indicate that the resistance mechanisms of CZA are generally random and diverse and provide a reference basis for the clinical use of CZA. These findings indicate that the resistance mechanisms of CZA are generally random and diverse, and that CZA-resistant strains with different resistance mechanisms may also appear in the same patient. This provides a reference basis for the clinical use of CZA and the selection of antibiotic after CZA resistance. This also emphasizes the importance of preventing the spread of CZA-resistant strains and the importance of drug sensitivity testing.

Ethical Approval

The research was conducted in accordance with the Declaration of Helsinki and according to the recommendations of the Ethics Committee of The Second Affiliated Hospital of Chongqing Medical University. The clinical samples were part of the routine hospital laboratory procedure, the patient data came from the medical record system, and confidentiality was assured throughout the study. The written informed consent has been provided by the patients to have the case details published.

Author Contributions

Min Jiang and Bin Sun share first authorship. All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed to submit to the current journal; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Funding

The work was supported by funds (to Di Mu) from the National Natural Science Foundation of China (82102376) and Kuanren Talents Program of the second affiliated hospital of Chongqing Medical University. This work was supported by funds (to Bin Sun) from the National Natural Science Foundation of China (81802062).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Yahav D, Giske CG, Grāmatniece A, Abodakpi H, Tam VH, Leibovici L. New β -lactam- β -lactamase inhibitor combinations. *Clin Microbiol Rev*. 2020;34(1):e00115–20. doi:10.1128/CMR.00115-20
2. Sharma R, Park TE, Moy S. Ceftazidime-Avibactam: a Novel Cephalosporin/ β -Lactamase Inhibitor Combination for the Treatment of Resistant Gram-negative Organisms. *Clin Ther*. 2016;38(3):431–444. doi:10.1016/j.clinthera.2016.01.018
3. Jorgensen SCJ, Trinh TD, Zasowski EJ, et al. Real-world experience with ceftazidime-avibactam for multidrug-resistant gram-negative bacterial infections. *Open Forum Infect Dis*. 2019;6(12):ofz522. doi:10.1093/ofid/ofz522
4. Strich JR, Ricotta E, Warner S, et al. Pharmacoepidemiology of ceftazidime-avibactam use: a retrospective cohort analysis of 210 US hospitals. *Clin Infect Dis*. 2021;72(4):611–621. doi:10.1093/cid/ciaa061
5. Isler B, Ezure Y, Romero JLG, Harris P, Stewart AG, Paterson DL. Is ceftazidime/avibactam an option for serious infections due to extended-spectrum- β -lactamase- and AmpC-producing *Enterobacterales*? a systematic review and meta-analysis. *Antimicrob Agents Chemother*. 2020;65(1):e01052–20. doi:10.1128/AAC.01052-20
6. Dietl B, Martinez LM, Calbo E, Garau J. Update on the role of ceftazidime-avibactam in the management of carbapenemase-producing *Enterobacterales*. *Future Microbiol*. 2020;15(7):473–484. doi:10.2217/fmb-2020-0012
7. Mirzaie A, Ranjbar R. Antibiotic resistance, virulence-associated genes analysis and molecular typing of *Klebsiella pneumoniae* strains recovered from clinical samples. *AMB Express*. 2021;11(1):122. doi:10.1186/s13568-021-01282-w
8. Ahmadi M, Ranjbar R, Behzadi P, et al. Virulence factors, antibiotic resistance patterns, and molecular types of clinical isolates of *Klebsiella pneumoniae*. *Expert Rev Anti Infect Ther*. 2022;20(3):463–472. doi:10.1080/14787210.2022.1990040
9. Wang G, Zhao G, Chao X, Xie L, Wang H. The characteristic of virulence, biofilm and antibiotic resistance of *Klebsiella pneumoniae*. *Int J Environ Res Public Health*. 2020;17(17):6278. doi:10.3390/ijerph17176278
10. Zhen X, Stålsby Lundborg C, Sun X, Zhu N, Gu S, Dong H. Economic burden of antibiotic resistance in China: a national level estimate for inpatients. *Antimicrob Resist Infect Control*. 2021;10(1):5. doi:10.1186/s13756-020-00872-w

11. Zong Z, Wu A, Hu B. Infection control in the era of antimicrobial resistance in China: progress, challenges, and opportunities. *Clin Infect Dis*. 2020;71(Suppl 4):S372–S378. doi:10.1093/cid/ciaa1514
12. Wang Y, Wang J, Wang R, Cai Y. Resistance to ceftazidime-avibactam and underlying mechanisms. *J Glob Antimicrob Resist*. 2020;22:18–27. doi:10.1016/j.jgar.2019.12.009
13. Karakostas S, Kritsotakis EI, Gikas A. Treatment options for *K. pneumoniae*, *P. aeruginosa* and *A. baumannii* co-resistant to carbapenems, aminoglycosides, polymyxins and tigecycline: an approach based on the mechanisms of resistance to carbapenems. *Infection*. 2020;48(6):835–851. doi:10.1007/s15010-020-01520-6
14. Coppi M, Di Pilato V, Monaco F, Giani T, Conaldi PG, Rossolini GM. Ceftazidime-avibactam resistance associated with increased *bla*_{KPC-3} gene copy number mediated by pKpQIL plasmid derivatives in sequence type 258 *Klebsiella pneumoniae*. *Antimicrob Agents Chemother*. 2020;64(4):e01816–19. doi:10.1128/AAC.01816-19
15. Humphries RM, Ambler J, Mitchell SL, et al. CLSI methods development and standardization working group best practices for evaluation of antimicrobial susceptibility tests. *J Clin Microbiol*. 2018;56(4):e01934–17. doi:10.1128/JCM.01934-17
16. Ding L, Shi Q, Han R, et al. Comparison of four carbapenemase detection methods for *bla*_{KPC-2} variants. *Microbiol Spectr*. 2021;9(3):e0095421. doi:10.1128/Spectrum.00954-21
17. Dallenne C, Da Costa A, Decré D, Favier C, Arlet G. Development of a set of multiplex PCR assays for the detection of genes encoding important beta-lactamases in Enterobacteriaceae. *J Antimicrob Chemother*. 2010;65(3):490–495. doi:10.1093/jac/dkp498
18. Tan D, Zhang Y, Cheng M, et al. Characterization of *Klebsiella pneumoniae* ST11 isolates and their interactions with lytic phages. *Viruses*. 2019;11(11):1080. doi:10.3390/v11111080
19. Nelson K, Hemarajata P, Sun D, et al. Resistance to ceftazidime-avibactam is due to transposition of KPC in a porin-deficient strain of *Klebsiella pneumoniae* with increased efflux activity. *Antimicrob Agents Chemother*. 2017;61(10):e00989–17. doi:10.1128/AAC.00989-17
20. Shields RK, Chen L, Cheng S, et al. Emergence of ceftazidime-avibactam resistance due to plasmid-borne *bla*_{KPC-3} mutations during treatment of carbapenem-resistant *Klebsiella pneumoniae* infections. *Antimicrob Agents Chemother*. 2017;61(3):e02097–16. doi:10.1128/AAC.02097-16
21. Li D, Li K, Dong H, et al. Ceftazidime-avibactam resistance in *Klebsiella pneumoniae* sequence type 11 due to a mutation in plasmid-borne *bla*_{KPC-2} to *bla*_{KPC-33}, in Henan, China. *Infect Drug Resist*. 2021;14:1725–1731. doi:10.2147/IDR.S306095
22. Galani I, Karaikos I, Angelidis E, et al. Emergence of ceftazidime-avibactam resistance through distinct genomic adaptations in KPC-2-producing *Klebsiella pneumoniae* of sequence type 39 during treatment. *Eur J Clin Microbiol Infect Dis*. 2021;40(1):219–224. doi:10.1007/s10096-020-04000-9
23. Bianco G, Boattini M, Iannaccone M, et al. Carbapenemase detection testing in the era of ceftazidime/avibactam-resistant KPC-producing Enterobacteriales: a 2-year experience. *J Glob Antimicrob Resist*. 2021;24:411–414. doi:10.1016/j.jgar.2021.02.008
24. Niu S, Chavda KD, Wei J, et al. A ceftazidime-avibactam-resistant and carbapenem-susceptible *Klebsiella pneumoniae* strain harboring *bla*_{KPC-14} isolated in New York City. *mSphere*. 2020;5(4):e00775–20. doi:10.1128/mSphere.00775-20
25. Winkler ML, Papp-Wallace KM, Bonomo RA. Activity of ceftazidime/avibactam against isogenic strains of *Escherichia coli* containing KPC and SHV β -lactamases with single amino acid substitutions in the Ω -loop. *J Antimicrob Chemother*. 2015;70(8):2279–2286. doi:10.1093/jac/dkv094

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>