



# Current Perspectives on Indications for Ileal Pouch-Anal Anastomosis in Older Patients

Brandon M Shore<sup>1</sup>, Bharati Kochar<sup>2,3</sup>, Hans H Herfarth<sup>4,5</sup> , Edward L Barnes<sup>4,5</sup> 

<sup>1</sup>Department of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA; <sup>2</sup>Division of Gastroenterology, Massachusetts General Hospital, Boston, MA, USA; <sup>3</sup>The Mongan Institute, Boston, MA, USA; <sup>4</sup>Division of Gastroenterology and Hepatology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA; <sup>5</sup>Center for Gastrointestinal Biology and Diseases, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

Correspondence: Edward L Barnes, Division of Gastroenterology and Hepatology, University of North Carolina at Chapel Hill, Campus Box #7080, 130 Mason Farm Road, Chapel Hill, NC, 27599-7080, USA, Tel +1 919-962-3112, Fax +1 919-966-6842, Email [edward\\_barnes@med.unc.edu](mailto:edward_barnes@med.unc.edu)

**Abstract:** The population of older patients with inflammatory bowel disease (IBD) is expected to continue to increase in the coming decades, which necessitates and improved understanding of the critical issues faced by patients in this population. Although restorative proctocolectomy with IPAA remains the surgical procedure of choice for the majority of patients with medically refractory ulcerative colitis (UC) and UC-related dysplasia, the evidence surrounding surgery for older patients UC remains sparse. In particular, comparisons of outcomes among older and younger patients undergoing IPAA and comparisons between older patients undergoing IPAA and those undergoing proctocolectomy with end ileostomy remain an understudied and important issue, as evidence in this area will be used to guide patient-centered surgical choices among older patients who require colectomy for UC. In this narrative review, we review the available literature regarding IPAA for older patients, as well as the pre-, peri-, and postoperative factors that may influence outcomes in this population.

**Keywords:** pouchitis, surgery, older age, ulcerative colitis, J-pouch

## Introduction and Epidemiology

Proctocolectomy with ileal pouch-anal anastomosis (IPAA) is the preferred choice of restorative surgery among patients who require colectomy for medically refractory ulcerative colitis (UC) or UC-related dysplasia. As the prevalence of patients with inflammatory bowel disease (IBD) among elderly individuals increase,<sup>1</sup> it is natural that providers will increasingly face complex and multidisciplinary decision-making regarding the need for and timing of restorative proctocolectomy with IPAA in older adults. However, data among older patients undergoing IPAA is limited. In this narrative review, we will discuss the relevant available literature with respect to IPAA among older individuals, with a particular focus on epidemiology, pre-operative and intra-operative planning, and post-operative outcomes among older patients with IBD.

Up to 15% of patients diagnosed with IBD are over 65 years in age, which is generally considered older than the traditional patient with IBD.<sup>2</sup> Additionally, as treatments for IBD improve and the general population life expectancy increases, the proportion of older patients with IBD is expected to increase in the coming years.<sup>1</sup> Even though restorative proctocolectomy with IPAA remains the surgical procedure of choice for most patients with UC, the evidence surrounding surgery for older patients with severe UC remains an understudied and important issue as most studies have been small in size with limited generalizability. In a systematic review of over 5600 publications, a paucity of studies evaluated rates of colectomy for UC in older patients as well as rates of surgery for all older patients with IBD.<sup>1</sup> A multicenter study in Greece showed patients above the age of 60 with UC were less likely to undergo any surgery for UC compared to younger patients (5.9% vs 18.2%).<sup>3</sup> Another small study showed low rates of total colectomy (less than 2%) in the older population but did not comment specifically on the rate of IPAA.<sup>4</sup> Studies looking at surgery for colorectal cancer found patients with IPAA had a mean age of 44 compared to 54 in those with a permanent ileostomy.<sup>5</sup>

A surgeon's preference to reserve IPAA for younger patients may be a result of the extensive pelvic dissection and concern for more postoperative complications in older patients compared to younger, healthier patients. However, several recent publications have demonstrated favorable post-operative outcomes among surgery when preoperative and intra-operative factors are considered.<sup>6,7</sup> Consideration of the patient's comorbidities and goals can allow the provider to create an individualized approach and select among multiple surgical options for the best long-term outcomes.

## Surgical Options

For older patients (defined as age 60 or older)<sup>8,9</sup> with severe or medically refractory UC, several surgical options exist: proctocolectomy with IPAA, total colectomy with end ileostomy, or ileorectal anastomosis. Currently, none of these options are definitively restricted by age. In many cases, an IPAA is the preferred restorative surgical procedure for patients undergoing colectomy for UC in large part due to reported significant improvements in quality of life after IPAA<sup>10,11</sup> (Table 1). In many cases, a major objective of IPAA surgery is to provide a relief from the burden of underlying disease while also maintaining continence.<sup>12</sup> However, it should be noted that there is a paucity of literature demonstrating the superiority of IPAA over total proctocolectomy with end ileostomy, with one systematic review demonstrating equivalence.<sup>13</sup> IPAA surgery also has short- and long-term risks of adverse events that should be considered including immediate postoperative complications such as leaks, hemorrhage, and pelvic sepsis,<sup>12</sup> as well as long-term complications such as pouchitis and other chronic inflammatory conditions of the pouch.<sup>11,12</sup> Given these considerations, shared decision-making is critical for any patient considering surgical options for UC.

Total colectomy with ileorectal anastomosis (IRA) is a rectal sparing surgery that involves removal of the diseased colon with re-anastomosis to the rectum. IRA surgery does not involve an extensive pelvic dissection, thus the risk of sexual dysfunction and incontinence is lower.<sup>14</sup> Potential benefits of IRA include a perceived decreased difficulty or complexity of surgery compared to IPAA, lower complication rates and a similar long-term functional outcome in selected patients.<sup>14</sup> However, patient selection is important as those with extensive rectal disease should be carefully considered for IPAA over IRA given the risk of rectal dysplasia over time, with studies demonstrating dysplasia rates increasing from 9% at 10 years to 25% at 20 years in IRA.<sup>14,15</sup> Additionally, elderly patients without good rectal compliance or adequate anal sphincter tone may be poor surgical candidates.<sup>14</sup> In evaluations not restricted or analyzed by age at the time of surgery, morbidity and mortality after IRA were found to be as high as 28%<sup>16</sup> and 4.2%,<sup>17</sup> respectively.

On average, patients undergoing IRA record 6 bowel movements per day, minimal nighttime seepage, and 1 nocturnal bowel movement.<sup>15</sup> Despite these favorable immediate outcomes, rates of cumulative failure have been reported at 10.1% and 24.1% at 5 and 10 years, with many needing proctectomy with IPAA or continent ileostomy.<sup>18</sup> These data should be interpreted based on each unique clinical scenario; it may be reasonable for an older patient with many comorbidities and shorter life expectancy to undertake these risks compared to a younger patient. However, in the current practice paradigm, IRA remains a choice for a very selected patient population.

**Table 1** Advantages and Disadvantages of Surgical Approaches for Medically Refractory Ulcerative Colitis

	Ileal Pouch-Anal Anastomosis	Ileorectal Anastomosis	End Ileostomy
<b>Advantages</b>	Restorative surgery Low rate of incontinence Low rate of rectal cancer Favorable quality of life	Restorative surgery Lower rates of sexual dysfunction Lower rates of incontinence Favorable functional outcomes	No risk for rectal cancer if proctectomy is performed
<b>Disadvantages</b>	Motility and pelvic floor issues may affect pouch function Inflammatory conditions of the pouch are common	Higher rates of dysplasia and/or rectal cancer Increased risk for recurrent disease and need for maintenance therapy	Non-restorative surgery Decreased quality of life

## Pre-Operative Considerations in Older vs Younger Patients

Patient comorbidities, frailty, and sphincter function are all integral parts of the surgery planning process and can help guide functional assessments to help risk stratify for post-operative adverse outcomes. The American College of Surgeons recommends the use of the National Surgical Quality Improvement Program (NSQIP) risk calculator to estimate risk of adverse outcomes after surgery.<sup>19,20</sup> Important preoperative factors to assess within the calculator include age, functional status, steroid use, disseminated cancer, previous cardiac event, chronic obstructive pulmonary disease (COPD), dialysis and many more.<sup>20</sup> These factors help predict the presence of nonsurgical complications including morbidity, mortality, pneumonia, and cardiac events and should be evaluated in conjunction with data evaluating post-surgical outcomes.<sup>20</sup> However, there may be existing limitations to using this generalized calculator in the setting of IBD-related surgeries, particularly IPAA. The American College of Surgeons NSQIP-IBD Collaborative is a disease-specific collaborative focusing on variables specific to IBD-related surgery and outcomes, including IPAA.<sup>21</sup> This may have significant advantages for risk stratification and future research efforts.

Frailty is an emerging factor used to help assess risk of postoperative complications<sup>7</sup> and describes a state of reduced physiologic reserve and includes cognition, functionality, nutrition and comorbidities.<sup>22</sup> Many of the factors involved in determining frailty are similar to those described in the NSQIP risk calculator.<sup>23</sup> One study by Telemi et al showed that higher modified frailty indices correlated with increased rates of morbidity and cardiopulmonary complications.<sup>23</sup>

Similarly, pre-operative sarcopenia is an important factor that may help providers determine risk of non-operative complications. Sarcopenia describes loss of skeletal muscle mass over time and is higher in those with chronic medical conditions and malnourishment such as IBD patients.<sup>24</sup> It could be a surrogate for physiologic reserve; sarcopenia is strongly associated with longer and more difficult post-operative recovery.<sup>24</sup> Some studies have measured sarcopenia through visceral adiposity on abdominal Computed Tomography (CT) imaging and shown a direct correlation between lower visceral adiposity and increased rates of the need for rescue therapy in hospitalized IBD patients.<sup>25</sup> In a retrospective study of 69 patients, sarcopenia was associated with an independent risk factor for surgical site infection after restorative proctocolectomy with IPAA.<sup>26</sup> Similarly, among patients undergoing surgery for colorectal cancer, sarcopenia has been associated with an increased risk for complications such as non-surgical site infections and longer hospital stays.<sup>27</sup> However, standardized definitions and measures of both frailty and sarcopenia in a younger population of adults, such as patients with IBD, are not yet standardized. The combination of frailty, sarcopenia and the NSQIP risk calculators have the potential to serve as valuable tools to determine post-operative risk for non-operative complications.

Sphincter tone and pelvic floor health is another important factor that should be addressed pre-operatively. Increasing age has traditionally been associated with an increased risk for incontinence.<sup>28</sup> Studies using anorectal manometry both pre and post IPAA have demonstrated that patients who experience incontinence after IPAA are more likely to have had incontinence prior to IPAA.<sup>29</sup> A caveat for such an approach would be that assessment of sphincter function in a patient with medically refractory UC is not predictive of the sphincter function in the setting of a pouch with no significant cuff inflammation. Given these potential differences in incontinence and underlying sphincter function, some authors have argued that it is important to assess sphincter function prior to consideration of an IPAA, with a subsequent frank discussion of post-operative expectations particularly among elderly patients.<sup>28</sup> The best method of this assessment is not well defined however. Older literature suggests that although a significant decrease in mean resting pressure may be noted on anorectal manometry in some patients after the initial IPAA creation, this was not associated with worse clinical outcomes after ileostomy takedown. Additionally, there was no correlation between preoperative mean resting pressure and postoperative incontinence score.<sup>30</sup> At a minimum, a digital rectal examination should be performed for evaluation of sphincter tone and dysfunction in lieu of using chronologic age alone as a barrier to determine eligibility for IPAA.

## Intra-Operative Considerations in Older vs Younger Patients

In the current era, IPAA is typically performed in a staged approach, with delayed creation of the pouch being preferred.<sup>11</sup> In a multi-staged approach to IPAA, two forms of ostomy may be used. In a traditional 2-stage IPAA, a total proctocolectomy is performed with a loop ileostomy followed by an ostomy takedown. In a 3-stage approach, a total abdominal colectomy is performed with an end ileostomy, followed by proctectomy and IPAA with a diverting loop ileostomy (second stage), and

ultimately an ostomy takedown. Patients who develop high output in their diverting loop ileostomy (approximately 14% of patients) may require early closure of their ostomy, which increases their risk of morbidity and mortality.<sup>31</sup> Older patients with high output stomas are at greater risk for electrolyte imbalances, dehydration and readmission<sup>32</sup> and elderly patients with underlying sarcopenia or frailty may have even higher susceptibility to these issues.

In an evaluation of prospective data collected through the American College of Surgeons NSQIP, delayed pouch creation was associated with a decreased likelihood of unplanned reoperations and major and minor adverse events.<sup>33</sup> Importantly, this data should be taken in consideration of the need for an additional surgery, which may independently offset these adverse events outlined above in the older patient. Additionally, it may be reasonable to pursue IRA in the right patient to preserve continence and avoid the complicated pelvic dissection in IPAA if the risks of dysplasia are explained and understood, especially in those with a life expectancy of less than 10 years. Additionally, total proctocolectomy with end ileostomy may be considered in this population. For those who choose IPAA, data among rates of postoperative complications comparing one- versus two- or three-stage approaches to IPAA in the older population is limited and therefore should be an individualized decision among providers and patients.

## Post Operative Considerations in Older vs Younger Patients

### Immediate Outcomes

Early complications (within 30 days of surgery)<sup>34</sup> most commonly consist of pelvic sepsis, portal vein thrombosis, hemorrhage, pouch ischemia and anastomotic leak.<sup>12</sup> Late complications (>30 days postoperatively)<sup>34</sup> include but are not limited to chronic pelvic sepsis, pouch fistula, pouch dysfunction, pouchitis, small bowel obstruction, malignancy and pouch failure.<sup>12</sup> Pouchitis is the most common complication after IPAA, affecting 40% of patients within the first year of surgery<sup>35</sup> and more than 80% of patients within their lifetime.<sup>36,37</sup> An additional concern is that the incidence of pouchitis appears to be increasing, as evidenced by a recent evaluation of a population-based prospective cohort study from Denmark.<sup>38</sup>

In a meta-analysis studying postoperative complications among elderly patients undergoing IPAA, increasing age at the time of surgery was not associated with an increase in short- or long-term morbidity and mortality.<sup>39</sup> Immediate outcomes were assessed from 13 studies and found that the rate of surgical site infections, pelvic sepsis and ileus requiring nasogastric tube placement were 8.2%, 9.9% and 18.4%, respectively.<sup>39</sup> Around 8% required reoperation for SBO and 16.3% had an unplanned hospital readmission within 30 days.<sup>39</sup> Further analysis comparing those age 50–65 compared to those over 65 showed no significant difference in rates of SBO, anastomotic leaks, reoperation and readmission among these age groups.<sup>39</sup> Another retrospective study pooling the VA national database of elderly patients with UC found no significant difference in surgical morbidity or 30-day mortality comparing patients who underwent IPAA vs ileo-rectal anastomosis.<sup>40</sup> In a more recent analysis of 601 patients over 50 years of age undergoing IPAA for UC, numerically greater rates of pouchitis were seen in patients 70 and older compared to patients age 60–69 and 50–59 at the time of surgery.<sup>41</sup>

### Natural History

The majority of patients undergoing IPAA experience long-term success, with an estimated pouch failure rate of 6% and 9% at 5–10 years and more than 10 years follow-up, respectively.<sup>42</sup> However, the data on surgery in older patients is lacking as most existing literature consists of evaluations of patients under 45 years without significant co-morbidities or functional limitations who are otherwise healthy and fit for surgery.<sup>43</sup> A meta-analysis found no significant difference in long-term outcomes including incontinence, anastomotic leak, pouchitis, SBO, and chronic fistula/sinus tract development when comparing older and younger patients.<sup>39</sup>

In addition to the risk of acute pouchitis, chronic inflammatory conditions of the pouch place an additional burden on patients. Approximately 17% of patients will develop chronic pouchitis<sup>44</sup> with an additional 10% of patients developing Crohn's-like disease of the pouch despite a pre-operative diagnosis of UC.<sup>45</sup> Although the data are heterogenous, there are some indications that older patients undergoing IPAA may be at risk for chronic inflammatory conditions of the pouch including chronic antibiotic dependent pouchitis (CADP).<sup>46</sup> This is of particular concern given that chronic inflammation is thought to be a potential risk factor for the development of pouch dysplasia<sup>47</sup> and pouch failure.<sup>48</sup>

In more granular comparisons of functional outcomes and patient satisfaction, IPAA appears to be very well tolerated among older patients. One meta-analysis found that patients had an average of 6 daily bowel movements and 1.6 nocturnal bowel movements,<sup>39</sup> both of which are consistent with accepted values postoperatively. In additional assessments, 72% of patients above 60 years with IPAA could differentiate flatus from feces.<sup>39</sup> Another review pooled results from four studies and showed a similar number of daily bowel movements in patients 65 years of age compared to those 45 years old.<sup>1</sup> Other studies demonstrated good quality of life scores in individuals up to 70 years and more than 96% of patients over 65 years said they would recommend the surgery to others after 3 years.<sup>49</sup> Several prior studies have indicated that the average rate of incontinence in patients with IPAA may increase over time after IPAA; however, there may be confounding variables contributing to these findings including increasing age, neuropathy and pharmacologic therapies.<sup>39,50</sup> The assessment of both functional outcomes and specifically patient-reported outcomes among older patients undergoing IPAA is critical, as evidenced by the fact that defecation urge and incomplete evacuation are among the most important factors older patients use to determine whether the surgery was successful.<sup>51</sup>

Although pouch failure rates among elderly patients were similar to the general population, prior studies have shown that pouch failure increases with increasing time, indicating that the time from initial IPAA surgery may be a more important prognostic factor for pouch failure compared to age alone.<sup>28,52</sup> The most common causes of pouch failure include pelvic sepsis and anastomotic leak,<sup>39,41</sup> however the rates of these complications do not seem to differ when comparing age at time of surgery.<sup>39</sup> Rates of dysplasia and/or any neoplasia including pouch lymphoma after IPAA remain low, with estimates generated in a large single center study in the US around 5% at 25 years.<sup>53</sup> Patients undergoing proctocolectomy with IPAA for colonic dysplasia or cancer and individuals with chronic pouch inflammation appear to be at the highest risk for dysplasia complications, with annual pouchoscopy suggested in these patients.<sup>54</sup> Importantly, the impact of aging on pouch and rectal cuff neoplasia is unknown, and thus the evidence to support decisions regarding stopping surveillance pouchoscopy is lacking (Table 2).

## Motility-Related Disorders

Pelvic floor dysfunction and other motility disorders are a feared complication after IPAA in older patients given the extensive pelvic dissection performed in the surgery,<sup>55</sup> however literature surrounding this issue is limited. Moreover, some weakening in pelvic floor and anal sphincter muscles are expected with age, which may predispose patients undergoing IPAA to motility disorders.<sup>55</sup> The mechanism for this is thought to be from anal sphincter atrophy and shortening, and decreased sensation that occurs with age.<sup>56</sup> Furthermore, older patients commonly experience some degree of skeletal muscle loss, which also increase their risk of pelvic floor dyssynergia.<sup>56</sup> In patients undergoing IPAA, rates of incontinence may be even higher due to the association of decreased anal sphincter pressures after pouch creation and impaired relaxation during defecation.<sup>56,57</sup> However, this issue is common, and many patients find some relief from biofeedback therapy, an approach that focuses on improving pelvic and anal muscle relaxation during defecation.<sup>58</sup> A retrospective study by Quinn showed that biofeedback therapy provided improvement in symptoms in 91% of IPAA patients, but this study did not focus on the older patient population.<sup>58</sup> Overall, functional outcomes remain a high priority for elderly patients following IPAA, and surgical techniques should be optimized to preserve sphincter function as much as possible.

**Table 2** Current Knowledge Regarding Ileal Pouch-Anal Anastomosis Surgery in Older Patients and Opportunities for Future Research

<b>What is known</b>	Restorative surgery including IPAA is associated with improved quality of life Risks of short-term morbidity and mortality appear similar between older patients and younger patients undergoing IPAA Rates of pouch failure are similar to younger patients undergoing IPAA Underlying motility disorders may impact pouch function and outcomes
<b>What is unknown/future research questions</b>	How long should surveillance pouchoscopies be performed in older patients with increased risk for colorectal cancer (positive family history or previous colorectal Ca)? What is the impact of aging on neoplasia rates after IPAA? How do frailty and sarcopenia impact short and long-term outcomes after IPAA surgery? What patient preferences drive decisions for end ileostomy versus restorative surgery (IPAA)?



## Conclusion

IPAA has been shown to be safe and well tolerated among elderly patients with IBD. Despite the limitations of a narrative review, we identified no major differences in the literature in both short- and long-term outcomes among patients undergoing IPAA over 60 years compared to younger individuals. Additionally, quality of life indices among this patient population indicate favorable outcomes compared to those receiving an end ileostomy. It is important to note that two patients of the same chronologic age can be significantly different surgical candidates when considering both their comorbidities and functional status, and thus pre-operative screening and individualized counseling regarding all surgical options is critical. Overall, the decisions regarding pursuit of an IPAA should not be based upon age alone but rather a holistic picture taking into consideration the patient's wishes, comorbidities, and preoperative functional status as individualized patient selection likely offers the best opportunity for optimized postoperative outcomes.

## Funding

This research was supported by grants from the National Institutes of Health [K23DK127157-01 (ELB)] and [R03AG074059 (BK)].

## Disclosure

Brandon Shore has no relevant disclosures.

Bharati Kochar has served as a consultant for Pfizer.

Hans H. Herfarth has served as a consultant for Alivio, AMAG, BMS, ExeGI Finch, Gilead, Janssen, Lycera, Merck, Otsuka, Pfizer, PureTech, Seres and has received research support from Pfizer and Artizan Biosciences.

Edward L Barnes reports grants from National Institutes of Health, during the conduct of the study; personal fees from AbbVie, personal fees from Gilead, personal fees from Pfizer, personal fees from Target RWE, personal fees from Eli Lilly, outside the submitted work.

The authors report no other conflicts of interest in this work.

## References

- Shung DL, Abraham B, Sellin J, et al. Medical and surgical complications of inflammatory bowel disease in the elderly: a systematic review. *Dig Dis Sci*. 2015;60:1132–1140. doi:10.1007/s10620-014-3462-2
- Ananthakrishnan AN, McGinley EL, Binion DG. Inflammatory bowel disease in the elderly is associated with worse outcomes: a national study of hospitalizations. *Inflamm Bowel Dis*. 2009;15:182–189. doi:10.1002/ibd.20628
- Triantafyllidis JK, Emmanouilidis A, Pomonis E, et al. Ulcerative colitis in the elderly: clinical patterns and outcome in 51 Greek patients. *J Gastroenterol*. 2001;36:312–316. doi:10.1007/s005350170096
- Almogy G, Sachar DB, Bodian CA, et al. Surgery for ulcerative colitis in elderly persons: changes in indications for surgery and outcome over time. *Arch Surg*. 2001;136:1396–1400. doi:10.1001/archsurg.136.12.1396
- Snelgrove R, Brown CJ, O'Connor BI, et al. Proctocolectomy for colorectal cancer--is the ileal pouch anal anastomosis a safe alternative to permanent ileostomy? *Int J Colorectal Dis*. 2014;29:1485–1491. doi:10.1007/s00384-014-2027-3
- Cohan JN, Bacchetti P, Varma MG, et al. Impact of patient age on procedure type for ulcerative colitis: a national study. *Dis Colon Rectum*. 2015;58:769–774. doi:10.1097/DCR.0000000000000398
- Cohan JN, Bacchetti P, Varma MG, et al. Outcomes after ileoanal pouch surgery in frail and older adults. *J Surg Res*. 2015;198:327–333. doi:10.1016/j.jss.2015.04.014
- Pardi DS, Loftus EV Jr, Camilleri M. Treatment of inflammatory bowel disease in the elderly: an update. *Drugs Aging*. 2002;19:355–363. doi:10.2165/00002512-200219050-00004
- Vavricka SR, Rogler G. Treatment of severe ulcerative colitis: differences in elderly patients? *Dig Dis*. 2009;27:315–321. doi:10.1159/000228567
- Barnes EL, Herfarth HH, Kappelman MD, et al. Incidence, risk factors, and outcomes of pouchitis and pouch-related complications in patients with ulcerative colitis. *Clin Gastroenterol Hepatol*. 2021;19:1583–1591 e4. doi:10.1016/j.cgh.2020.06.035
- Barnes EL, Lightner AL, Regueiro M. Peri-operative and post-operative management of patients with crohn's disease and ulcerative colitis. *Clin Gastroenterol Hepatol*. 2020;18:1356–1366. doi:10.1016/j.cgh.2019.09.040
- Ng KS, Gonsalves SJ, Sagar PM. Ileal-anal pouches: a review of its history, indications, and complications. *World J Gastroenterol*. 2019;25:4320–4342. doi:10.3748/wjg.v25.i31.4320
- Murphy PB, Khot Z, Vogt KN, et al. Quality of life after total proctocolectomy with ileostomy or IPAA: a systematic review. *Dis Colon Rectum*. 2015;58:899–908. doi:10.1097/DCR.0000000000000418
- Scoglio D, Ahmed Ali U, Fichera A. Surgical treatment of ulcerative colitis: ileorectal vs ileal pouch-anal anastomosis. *World J Gastroenterol*. 2014;20:13211–13218. doi:10.3748/wjg.v20.i37.13211
- da Luz Moreira A, Kiran RP, Lavery I. Clinical outcomes of ileorectal anastomosis for ulcerative colitis. *Br J Surg*. 2010;97:65–69. doi:10.1002/bjs.6809

16. Borjesson L, Lundstam U, Oresland T, et al. The place for colectomy and ileorectal anastomosis: a valid surgical option for ulcerative colitis? *Tech Coloproctol*. 2006;10:237–241. doi:10.1007/s10151-006-0286-x
17. Oakley JR, Lavery IC, Fazio VW, et al. The fate of the rectal stump after subtotal colectomy for ulcerative colitis. *Dis Colon Rectum*. 1985;28:394–396. doi:10.1007/BF02560219
18. Andersson P, Norblad R, Soderholm JD, et al. Ileorectal anastomosis in comparison with ileal pouch anal anastomosis in reconstructive surgery for ulcerative colitis—a single institution experience. *J Crohn's Colitis*. 2014;8:582–589. doi:10.1016/j.crohns.2013.11.014
19. Liu Y, Cohen ME, Hall BL, et al. Evaluation and enhancement of calibration in the American college of surgeons NSQIP surgical risk calculator. *J Am Coll Surg*. 2016;223:231–239. doi:10.1016/j.jamcollsurg.2016.03.040
20. Bilimoria KY, Liu Y, Paruch JL, et al. Development and evaluation of the universal ACS NSQIP surgical risk calculator: a decision aid and informed consent tool for patients and surgeons. *J Am Coll Surg*. 2013;217:833–42 e1–e3. doi:10.1016/j.jamcollsurg.2013.07.385
21. Eisenstein S, Holubar SD, Hilbert N, et al. The ACS national surgical quality improvement program-inflammatory bowel disease collaborative: design, implementation, and validation of a disease-specific module. *Inflamm Bowel Dis*. 2019;25:1731–1739. doi:10.1093/ibd/izz044
22. Hubbard RE, Story DA. Patient frailty: the elephant in the operating room. *Anaesthesia*. 2014;69:26–34. doi:10.1111/anae.12490
23. Telemi E, Trofymenko O, Venkat R, et al. Frailty predicts morbidity after colectomy for ulcerative colitis. *Am Surg*. 2018;84:225–229. doi:10.1177/000313481808400229
24. Pedersen M, Cromwell J, Nau P. Sarcopenia is a predictor of surgical morbidity in inflammatory bowel disease. *Inflamm Bowel Dis*. 2017;23:1867–1872. doi:10.1097/MIB.0000000000001166
25. Cushing KC, Kordbacheh H, Gee MS, et al. Sarcopenia is a novel predictor of the need for rescue therapy in hospitalized ulcerative colitis patients. *J Crohn's Colitis*. 2018;12:1036–1041. doi:10.1093/ecco-jcc/jjy095
26. Fujikawa H, Araki T, Okita Y, et al. Impact of sarcopenia on surgical site infection after restorative proctocolectomy for ulcerative colitis. *Surg Today*. 2017;47:92–98. doi:10.1007/s00595-016-1357-x
27. Nakanishi R, Oki E, Sasaki S, et al. Sarcopenia is an independent predictor of complications after colorectal cancer surgery. *Surg Today*. 2018;48:151–157. doi:10.1007/s00595-017-1564-0
28. Chang S, Shen B, Remzi F. When not to pouch: important considerations for patient selection for ileal pouch-anal anastomosis. *Gastroenterol Hepatol*. 2017;13:466–475.
29. Dayton MT, Larsen KR. Should older patients undergo ileal pouch-anal anastomosis? *Am J Surg*. 1996;172:444–447. doi:10.1016/S0002-9610(96)00253-X
30. Morgado PJ Jr, Wexner SD, James K, et al. Ileal pouch-anal anastomosis: is preoperative anal manometry predictive of postoperative functional outcome? *Dis Colon Rectum*. 1994;37:224–228. doi:10.1007/BF02048159
31. Gunnarsson U, Karlborn U, Docker M, et al. Proctocolectomy and pelvic pouch—is a diverting stoma dangerous for the patient? *Colorectal Dis*. 2004;6:23–27. doi:10.1111/j.1463-1318.2004.00570.x
32. Arenas Villafranca JJ, Lopez-Rodriguez C, Abiles J, et al. Protocol for the detection and nutritional management of high-output stomas. *Nutr J*. 2015;14:45. doi:10.1186/s12937-015-0034-z
33. Kochar B, Barnes EL, Peery AF, et al. Delayed ileal pouch anal anastomosis has a lower 30-day adverse event rate: analysis from the national surgical quality improvement program. *Inflamm Bowel Dis*. 2018;24:1833–1839. doi:10.1093/ibd/izz082
34. Peyrin-Biroulet L, Germain A, Patel AS, et al. Systematic review: outcomes and post-operative complications following colectomy for ulcerative colitis. *Aliment Pharmacol Ther*. 2016;44:807–816. doi:10.1111/apt.13763
35. Gionchetti P, Calabrese C, Laureti S, et al. Pouchitis: clinical Features, diagnosis, and treatment. *Int J Gen Med*. 2021;14:3871–3879. doi:10.2147/IJGM.S306039
36. Barnes EL, Herfarth HH, Sandler RS, et al. Pouch-related symptoms and quality of life in patients with ileal pouch-anal anastomosis. *Inflamm Bowel Dis*. 2017;23:1218–1224. doi:10.1097/MIB.0000000000001119
37. Lightner AL, Mathis KL, Dozois EJ, et al. Results at up to 30 years after ileal pouch-anal anastomosis for chronic ulcerative colitis. *Inflamm Bowel Dis*. 2017;23:781–790. doi:10.1097/MIB.0000000000001061
38. Barnes EL, Allin KH, Iversen AT, et al. Increasing incidence of pouchitis between 1996 and 2018: a population-based Danish cohort study. *Clin Gastroenterol Hepatol*. 2022. doi:10.1016/j.cgh.2022.04.015
39. Pedersen KE, Jia X, Holubar SD, et al. Ileal pouch-anal anastomosis in the elderly: a systematic review and meta-analysis. *Colorectal Dis*. 2021;23:2062–2074. doi:10.1111/codi.15665
40. Longo WE, Virgo KS, Bahadursingh AN, et al. Patterns of disease and surgical treatment among United States veterans more than 50 years of age with ulcerative colitis. *Am J Surg*. 2003;186:514–518. doi:10.1016/j.amjsurg.2003.07.001
41. Duraes LC, Liang J, Steele SR, et al. Restorative proctocolectomy with ileal pouch-anal anastomosis in elderly patients - is advanced age a contraindication? *ANZ J Surg*. 2022. doi:10.1111/ans.17728
42. Alsafi Z, Snell A, Segal JP. Prevalence of 'pouch failure' of the ileoanal pouch in ulcerative colitis: a systematic review and meta-analysis. *Int J Colorectal Dis*. 2022;37:357–364. doi:10.1007/s00384-021-04067-6
43. Chapman JR, Larson DW, Wolff BG, et al. Ileal pouch-anal anastomosis: does age at the time of surgery affect outcome? *Arch Surg*. 2005;140:534–539. doi:10.1001/archsurg.140.6.534
44. Fazio VW, Kiran RP, Remzi FH, et al. Ileal pouch anal anastomosis: analysis of outcome and quality of life in 3707 patients. *Ann Surg*. 2013;257:679–685. doi:10.1097/SLA.0b013e31827d99a2
45. Barnes EL, Kochar B, Jessup HR, et al. The incidence and definition of crohn's disease of the pouch: a systematic review and meta-analysis. *Inflamm Bowel Dis*. 2019;25:1474–1480. doi:10.1093/ibd/izz005
46. Weaver KN, Kochar B, Hansen JJ, et al. Chronic antibiotic dependent pouchitis is associated with older age at the time of ileal pouch anal anastomosis (j-pouch) surgery. *Crohn's Colitis*. 2019;360:otz029. doi:10.1093/crocol/otz029
47. Ardalan ZS, Sparrow MP. A personalized approach to managing patients with an ileal pouch-anal anastomosis. *Front Med*. 2019;6:337. doi:10.3389/fmed.2019.00337
48. Fazio VW, Ziv Y, Church JM, et al. Ileal pouch-anal anastomoses complications and function in 1005 patients. *Ann Surg*. 1995;222:120–127. doi:10.1097/0000658-199508000-00003

49. Delaney CP, Fazio VW, Remzi FH, et al. Prospective, age-related analysis of surgical results, functional outcome, and quality of life after ileal pouch-anal anastomosis. *Ann Surg.* **2003**;238:221–228. doi:10.1097/01.sla.0000080825.95166.26
50. Takao Y, Gilliland R, Noguera JJ, et al. Is age relevant to functional outcome after restorative proctocolectomy for ulcerative colitis?: prospective assessment of 122 cases. *Ann Surg.* **1998**;227:187–194. doi:10.1097/00000658-199802000-00006
51. Brandsborg S, Chen TY, Nicholls RJ, et al. Difference between patients' and clinicians' perception of pouch dysfunction and its impact on quality of life following restorative proctocolectomy. *Colorectal Dis.* **2015**;17:O136–40. doi:10.1111/codi.12948
52. Mark-Christensen A, Erichsen R, Brandsborg S, et al. Pouch failures following ileal pouch-anal anastomosis for ulcerative colitis. *Colorectal Dis.* **2018**;20:44–52. doi:10.1111/codi.13802
53. Kariv R, Remzi FH, Lian L, et al. Preoperative colorectal neoplasia increases risk for pouch neoplasia in patients with restorative proctocolectomy. *Gastroenterology.* **2010**;139:806–12, 812 e1–2. doi:10.1053/j.gastro.2010.05.085
54. Derikx L, Nissen LHC, Smits LJ, et al. Risk of neoplasia after colectomy in patients with inflammatory bowel disease: a systematic review and meta-analysis. *Clin Gastroenterol Hepatol.* **2016**;14:798–806 e20. doi:10.1016/j.cgh.2015.08.042
55. Farouk R, Pemberton JH, Wolff BG, et al. Functional outcomes after ileal pouch-anal anastomosis for chronic ulcerative colitis. *Ann Surg.* **2000**;231:919–926. doi:10.1097/00000658-200006000-00017
56. Yu SW, Rao SS. Anorectal physiology and pathophysiology in the elderly. *Clin Geriatr Med.* **2014**;30:95–106. doi:10.1016/j.cger.2013.10.003
57. Khanna R, Li Y, Schroeder T, et al. Manometric evaluation of evacuatory difficulty (dyschezia) in ileal pouch patients. *Inflamm Bowel Dis.* **2013**;19:569–575. doi:10.1097/MIB.0b013e31827e78d6
58. Quinn KP, Tse CS, Lightner AL, et al. Nonrelaxing pelvic floor dysfunction is an underestimated complication of ileal pouch-anal anastomosis. *Clin Gastroenterol Hepatol.* **2017**;15:1242–1247. doi:10.1016/j.cgh.2017.02.024

## Clinical and Experimental Gastroenterology

Dovepress

### Publish your work in this journal

Clinical and Experimental Gastroenterology is an international, peer-reviewed, open access, online journal publishing original research, reports, editorials, reviews and commentaries on all aspects of gastroenterology in the clinic and laboratory. This journal is indexed on American Chemical Society's Chemical Abstracts Service (CAS). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-and-experimental-gastroenterology-journal>