


Prospective, Randomized, Fellow Eye-Controlled Study of Postoperative Pain and Inflammation Control with an Intracanalicular Dexamethasone 0.4 mg Ophthalmic Insert Following Small Incision Lenticule Extraction

Kathleen J Jee, Joseph Ling, Shamik Bafna, Thomas Chester , Jeffrey Augustine, William F Wiley

Cleveland Eye Clinic, Brecksville, OH, USA

Correspondence: Kathleen J Jee, Cleveland Eye Clinic, 7001 South Edgerton Road, Suite B, Brecksville, OH, 44141, USA, Tel +1 440-526-1974, Email kathy.jee@gmail.com

Purpose: To compare postoperative anterior chamber inflammation, pain, and patient preference following small incision lenticule extraction (SMILE) in eyes treated with a dexamethasone 0.4 mg intracanalicular insert (DEX) or topical prednisolone acetate (PRED).

Patients and Methods: In this prospective, randomized, fellow eye-controlled trial, 20 patients underwent same-day, bilateral SMILE. One randomly-selected eye of each patient received DEX placed immediately postoperatively, and the fellow eye received topical PRED tapered over 2 weeks. Postoperative evaluations were performed on day 1, week 1, month 1, and month 3. Primary outcomes included postoperative pain, incidence of anterior chamber cell and flare, and patient preference of steroid therapy.

Results: No eyes in either group had any clinically evident cell or flare at any postoperative time point. Mean pain scores (0–10 by subjective report) and incidence of any pain were statistically similar at all postoperative visits. Uncorrected distance visual acuity improved in all eyes, 91% of which achieved 20/25 or better. No eyes lost any lines of corrected distance visual acuity. Three eyes developed a steroid-related rise in intraocular pressure, all of which resolved with 2 of the 3 eyes requiring topical therapy. At 1 week, 1 month, and 3 months, 70%, 65%, and 53% of patients preferred DEX over PRED therapy, respectively.

Conclusion: The DEX insert was preferred by more patients and controlled postoperative inflammation and pain comparably to topical PRED in eyes undergoing SMILE. There were no statistically significant differences in visual outcomes between the two groups.

Keywords: refractive surgery, steroid, myopia, astigmatism

Introduction

Small incision lenticule extraction (SMILE) is a corneal refractive procedure performed with a femtosecond laser in which an intrastromal lenticule is dissected from its anterior and posterior stromal interfaces and removed through a small incision without the need for the creation of a flap.¹ The procedure was approved in the United States in 2016 for the correction of myopia and in 2018 for myopic astigmatism, and has been shown to be effective in correcting hyperopia and presbyopia as well.^{2–4} Meta-analyses suggest SMILE results in less reduction in corneal sensitivity and less incidence of postoperative dry eye compared to femtosecond-laser assisted in situ keratomileusis (FS-LASIK), supporting its preferential use in eyes with pre-existing dryness.^{5–8} The biomechanical strength of the cornea following SMILE is thought to be better preserved compared to FS-LASIK, potentially improving the safety of SMILE in patients with high refractive errors, those who engage in contact sports, or serve in active-duty military.^{8,9}

Following SMILE, the standard postoperative medication regimen includes topical antibiotic and corticosteroid eye drops multiple times per day from multiple bottles with a steroid taper. Recently, there has been a trend for cataract surgeons to move towards a drop-free postoperative experience to improve patient satisfaction and medication compliance following phacoemulsification.^{10–13} A similar shift away from frequent administration of topical therapies following keratorefractive surgery would be expected to improve patients' postoperative experience.

The dexamethasone (DEX) intracanalicular insert (Dextenza, Ocular Therapeutix) is a rod-shaped insert with 0.4 mg of preservative-free dexamethasone embedded within a hydrogel matrix. The DEX insert can be placed into either the superior or inferior punctum and delivers a tapering dose of steroid to the ocular surface for 30 days before dissolving without the need for removal. Phase 3 trials demonstrated the insert's ability to control postoperative pain and inflammation following phacoemulsification cataract surgery with high patient satisfaction.^{14–16}

This study compares postoperative pain, inflammation, and corticosteroid preference in patients undergoing bilateral SMILE treated with the DEX insert in one eye and topical prednisolone acetate in the fellow eye.

Patients and Methods

This prospective, randomized, fellow-eye controlled study was conducted at a single site (Clear Choice Custom LASIK Center) in 20 patients (40 eyes) with three surgeons between June 2020 and February 2021 and was performed according to guidelines established by the Declaration of Helsinki. The study protocol was approved by Salus IRB (Austin, TX), and the study was registered at ClinicalTrials.gov (NCT04380857). Written informed consent was obtained from all study participants. Patient data was anonymized and kept confidential.

Study participants were at least 18 years old and appropriate surgical candidates for SMILE as assessed by refraction and corneal tomography (Pentacam, OCULUS GmbH, Germany). Key exclusion criteria included active ocular or systemic infectious conditions, nasolacrimal duct obstruction, hypersensitivity to dexamethasone, and use of any immunosuppressant or immunomodulating therapies. Consecutive qualifying subjects were invited to participate to minimize selection bias.

Eligible subjects underwent same-day, bilateral SMILE performed with the VisuMax femtosecond laser (Carl Zeiss Meditec, Inc). One eye was randomly assigned to the DEX insert, which was placed in the inferior canaliculus immediately after the SMILE procedure, and the fellow eye was assigned to topical 1% prednisolone acetate (PRED) eye drops prescribed every 2 hours on the day of surgery, 4 times daily for the first week, two times daily for the second week, and then discontinued. All eyes received standard postoperative eye drops including 0.3% ofloxacin (every 2 hours on the day of surgery and then 4 times daily for the first week) and preservative-free artificial tears as needed. Rescue therapy with topical corticosteroids was permitted if anterior chamber cell scores were 2+ or higher, flare scores 3+ or higher, or pain was graded at level 4 or higher. Patients were evaluated 1 day, 1 week, 1 month, and 3 months postoperatively.

The primary outcomes were between-group (DEX versus PRED) differences in the incidence of anterior chamber cell and flare and ocular pain scores, and patient preference for postoperative steroid therapy. Anterior chamber cell and flare were graded using the Standardization of Uveitis Nomenclature Working Group's clinical quantification scale for cell and flare.¹⁷ Ocular pain was assessed subjectively on a 0–10 scale, with 0 being no pain and 10 being worst pain imaginable. Patient preference for steroid treatment was assessed using a 5-point scale consisting of moderately or much preferring one treatment over the other versus no preference.

Secondary outcomes were uncorrected distance visual acuity (UDVA) and corrected distance visual acuity (CDVA) at 1 week, 1 month, and 3 months postoperatively using the ETDRS chart at 4 meters, and the incidence of unscheduled patient encounters (via phone or in-person office visits). Additional data collected included intraocular pressure (IOP) using Goldmann tonometry, surgeon-rated ease of DEX insertion (easy/average/difficult), and post-insertion DEX visualization. No formal power analysis was conducted, and the sample size was set at 40 eyes based on the sample size used in several similar prior studies.^{18–20} Chi-square tests were used to compare proportions and *t*-tests to compare means. The level of significance was set at 0.05.

Results

Twenty-three patients were enrolled in the study. One patient failed screening and was not randomized. Two additional patients were randomized, but the DEX insert could not be successfully placed in the assigned eye at the time of surgery. Analysis was conducted on the remaining 20 patients (40 eyes). Seventeen (85%) patients were White, 11 (55%) were women, and 9 (45%) were men. The mean age was 29.9 years (range: 20 to 37 years). The DEX insert was visualized in all 20 eyes after placement. Surgeon-rated DEX insertion was graded as easy in 12 eyes, average in 3 eyes, and difficult in 5 eyes. All subjects were seen through week 1; three were lost to follow up thereafter.

No eyes had evidence of inflammation with anterior chamber cell or flare at any time point, and no eyes required corticosteroid rescue therapy. Mean pain scores were consistently low (<1 on a 0–10 scale) and were statistically similar between DEX and PRED eyes preoperatively and at each postoperative timepoint (Figure 1). Preoperatively, two patients reported mild bilateral pain and one reported mild unilateral pain in the eye randomized to PRED. The DEX insert was preferred over topical PRED in 70% of patients (14/20) at 1 week, 64.7% of patients (11/17) at 1 month, and 52.9% (9/17) at 3 months; 10%, 17.6%, and 17.6%, respectively, preferred PRED over DEX, and 20%, 17.6%, and 29.4%, respectively, had no preference (Figure 2).

UDVA and CDVA are detailed in Table 1. At 1 week postoperatively, mean logMAR UDVA was 0.013 ± 0.183 and -0.010 ± 0.0114 in PRED and DEX eyes, respectively, which was statistically similar ($P = 0.505$). At 1 month postoperatively, the PRED and DEX eyes had mean UDVA of -0.026 ± 0.160 and -0.014 ± 0.181 , respectively ($P = 0.762$). At 3 months postoperatively, the PRED and DEX eyes had mean UDVA of -0.068 ± 0.156 and -0.071 ± 0.108 , respectively ($P = 0.941$) (Figure 3). The postoperative mean UDVA over time for both groups is shown in Figure 4. No patient lost any lines of CDVA (Figure 5). At 1 week postoperatively, mean CDVA was -0.071 ± 0.101 and -0.083 ± 0.070 in PRED and DEX eyes, respectively ($P = 0.332$). At 1 month postoperatively, the PRED and DEX eyes had mean CDVA of -0.114 ± 0.079 and -0.112 ± 0.087 , respectively ($P = 0.869$). At 3 months postoperatively, the PRED and DEX eyes had mean CDVA of -0.139 ± 0.091 and -0.121 ± 0.078 , respectively ($P = 0.211$). Mean manifest refraction spherical equivalent (MRSE) did not statistically differ between PRED and DEX eyes at 1 week, 1 month, or 3 months ($P = 0.821$, 0.482 , and 0.369 , respectively). At 3 months, 88.2% of eyes in both the PRED and DEX groups were within ± 0.5 D of intended target (Figure 6).

No intraoperative adverse events or complications were recorded. Two subjects experienced bilateral dry eye, one of which resolved within 1 month. Mean IOP was similar between DEX and PRED eyes at 1 week ($P = 0.241$), higher in




Figure 1 Mean pain scores (0–10) postoperatively after small incision lenticule extraction by treatment group.




Figure 2 Patient preference for corticosteroid treatment type at month 3 (n=17).

DEX eyes than PRED eyes at 1 month (consistent with PRED cessation at 2 weeks) ($P = 0.006$), and similar again at 3 months ($P = 0.601$) (Figure 7). Three DEX eyes of 3 subjects experienced elevated IOP at the 1-month visit presumed related to steroid response. Two were moderate (peak IOP 25 and 26 mmHg) and one was more significant, with IOP peak of 45 mmHg and corneal stromal haze with reduced visual acuity. Two of the eyes received a brief course of topical IOP-lowering therapy; IOP in all three eyes normalized within 1 week. These adverse events resulted in a total of 9 unscheduled visits.

Discussion

In this randomized clinical trial, the DEX insert controlled postoperative pain and anterior chamber inflammation comparably to topical PRED in paired fellow eyes undergoing SMILE. At 3 months, 53% of patients preferred the DEX insert compared to only 18% who preferred topical PRED. Both treatments were safe and effective; adverse events were uncommon and not sight-threatening.

Clinically significant inflammation is uncommon after keratorefractive surgery but can adversely affect both visual outcomes and patient satisfaction. LASIK causes a transient increase in anterior chamber cell and/or flare on day 1 that

Table 1 Mean UDVA and CDVA at Postoperative Week 1, Postoperative Month 1, and Postoperative Month 3

	DEX	PRED	P
Postoperative week 1 (n=20)			
Mean logMAR UDVA	-0.010 ± 0.0114	0.013 ± 0.183	0.505
Mean logMAR CDVA	-0.083 ± 0.070	-0.071 ± 0.101	0.332
Postoperative month 1 (n=17)			
Mean logMAR UDVA	-0.014 ± 0.181	-0.026 ± 0.160	0.762
Mean logMAR CDVA	-0.112 ± 0.087	-0.114 ± 0.079	0.869
Postoperative month 3 (n=17)			
Mean logMAR UDVA	-0.071 ± 0.108	-0.068 ± 0.156	0.941
Mean logMAR CDVA	-0.121 ± 0.078	-0.139 ± 0.091	0.211




Figure 3 Uncorrected distance visual acuity (UDVA) three months postoperatively following small incision lenticule extraction in the (A) Dextenza (n=17) and (B) Drops (n=17) groups.




Figure 4 Mean uncorrected distance visual acuity (UDVA) postoperatively over time.

normalizes by day 7.^{21–23} Inflammation is generally less common and less severe after SMILE than FS-LASIK in animal and ex-vivo human studies.^{24,25} Diffuse lamellar keratitis (DLK) has a low reported incidence (0.45% to 1.6%) after SMILE.^{26,27} Recognizing the adverse effects of postoperative inflammation, the American Academy of Ophthalmology's Preferred Practice Pattern for Refractive Errors and Refractive Surgery includes topical corticosteroids among the standard postoperative care following keratorefractive surgery.²⁸ In the current study, no eyes had clinically evident anterior chamber cell or flare, or evidence of DLK, in either treatment group.

Pain occasionally occurs after SMILE, although it is typically mild in nature. In one study of 53 eyes, 21% of eyes manifested mild pain within the first 24 hours post-procedure; no moderate or severe pain was reported, and no new reports of pain arose more than 4 hours into the first 24-hour postoperative period.²⁹ In the current study, half of the eyes in each group reported mild-moderate pain in the first 24 hours, half of which resolved by 1 week, and all of which resolved by month 3. The etiology of the pain in eyes in our study cannot be easily determined. Postoperative pain following keratorefractive surgery can be related to the incision, postoperative inflammation, or a well-described increase




Figure 5 Change in Snellen lines of corrected distance visual acuity (CDVA) three months postoperatively following small incision lenticule extraction in the (A) Dextenza and (B) Drops groups.




Figure 6 Manifest spherical equivalent refractive accuracy three months postoperatively following small incision lenticule extraction in the (A) Dextenza and (B) Drops groups.

in the prevalence and severity of dry eye symptoms. Dry eye symptoms are a significant contributor to patient dissatisfaction with keratorefractive surgery, accounting for 20–30% in published series of dissatisfied patients.^{30,31} A meta-analysis found that SMILE is less likely to cause or aggravate dry eye symptoms compared to FS-LASIK, presumably attributable to avoidance of flap creation and attendant corneal denervation.^{6,32}

Visual outcomes were comparable between treatment groups in this study and were excellent as evidenced at 3 months (Figures 3, 5, 6, 8, 9, and 10). No eyes lost any lines of CDVA. At 3 months, 82.4% of eyes in both groups achieved UDVA of 20/20 or better with 88.2% of eyes with MRSE within ± 0.5 D. These outcomes are consistent with prior studies of SMILE.^{33,34}

IOP elevations occurred in 3 eyes at 1 month and were presumed related to steroid response. Two eyes were treated with topical IOP-lowering therapy that quickly restored IOP control, while the third eye's IOP normalized without therapy. In phase 3 studies of DEX for control of postoperative inflammation and pain following cataract surgery, IOP elevations were reported in 4.4–7.4% of dexamethasone eyes.^{14,15} Steroid responses may be more common in myopic




Figure 7 Mean intraocular pressure (mmHg) postoperatively over time by treatment group.




Figure 8 Attempted versus achieved manifest spherical equivalent refraction three months postoperatively following small incision lenticule extraction in the (A) Dextenza and (B) Drops groups.




Figure 9 Refractive astigmatism three months postoperatively following small incision lenticule extraction in the (A) Dextenza and (B) Drops groups.




Figure 10 Stability of spherical equivalent refraction after small incision lenticule extraction in the (A) Dextenza and (B) Drops groups.

eyes.³⁵ A statistically significant difference in mean IOP between groups was seen at 1 month, but this was attributable to a reduction in mean IOP in the PRED group and not a rise in mean IOP in the DEX group. Future applications of the DEX insert in conjunction with SMILE could decrease the dose to reduce steroid response.

This study demonstrates that postoperative inflammation and pain following SMILE can be controlled without the need for patient-administered topical corticosteroid eye drop therapy. The DEX insert can eliminate the risk of nonadherence with topical steroid therapy. Half of patients administer fewer than half of their postoperative antibiotic and steroid eye drops after cataract surgery.³⁶ Even though SMILE patients are still responsible for administering topical antibiotics, there is a benefit to simplifying the postoperative medical regimen by eliminating the steroid taper. Regimen complexity is associated with poorer adherence, which increases the risk of inflammation, pain, delayed visual recovery, and patient dissatisfaction.^{37–39} Minimizing the frequency of eye drop instillation also reduces hand-face contact, an important consideration in the COVID-19 era, particularly given that about 30% of patients report not washing hands when instilling postoperative eye drops after anterior segment surgery.⁴⁰

The key strength of this study is its prospective, randomized design. Assessments and outcomes were standardized, and all subjects experienced both treatments, serving as their own controls and giving them the ideal opportunity to identify their preferred treatment. Limitations included the relatively small sample size, although our sample size was consistent with similar studies in the recent literature.^{18–20} In addition, validated tests or questionnaires were not used for analysis of ocular pain and steroid preference, which were primary endpoints. Because validated tests were not used, full understanding of the implications of statistical tests is difficult. Moreover, the possibility of cross effects of a drug in both eyes should be considered; however, effects are expected to be minimal for drugs with low penetration into the blood-brain barrier and low systemic absorption.

In summary, the DEX insert controls postoperative inflammation and pain comparably to topical PRED in eyes undergoing SMILE, with similar visual outcomes achieved in both groups and with more patients preferring the DEX insert (53%) over PRED (18%). The DEX insert may reduce the postoperative medication burden, which may improve adherence and reduce the risk of adverse outcomes.

Data Sharing Statement

The data generated and analyzed for this study is not publicly available but may be requested from the corresponding author, Kathleen Jee.

Ethics Approval and Informed Consent

The study protocol was approved by Salus IRB (Austin, TX), and the study was registered at ClinicalTrials.gov (NCT04380857). Written informed consent was obtained from all study participants.

Funding

Ocular Therapeutix provided a research grant that funded this study but were not involved in the study design or submission of the paper for publication.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Reinstein DZ, Archer TJ, Gobbe M. Small incision lenticule extraction (SMILE) history, fundamentals of a new refractive surgery technique and clinical outcomes. *Eye Vis*. 2014;1(1):3. doi:10.1186/s40662-014-0003-1
2. U. S. Food and Drug Administration. Premarket Approval (PMA): VisuMax femtosecond laser. Available from: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P150040>. Accessed July 6, 2021.
3. Moshirfar M, Bruner CD, Skanchy DF, Shah T. Hyperopic small-incision lenticule extraction. *Curr Opin Ophthalmol*. 2019;30(4):229–235. doi:10.1097/ICU.0000000000000580
4. Luft N, Siedlecki J, Sekundo W, et al. Small incision lenticule extraction (SMILE) monovision for presbyopia correction. *Eur J Ophthalmol*. 2018;28(3):287–293. doi:10.5301/ejo.5001069
5. He M, Huang W, Zhong X. Central corneal sensitivity after small incision lenticule extraction versus femtosecond laser-assisted LASIK for myopia: a meta-analysis of comparative studies. *BMC Ophthalmol*. 2015;15:141. doi:10.1186/s12886-015-0129-5
6. Kobashi H, Kamiya K, Shimizu K. Dry eye after small incision lenticule extraction and femtosecond laser-assisted LASIK: meta-analysis. *Cornea*. 2017;36(1):85–91. doi:10.1097/ICO.0000000000000999
7. Zhang Y, Shen Q, Jia Y, Zhou D, Zhou J. Clinical outcomes of SMILE and FS-LASIK used to treat myopia: a meta-analysis. *J Refract Surg*. 2016;32(4):256–265. doi:10.3928/1081597X-20151111-06
8. Titiyal JS, Kaur M, Shaikh F, Gagrani M, Brar AS, Rathi A. Small incision lenticule extraction (SMILE) techniques: patient selection and perspectives. *Clin Ophthalmol*. 2018;12:1685–1699. doi:10.2147/OPTH.S157172
9. Sia RK, Ryan DS, Beydoun H, et al. Visual outcomes after SMILE from the first-year experience at a U.S. military refractive surgery center and comparison with PRK and LASIK outcomes. *J Cataract Refract Surg*. 2020;46(7):995–1002. doi:10.1097/j.jcrs.0000000000000203
10. Shorstein NH, Myers WG. Drop-free approaches for cataract surgery. *Curr Opin Ophthalmol*. 2020;31(1):67–73. doi:10.1097/ICU.0000000000000625
11. Lindstrom RL, Galloway MS, Grzybowski A, Liegner JT. Dropless cataract surgery: an overview. *Curr Pharm Des*. 2017;23(4):558–564. doi:10.2174/1381612822666161129150628
12. Assil KK, Greenwood MD, Gibson A, Vantipalli S, Metzinger JL, Goldstein MH. Dropless cataract surgery: modernizing perioperative medical therapy to improve outcomes and patient satisfaction. *Curr Opin Ophthalmol*. 2021;32(Suppl 1):S1–S12. doi:10.1097/ICU.0000000000000708
13. Bardoloi N, Sarkar S, Paliania A, Das H. Efficacy and safety of dropless cataract surgery. *Indian J Ophthalmol*. 2020;68(6):1081–1085. doi:10.4103/ijo.IJO_1186_19
14. Walters T, Bafna S, Vold S, et al. Efficacy and safety of sustained release dexamethasone for the treatment of ocular pain and inflammation after cataract surgery: results from two phase 3 studies. *J Clin Exp Ophthalmol*. 2016;7:1–11. doi:10.4172/2155-9570.1000572
15. Tyson SL, Bafna S, Gira JP, et al. Multicenter randomized phase 3 study of a sustained-release intracanalicular dexamethasone insert for treatment of ocular inflammation and pain after cataract surgery. *J Cataract Refract Surg*. 2019;45(2):204–212. doi:10.1016/j.jcrs.2018.09.023
16. Gira JP, Sampson R, Silverstein SM, Walters TR, Metzinger JL, Talamo JH. Evaluating the patient experience after implantation of a 0.4 mg sustained release dexamethasone intracanalicular insert (Dextenza): results of a qualitative survey. *Patient Prefer Adherence*. 2017;11:487–494. doi:10.2147/PPA.S126283
17. Jabs DA, Nussenblatt RB, Rosenbaum JT. Standardization of uveitis nomenclature for reporting clinical data. results of the first international workshop. *Am J Ophthalmol*. 2005;140(3):509–516.
18. Greenwood MD, Gorham RA, Boever KR. A randomized fellow-eye clinical trial to evaluate patient preference for dexamethasone intracanalicular insert or topical prednisolone acetate for control of postoperative symptoms following bilateral femtosecond Laser in Site Keratomileusis (LASIK). *Clin Ophthalmol*. 2020;14:2223–2228. doi:10.2147/OPTH.S265311
19. Larsen J, Whitt T, Parker B, Swan R. A randomized, controlled, prospective study of the effectiveness and safety of an intracanalicular dexamethasone ophthalmic insert (0.4 mg) for the treatment of post-operative inflammation in patients undergoing Refractive Lens Exchange (RLE). *Clin Ophthalmol*. 2021;15:2211–2217. doi:10.2147/OPTH.S311070
20. Ibach MJ, Shafer BM, Wallin DD, Puls-Boever KR, Thompson VM, Berdahl JP. The effectiveness and safety of dextenza 0.4 mg for the treatment of postoperative inflammation and pain in patients after photorefractive keratectomy: the RESTORE trial. *J Refract Surg*. 2021;37(9):590–594. doi:10.3928/1081597X-20210610-05
21. El-Harazi SM, Chuang AZ, Yee RW. Assessment of anterior chamber flare and cells after laser in situ keratomileusis. *J Cataract Refract Surg*. 2001;27(5):693–696. doi:10.1016/S0886-3350(01)00798-2
22. Pisella PJ, Albou-Ganem C, Bourges JL, Debbasch C, Limon S. Evaluation of anterior chamber inflammation after corneal refractive surgery. *Cornea*. 1999;18(3):302–305. doi:10.1097/00003226-199905000-00011
23. Vita RC, Campos M, Belfort R Jr, Paiva ER. Alterations in blood-aqueous barrier after corneal refractive surgery. *Cornea*. 1998;17(2):158–162. doi:10.1097/00003226-199803000-00007
24. Luft N, Schumann RG, Dirisamer M, et al. Wound healing, inflammation, and corneal ultrastructure after SMILE and femtosecond laser-assisted LASIK: a human ex vivo study. *J Refract Surg*. 2018;34(6):393–399. doi:10.3928/1081597X-20180425-02
25. Liu L, Cheng W, Wu D, et al. The differential expression of cytokines and growth factors after SMILE compared with FS-LASIK in rabbits. *Invest Ophthalmol Vis Sci*. 2020;61(5):55. doi:10.1167/iovs.61.5.55

26. Reinstein DZ, Stuart AJ, Vida RS, Archer TJ, Carp GI. Incidence and outcomes of sterile multifocal inflammatory keratitis and diffuse lamellar keratitis after SMILE. *J Refract Surg.* **2018**;34(11):751–759. doi:10.3928/1081597X-20181001-02
27. Zhao J, He L, Yao P, et al. Diffuse lamellar keratitis after small-incision lenticule extraction. *J Cataract Refract Surg.* **2015**;41(2):400–407. doi:10.1016/j.jcrs.2014.05.041
28. Chuck RS, Jacobs DS, Lee JK, et al. *Refractive Errors and Refractive Surgery Preferred Practice Pattern.* San Francisco, CA: American Academy of Ophthalmology; **2017**.
29. Liu T, Dan T, Luo Y. Small incision lenticule extraction for correction of myopia and myopic astigmatism: first 24-hour outcomes. *J Ophthalmol.* **2017**;2017:5824534. doi:10.1155/2017/5824534
30. Levinson BA, Rapuano CJ, Cohen EJ, Hammersmith KM, Ayres BD, Laibson PR. Referrals to the Wills Eye Institute cornea service after laser in situ keratomileusis: reasons for patient dissatisfaction. *J Cataract Refract Surg.* **2008**;34(1):32–39.
31. Jabbur NS, Sakatani K, O'Brien TP. Survey of complications and recommendations for management in dissatisfied patients seeking a consultation after refractive surgery. *J Cataract Refract Surg.* **2004**;30(9):1867–1874. doi:10.1016/j.jcrs.2004.01.020
32. Liu YC, Jung ASJ, Chin JY, Yang LWY, Mehta JS. Cross-sectional study on corneal denervation in contralateral eyes following SMILE versus LASIK. *J Refract Surg.* **2020**;36(10):653–660. doi:10.3928/1081597X-20200730-01
33. Han T, Xu Y, Han X, et al. Three-year outcomes of small incision lenticule extraction (SMILE) and femtosecond laser-assisted laser in situ keratomileusis (FS-LASIK) for myopia and myopic astigmatism. *Br J Ophthalmol.* **2019**;103(4):565–568. doi:10.1136/bjophthalmol-2018-312140
34. Ang M, Farook M, Htoon HM, Mehta JS. Randomized clinical trial comparing femtosecond LASIK and small-incision lenticule extraction. *Ophthalmology.* **2020**;127(6):724–730. doi:10.1016/j.ophtha.2019.09.006
35. Chang DF, Tan JJ, Tripodis Y. Risk factors for steroid response among cataract patients. *J Cataract Refract Surg.* **2011**;37(4):675–681. doi:10.1016/j.jcrs.2010.10.051
36. Hermann MM, Ustundag C, Diestelhorst M. Electronic compliance monitoring of topical treatment after ophthalmic surgery. *Int Ophthalmol.* **2010**;30(4):385–390. doi:10.1007/s10792-010-9362-3
37. Tsai JC. A comprehensive perspective on patient adherence to topical glaucoma therapy. *Ophthalmology.* **2009**;116(11 Suppl):S30–S36. doi:10.1016/j.ophtha.2009.06.024
38. Sleath B, Carpenter DM, Blalock SJ, et al. Applying the resources and supports in self-management framework to examine ophthalmologist-patient communication and glaucoma medication adherence. *Health Educ Res.* **2015**;30(5):693–705. doi:10.1093/her/cyv034
39. Newman-Casey PA, Robin AL, Blachley T, et al. The most common barriers to glaucoma medication adherence: a cross-sectional survey. *Ophthalmology.* **2015**;122(7):1308–1316. doi:10.1016/j.ophtha.2015.03.026
40. An JA, Kasner O, Samek DA, Levesque V. Evaluation of eyedrop administration by inexperienced patients after cataract surgery. *J Cataract Refract Surg.* **2014**;40(11):1857–1861. doi:10.1016/j.jcrs.2014.02.037

Clinical Ophthalmology

Dovepress

Publish your work in this journal

Clinical Ophthalmology is an international, peer-reviewed journal covering all subspecialties within ophthalmology. Key topics include: Optometry; Visual science; Pharmacology and drug therapy in eye diseases; Basic Sciences; Primary and Secondary eye care; Patient Safety and Quality of Care Improvements. This journal is indexed on PubMed Central and CAS, and is the official journal of The Society of Clinical Ophthalmology (SCO). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-ophthalmology-journal>