

MTHFR and MTRR Genetic Polymorphism of Methotrexate Therapy Outcomes in Early Rheumatoid Arthritis

Qian Zhang^{1,2}, Pan Fu², Zhanglei Cao², Hua Huang³, Qinwen Wen³, Kaizhe Wang², Tong Kong², Xiudi Wu³, Jianping Zheng²

¹Jiangxi Key Laboratory for Rare Earths Magnetic Materials and Devices, College of Rare Earths, Jiangxi University of Science and Technology, Ganzhou, 341000, People's Republic of China; ²Cixi Institute of Biomedical Engineering, Ningbo Institute of Materials Technology & Engineering, Chinese Academy of Sciences (CAS), Ningbo, 315300, People's Republic of China; ³Department of Rheumatology and Immunology, Ningbo First Hospital, Ningbo, 315010, People's Republic of China

Correspondence: Jianping Zheng; Xiudi Wu, Tel +86-18091984088; +86-13857826442, Email zhengjianping@nimte.ac.cn; awu1100@aliyun.com

Purpose: Methotrexate (MTX) is used as an anchor drug for the treatment of rheumatoid arthritis (RA) and there may be differences in drug action between genotypes. The purpose of this study was to investigate the relationship between clinical efficacy response and disease activity of MTX monotherapy with methylenetetrahydrofolate reductase (MTHFR) and methionine synthase reductase (MTRR) polymorphisms.

Patients and Methods: In the study, a population of 32 patients in East China with early RA fulfilling the diagnostic standards of the American College of Rheumatology (ACR) were enrolled, all of them received MTX monotherapy. Genotyping of patients MTHFR C677T and A1298C, MTRR A66G using tetra-primer ARMS-PCR method and sanger sequencing to verify its accuracy.

Results: The distribution of three polymorphic genotypes that were studied is in accordance with the Hardy-Weinberg genetic equilibrium. The patient pathology variables smoke (OR = 0.088, P = 0.037), drink alcohol (OR = 0.039, P = 0.016) and males (OR = 0.088, P = 0.037) were significantly associated with non-response to MTX. Genotype, allele distribution and genetic statistical models were not found to be related to MTX treatment response and disease activity in both the response groups and non-response groups.

Conclusion: Our findings suggest that the MTHFR C677T, MTHFR A1298C and MTRR A66G polymorphisms may not predict MTX clinical treatment response and disease activity in patients with early RA. The study revealed that smoke, alcohol, and males were possible influential factors for MTX non-response.

Keywords: MTHFR, MTRR, methotrexate, rheumatoid arthritis, DAS28

Introduction

Rheumatoid arthritis (RA) is a chronic systemic autoimmune disease, mainly characterised by symmetrical pain and swelling of small joints in the hands and the feet.¹ When chronic inflammation is not properly controlled over a long period of time, it can eventually lead to irreversible damage to the joint structures. Methotrexate (MTX) is a folic acid inhibitor with structural and physicochemical properties similar to folic acid, which acts as an anti-inflammatory and anti-proliferative agent in the body.² MTX is used extensively for the treatment of RA because of its efficacy, safety profile and affordability. The European League Against Rheumatism (EULAR) recommended MTX monotherapy for patients with early RA, and MTX in combination with other disease-modifying antirheumatic drugs (DMARDs) for patients with moderate or severe disease activity.³ However, 30–50% of patients with RA still have poor response to MTX treatment or relapse with inadequate response to re-treatment, resulting in drug resistance, and have to stop treatment or change to other medicines.⁴ The personalized response between patients may be due to differences in the gene expression or activity in the folic acid-MTX metabolic pathway, and such differences may arise to alter drug pharmacokinetics and affect MTX response.⁵

The methylenetetrahydrofolate reductase (MTHFR) and methionine synthase reductase (MTRR) are closely related to the folic acid metabolic pathway. MTHFR catalyzes the transformation of 5,10-methylenetetrahydrofolate (5,10-CH₂-THF) into 5-methyltetrahydrofolate (5-MTHF), which provides methionine-methylation for homocysteine (Hcy).⁶ MTRR generates functionally active methionine synthase (MTR), and allows it to catalyse the conversion of Hcy to methionine, maintaining normal levels of Hcy.⁷ These two enzymes are significant regulation factors, each regulated by its separate gene, and their associated regulatory genes are essential for folic acid metabolism and have therefore been extensively studied.

The MTHFR gene is located on chromosome 1p36.3 and the encoded production is a key enzyme in the metabolic process of the body. In existing studies, multiple polymorphic sites in the MTHFR gene have been reported, with the most widely studied SNP sites being mainly C677T (rs1801133) and A1298C (rs1801131). The early studies showed that the C677T mutation causes a decrease in enzyme activity, with carriers of the 677TT genotype and 677CT genotype having 30% and 65% of the enzyme activity of the 677CC genotype, respectively.⁸ Ashfield-Watt et al reported that the C677T mutation resulted in reduced levels of ingested folic acid, and that 677TT had lower levels of folic acid than the 677CT or 677CC genotypes, requiring higher intake to achieve equilibrium Hcy concentrations.⁹ Whether the enzyme activity was altered by the A1298C mutation remains controversial.^{10,11} Neither homozygote nor heterozygote of A1298C resulted in increased or decreased in folic acid concentrations.¹² MTRR gene mutations are one of the main causes of Hcy and abnormal folic acid metabolism, with A66G (rs1801394) as the most prominent and most studied mutation site. MTRR gene is located on chromosome 5p15.2–15.3, and A66G mutation results in methionine substitution at the 22nd isoleucine. The ability to metabolize folic acid and folic acid deficiency were associated with MTRR gene polymorphisms.¹³ MTRR A66G mutation leads to abnormal plasma hcy levels, with significantly higher hcy concentrations in the 66AG and 66GG genotypes than in the 66AA genotype, which may have an impact on MTX treatment response.¹⁴ In exploring the relationship between MTX therapeutic response with MTHFR C677T and MTHFR A1298C polymorphisms, it produced remarkably heterogeneous results, making this influential relationship full of uncertainty.

Therefore, in this study, we investigated a potential correlation between clinical efficacy response and disease activity of MTX monotherapy and MTHFR C677T, MTHFR A1298C and MTRR A66G polymorphisms in RA patients in East China for the first time, hoping to provide a possibility to achieve personalized treatment for RA patients.

Materials and Methods

Workflow Chart

Figure 1 demonstrates the general design of this study.




Figure 1 Workflow chart of the relationship between clinical efficacy response and disease activity of methotrexate (MTX) monotherapy with MTHFR C677T, MTHFR A1298C and MTRR A66G SNPs polymorphisms.

Characteristics of the Population

During 2019–2021, we diagnosed 32 patients with RA in the Department of Rheumatology and Immunology of the Ningbo First Hospital according to the 1987 diagnostic standards of the ACR,¹⁵ and all patients were diagnosed with RA for the first time. The early patients included in this study were treated with methotrexate monotherapy (no other DMARDs in combination) and initial methotrexate treatment dose was $7.5\text{mg} \pm 2.5\text{mg/week}$. To relieve the methotrexate negative effects, patients were supplemented with folic acid 10mg/week during treatment. The average treatment period was three months, according to the DAS28 score method and the standards for evaluating the degree of rheumatoid arthritis disease of ACR.¹⁶ Quantitative indicators to assess patients at baseline and post-treatment, including tender joints counts (TJC), swollen joints counts (SJC), erythrocyte sedimentation rate (ESR) and pain visual analogue scales (VAS). Based on EULAR's response criteria according to DAS28,¹⁷ MTX treatment response was determined by a combination of the DAS28 difference between the baseline and post-treatment periods, the ΔDAS28 , and the DAS28 at post-treatment period, with $\Delta\text{DAS28} \leq 0.6$ and $\Delta\text{DAS28} > 0.6 \leq 1.2$ but $\text{DAS28} > 5.1$ defined as treatment ineffective and the rest as moderate or good response. In this study, we divided patients into response and non-response groups according to treatment response, with moderate or good effect patients in the response group and ineffective patients in the non-response group. To explore the correlation between genotypes and response to MTX treatment by comparing the genotype distribution of patients in the response and non-response groups. The study was approved by the local ethics committee, complied with the standard procedures of the Declaration of Helsinki, and informed consent was obtained from all subjects.

DNA Extraction

Whole blood samples were taken intravenously from patients with EDTA anticoagulated blood collection tubes and marked with the date and patient information. Genomic DNA was extracted according to the kit manufacturer's recommendations (DNeasy Blood&Tissue Kit, QIAGEN, Germany). DNA was quantified using NanoDrop 2000 (Thermo Fisher Scientific, Massachusetts, US) and Qubit 2.0 (Thermo Fisher Scientific, Massachusetts, US). DNA integrity was tested by agarose gel electrophoresis. Purified gDNA samples were stored at -80°C .

Genotype Identification

Identification of allelic genotypes for the MTHFR C677T, MTHFR A1298C and MTRR A66G polymorphisms by primers designed based on the tetra-primer amplification refractory mutation system PCR (ARMS-PCR) proposed by Ye et al.¹⁸ The 3' terminus of the inner primer was designed according to the mutation site, and allele-specific bases were introduced, as well as a mismatched base at the penultimate third of the 3' terminus to enhance the specificity of the PCR amplification. The primers sequences and working concentrations, and annealing temperatures are shown in [Table S1](#). Both outer primer and inner primer concentrations were optimized for the system, and the amplification of the two shorter allele-specific products was enhanced at a final concentration ratio of 1:4, with clear genotyping results. The $10\text{ }\mu\text{L}$ PCR reaction system includes 10ng DNA, outer primer final concentration of $0.1\text{--}0.2\text{ }\mu\text{M}$, inner primer final concentration of $0.4\text{--}0.8\text{ }\mu\text{M}$, and 0.5U Taq polymerase (AmpliTaq Gold™ 360 DNA polymerase, Thermo Fisher Scientific, Massachusetts, US). The PCR amplification procedure was an initial denaturation at 95°C for 5 minutes, cycle conditions of denaturation at 95°C for 30 seconds, annealing temperature for 30 seconds depending on the polymorphic site, extension at 72°C for 30 seconds and final extension at 72°C for 7 minutes. Mixed $4\text{ }\mu\text{L}$ of PCR products with $2\text{ }\mu\text{L}$ of loading dye and analyzed genotypes in 8% polyacrylamide gel electrophoresis (PAGE).

Meta-Analysis

This section of the study was searched for literature using PubMed and Web of Science databases. Searches were conducted using the combination of keywords: “methotrexate” AND “rheumatoid arthritis” AND “polymorphism”. The title and abstract were used to obtain information related to the study for literature screening. The criteria for inclusion in the literature were: (1) study of the association of MTHFR C677T, MTHFR A1298C and MTRR A66G with MTX treatment outcomes in RA patients; (2) literature publication date in the last decade; (3) use of a case-

control or cohort design and inclusion of detailed genotype data, clinical characteristics of patients and baseline drugs in the literature.

Statistical Analysis

Statistical analysis of data was performed with IBM SPSS Statistics for Windows, version 26 (IBM Corp., Armonk, NY, USA). The chi-square test assessed whether the genotype distribution was consistent with Hardy-Weinberg equilibrium (HWE), $P > 0.05$ indicated genetic equilibrium and the sample was statistically significant. Patients' sample measurement data were described with the means and standard deviations (\pm SD) or interquartile ranges (IQRs). Results were compared between two groups using the Student's *t*-test and between multiple groups (or three different SNPs analyzed in the same sample) with Kruskal–Wallis test. When the frequency was less than 5, we used Fisher's exact test instead of the chi-square test. *P* values were considered statistically significant at $p < 0.05$. Genotype and allele frequencies were compared based on chi-square tests, and allelic and genotypic risks were assessed with used odds ratios (ORs) and 95% confidence intervals (CIs).

Meta-analysis was conducted using STATA 17.0 (Stata Corporation, TX, USA) software. The Mantel-Haenszel test was used to statistically analyze the genetic models such as dominant model, recessive model and co-dominant model to determine the correlation between MTHFR C677T, MTHFR A1298C and MTRR A66G polymorphisms and MTX treatment efficacy. Risk estimates were expressed using OR and 95% CIs, and Z-test *P* values < 0.05 were considered statistically significant.

Results

Patient Genotype

To determine the patient genotype, we designed outer and inner primers for the C677T, A1298C and A66G sites, respectively. In the tetra-primer ARMS-PCR method, optimization of the inner and outer primer concentrations was essential for accurate genotyping. To further determine the optimal primer concentration, we used PCR amplification with different ratios of outer primers and inner primers of 1:1, 1:2, 1:4 and 1:10, respectively, and found that at the primer ratio of 1:4, the PCR products obtained by 8% PAGE resulted in clear separation of each genotype (Figure 2A–C) (size of PCR amplification products refer to Table S1). The genotype identification results of 32 patients with regard to C677T, A1298C and A66G are shown in Table S2. To verify the genotyping accuracy, PCR amplification was done using outer primers for each SNP site and sanger sequencing was performed on all samples. From further comparison, we found that the patient genotype obtained by the tetra-primer amplification method was 100% consistent with the sanger sequencing results (Figure 2D–F). Therefore, we identified the genotype of the patient and provided a basis for subsequent correlation analysis.

Clinical Features of the Patients

A total of 32 patients with early RA were reported in this study, and the clinical characteristics of the patients at baseline and the variables assessed by DAS28 post-treatment period are listed in Table 1. All patients were treated with MTX monotherapy from the first date of diagnosis of RA, with a mean treatment period of 86 days. Based on Δ DAS28, 21 of the 32 patients were determined to be in the response group and 11 in the non-response group. Our results revealed no significant differences in age, BMI, Anti-CCP positivity, RF positivity, MTX treatment measures, or treatment duration between the response and non-response groups ($p > 0.05$). In contrast, smoke (current smokers at the time of data collection, OR = 0.088, $P = 0.037$), drink (alcohol consumption ≥ 1 unit, OR = 0.039, $P = 0.016$) and male (OR = 0.088, $P = 0.037$) were significantly associated with non-response to MTX. The average levels of TJC, SJC, ESR and DAS28 after the post-treatment in patients were 1 (0–2), 0 (0–1), 19.5 (10–33) mm/h and 3.12 ± 1.55 , respectively, which were clearly different in the response and non-response groups, but not statistically significant ($P > 0.05$).




Figure 2 Genotyping results of patients with MTHFR C677T, A1298C and MTRR A66G and validation of PCR product sanger sequencing. (A): The SNP of MTHFR C677T (rs1801133). M: Markers: 20bp; lanes 1, 3, 4, 5, 7, 9 heterozygotes alleles (CT); lanes 2, 6, 7 mutant homozygotes alleles (TT); (B): The SNP of MTHFR A1298C (rs1801131). M: Markers: 20bp; lanes 1, 2, 4, 6, 8 wild type homozygotes (AA); lanes 3, 5, 7, 9 heterozygotes alleles (AC); (C): The SNP of MTRR CA66G (rs1801394). M: Markers: 20bp; lanes 1, 2, 3, 4, 8, 9 wild type homozygotes (AA); lanes 3, 6, 7 heterozygotes alleles (AG). (D): Sanger sequencing chromatogram of patient samples in the MTHFR C677T polymorphism showing three genotypes (CC, CT, TT). (E): Direct sequencing chromatogram of patient samples in the MTHFR A1298C polymorphism showing two genotypes (AA, AC). (F): Direct sequencing chromatogram of patient samples in the MTRR A66G polymorphism showing two genotypes (AA, AG).

Association of MTX Clinical Response with C677T, A1298C and A66G Polymorphisms

Table 2 shows the distribution of MTHFR C677T, MTHFR A1298C and MTRR A66G polymorphism genotypes between the response and non-response groups and comparison of statistical models. The results indicated that the

Table 1 Clinical Characteristics Variables of the RA Patients Studied

Characteristics	RA Patients (n=32)	Responders (n=21)	Non-Responders (n=11)	OR (95% CI)	P-value
Patients-related					
Gender (Female/Male)	27/5	20/1	7/4	0.088 (0.008–0.921)	0.037
Age (years) (mean ± SD)	50.4 ± 13.28	49.81 ± 12.6	51.55 ± 15.06		0.748
BMI median (IQR) kg/m ²	21.9 (17.6–28.3)	21.83 (20.0–23.1)	22.04 (19.4–24.7)		0.329
Height (cm) (mean ± SD)	162.1 ± 5.85	160.37 ± 4.55	164.64 ± 6.96		0.089
Weight (kg) (mean ± SD)	57.4 ± 8.56	55.11 ± 6.09	61.45 ± 10.84		0.1
Current smoking, n (%)	5 (15.6)	1 (4.8)	4 (36.36)	0.088 (0.008–0.921)	0.037
Alcohol consumption (≥1 unit per week), n (%)	4 (12.5)	0 (0)	4 (36.36)	0.039 (0.002–0.808)	0.016
Disease-related					
Anti-CCP positivity, n (%)	29 (90.63)	19 (90.48)	10 (90.91)	0.950 (0.076–11.803)	0.968
RF positivity, n (%)	30 (93.75)	20 (95.24)	10 (90.91)	2.000 (0.113–35.411)	0.636
Treatment related					
MTX dose (mg/week) (mean ± SD)	9.6 ± 0.92	9.52 ± 1.0	9.77 ± 0.75		0.297
Treatment duration, median (IQR), days	86 (82–91)	86 (84–91)	84 (73–106)		0.504
Folic acid dose (mg/week) (mean ± SD)	8.9 ± 2.1	8.57 ± 2.31	9.55 ± 1.51		0.163
Individual variable of DAS28 (post-treatment)					
TJC (out of 28), median (IQR)	1 (0–2)	0 (0–1)	2 (1–4)		0.158
SJC (out of 28), median (IQR)	0 (0–1)	0 (0–1)	2 (1–3)		0.130
ESR, median (IQR)	19.5 (10–33)	19 (9–28)	23 (10–67)		0.330
DAS 28 (mean ± SD)	3.12 ± 1.55	2.21 ± 0.70	3.93 ± 1.6		0.061

Notes: P value < 0.05 is considered to be of statistical significance (bolded display).

Abbreviations: n, number; SD, Standard deviation; IQR, interquartile range; BMI, body mass index; Anti-CCP, anti-cyclic citrullinated peptide antibodies; RF, rheumatoid factor; TJC, tender joints counts; SJC, swollen joints counts; ESR, erythrocyte sedimentation rate; DAS28, Disease Activity Score in 28 joints.

Table 2 Distribution of Polymorphic Genotypes, Alleles and Genetic Statistical Models in Response and Non-Response Groups

	Responders n (%)	Non-Responders n (%)	P-value	Compared Model	P-value	OR (95% CI)
MTHFR C677T	n = 21	n = 11				
Genotypes						
CC	8 (38.10)	3 (27.27)	Ref.	CC vs CT+TT	0.703	1.641 (0.334–8.068)
CT	9 (42.86)	6 (54.55)	0.402	CT vs CC+TT	0.530	0.625 (0.144–2.713)
TT	4 (19.05)	2 (18.18)	0.605	TT vs CC+CT	1	1.059 (0.162–6.938)
Allele						
C	25 (59.52)	12 (54.55)	Ref.			
T	17 (40.48)	10 (45.45)	0.702			
MTHFR A1298C						
AA	13 (61.90)	9 (81.82)	Ref.	AA vs AC+CC	0.425	0.361 (0.062–2.114)
AC	8 (38.10)	2 (18.18)	0.229	AC vs AA+CC	0.425	2.769 (0.473–16.213)
Allele						
A	34 (80.95)	20 (90.90)	Ref.			
C	8 (19.05)	2 (9.09)	0.254			
MTRR A66G						
AA	13 (61.90)	8 (72.73)	Ref.	AA vs AG+GG	0.703	0.609 (0.124–2.996)
AG	7 (33.33)	3 (27.27)	0.490	AG vs AA+GG	1	1.333 (0.267–6.653)
GG	1 (4.76)	0 (0)	0.636	GG vs AA+AG	1	1.683 (0.063–44.772)
Allele						
A	33 (78.57)	19 (86.36)	Ref.			
G	9 (21.43)	3 (13.64)	0.521			

Notes: Fisher's exact test is used when the frequency is less than 5; Ref. as reference category.

Abbreviations: n, number; OR, odds ratio; CI, confidence interval.

genotypes, alleles and the distribution of the three statistical models (recessive model, dominant model and co-dominant model) at each SNP site were not significantly correlated with MTX clinical treatment response.

Association of Disease Activity with C677T, A1298C and A66G Polymorphisms

To determine the relationship between different genotypes and disease activity in patients, the distribution between MTHFR C677T, MTHFR A1298C and MTRR A66G polymorphisms genotypes and disease activity parameters were compared, as shown in Table 3. The results showed that ESR levels were not remarkably different across the C677T

Table 3 Disease Activity Parameters Related to Polymorphic Genotypes

	ESR		TJC		SJC		DAS28	
	Mean ± SD	K-W	Median (IQR)	K-W	Median (IQR)	K-W	Mean ± SD	K-W
MTHFR C677T genotype								
CC	26.45 ± 24.65	0.302	1 (0–2)	0.867	1 (0–1)	0.868	3.03 ± 1.23	0.975
CT	26.73 ± 20.20	0.302	1 (0–2)	0.867	0 (0–1)	0.868	3.24 ± 1.50	0.975
TT	22.17 ± 14.88	0.302	1 (0–5)	0.867	0 (0–2)	0.868	2.98 ± 1.03	0.975
MTHFR A1298C genotype								
AA	30.41 ± 21.95	0.056	1 (0–2)	1	0 (0–2)	0.571	3.35 ± 1.42	0.186
AC	15.60 ± 12.34	0.056	1.5 (0–2)	1	0 (0–1)	0.571	2.61 ± 0.66	0.186
MTRR A66G genotype								
AA	26.52 ± 20.80	0.766	1 (0–2)	0.666	0 (0–1)	0.591	3.10 ± 1.36	0.883
AG	25.50 ± 21.51	0.766	1 (0–2)	0.666	0.5 (0–1)	0.591	3.14 ± 1.22	0.883
GG	13	0.766	2	0.666	1	0.591	3.26	0.883

Notes: K-W, Kruskal–Wallis, K-W test for comparison between multiple groups.

Abbreviations: SD, Standard deviation; IQR, interquartile range; ESR, erythrocyte sedimentation rate; TJC, tender joints counts; SJC, swollen joints counts; DAS28, Disease Activity Score in 28 joints.

genotypes and were reduced in the 1298AC and 66GG genotypes ($P > 0.05$). DAS28 was decreased after the post-treatment in 677TT and 1298AC, but was not statistically significant. TJC and SJC were not found to be significantly different across genotypes in the three SNPs. To further compare the differences in each disease activity parameter between the response and non-response groups at baseline and after the post-treatment, [Figure S1](#) shows that ESR, TJC, SJC and DAS28 were remarkably lower in the response group ($p < 0.001$), while the differences were not statistically significant in the non-response group.

Meta-Analysis

A total of 373 publications were identified after the initial search, and 37 publications were included after analysis of titles and abstracts. The full literature was interpreted in detail, and 13 of these publications met our inclusion criteria, with reasons for exclusion: (1) no detailed data for genotype; (2) letter or comment.

MTHFR C677T: The meta-analysis of MTHFR C677T included 12 studies,^{19–30} which included 958 responders and 840 non-responders, and the main characteristic information is presented in [Table 4](#). For all samples, dominant model (OR = 1.095, 95% CI = 0.81–1.362, $P = 0.412$) ([Figure 3](#)), recessive model (OR = 0.701, 95% CI = 0.81–1.362, $P = 0.087$) ([Figure S2A](#)) and codominant model (OR = 1.059, 95% CI = 0.844–1.329, $P = 0.621$) ([Figure S2B](#)) did not observe an association between the C677T polymorphism and MTX treatment response, nor was significant heterogeneity observed between studies. Study cohort stratified by ethnicity found that C677T polymorphism was significantly associated with MTX treatment response in the recessive model (OR = 0.497, 95% CI = 0.279–0.886, $P = 0.018$) and codominant model (OR = 1.492, 95% CI = 1.046–2.126, $P = 0.027$) in the European population, no association was found in other models and populations ([Table 5](#)).

MTHFR A1298C: The meta-analysis of MTHFR A1298C included 9 studies,^{20–24,26–29} which included 688 responders and 615 non-responders, and the main characteristic information is presented in [Table 6](#). For all samples, dominant model (OR = 1.023, 95% CI = 0.730–1.435, $P = 0.894$) ([Figure S3A](#)), recessive model (OR = 0.784, 95% CI = 0.390–1.578, $P = 0.495$) ([Figure S3B](#)) and codominant model (OR = 1.075, 95% CI = 0.760–1.520, $P = 0.682$) ([Figure 4](#)) did not observe an association between the A1298C polymorphism and MTX treatment response. The study cohort was stratified by ethnicity, and there was a significant association between the A1298C polymorphism and MTX treatment response in the recessive model (OR = 0.432, 95% CI = 0.201–0.926, $P = 0.031$) and the codominant model (OR = 1.981, 95% CI = 0.1108–3.543, $P = 0.021$) in the South Asian population. In addition, there was significant between study heterogeneity in the recessive model ($I^2=55\%$, $P = 0.023$) ([Table 5](#)).

MTRR A66G: The meta-analysis of MTRR A66G included 3 studies,^{21,28,31} which included 376 responders and 338 non-responders, and the main characteristic information is presented in [Table 7](#). For all samples, dominant model (OR = 0.879, 95% CI = 0.619–1.249, $P = 0.473$) ([Figure S4A](#)), recessive model (OR = 0.910, 95% CI = 0.594–1.392, $P = 0.663$) ([Figure S4B](#)) and codominant model (OR = 1.177, 95% CI = 0.850–1.629, $P = 0.327$) ([Figure 5](#)) did not observe an association between the A1298C polymorphism and MTX treatment response. No significant heterogeneity between studies was observed in any of the three genetic models ([Table 5](#)).

Discussion

Since 1988, when the US Food and Drug Administration approved MTX as therapy for RA,³² till the EULAR recognized it to be the anchor drug for the treatment of RA,³ MTX has played an important role in the treatment of RA. Low-dose MTX has long been considered an effective and safe anti-rheumatic drug, as well as a base drug for other DMARDs combinations.³³ However, because of individual differences in clinical response to MTX, at least one-third of patients with RA have no response or significantly lower efficacy when they take it.³⁴ Similar differences in clinical response were observed in our study cohort of the Zhejiang, East China population, with 34.4% of patients with early RA poorly treated with MTX monotherapy. In this study, we explored the correlations between MTX therapy response and disease activity with MTHFR C677T, MTHFR A1298C and MTRR A66G polymorphisms.

Our patient clinical pathology characteristic statistics demonstrated that age and BMI were not relevant to clinical response to MTX, while smoke status, drink and male may be factors associated with non-response to MTX. In early

Table 4 Summary of the Analyzed Studies and the Distribution of MTHFR C677T Genotypes

Study	Year	Ethnicity	Genotyping Method	Responders			Nonresponders			Mean Age, Years	Mean Disease Duration (Years)	Sex (Female/Male) (%)	MTX Dose (mg/Week)	Therapy Method	Date of End Point (Week)
				CC	CT	TT	CC	CT	TT						
Soukup T et al ²⁶	2015	Caucasian	Taqman	36	36	8	16	21	3	58.5 ± 12.6	No information	26.7/73.3	11 ± 2.7	Mixed	24
Ghodke-Puranik Y et al ²⁸	2015	South Asian	PCR RFLP	38	10	1	128	39	1	43.8 ± 10.4	5.6 ± 4.9	86/14	7.5–20	MTX	48
Boughrara W et al ²²	2017	Africa	Taqman	24	36	5	17	27	1	48.8 ± 13.44	9.28 ± 8.936	82.3/17.7	12.67 ± 2.33	MTX	24
Lima A et al ³⁰	2014	Caucasian	PCR RFLP	52	46	7	53	53	22	52 ± 11.9	8.0 (0.5–53.0)	84.1/15.9	15.0 (2.5–25.0)	MTX	24
Salazar J et al ²⁹	2014	Europe	Sanger sequence	21	17	1	9	6	1	55.62 ± 1.297	5.55	81.5/18.5	7.5–25	Mixed	24
Lv S et al ²¹	2018	East Asian	Sanger sequence	19	51	28	10	39	14	52.99 ± 13.81	4	82.4/17.6	8.92 ± 2.26	MTX	12
Berkani LM et al ²³	2017	Africa	Taqman	14	19	8	7	3	3	44.26 ± 14.41	<2	87.04/12.96	15.05 ± 2.4	Mixed	24
Lima A et al ²⁴	2016	Europe	Sequenom iPLEX	52	46	7	53	53	22	51 ± 15.6	0.3–51	84.1/15.9	2.5–25	MTX	12
Wang S et al ²⁰	2020	East Asian	PCR RFLP	102	62	12	58	48	14	54.6 ± 11.6	5.93	70.3/29.7	13.89 ± 1.99	MTX	24
Uribarri M et al ²⁵	2015	Europe	Taqman	13	28	7	5	8	7	61.5 ± 13.2	13.91 ± 8.11	72.1/27.9	No information	MTX	No information
Kolan SS et al ¹⁹	2022	Europe	Sequenom iPLEX	53	57	14	45	22	9	53.95	0.23–0.83	61/39	15	MTX	16
Iqbal MP et al ²⁷	2015	South Asian	PCR-RFLP	19	7	2	16	5	2	42.87 ± 13.5	6.2–7.0	86.6/13.4	15–25	MTX	24

Notes: Mean age was measured using mean±SD; Mean disease duration and MTX dose were measured using mean±SD or range.




Figure 3 Forest plot of the correlation between MTHFR C677T polymorphism and the efficacy of MTX in RA patients (CC vs CT+TT (dominant model)).

Abbreviations: OR, odds ratio; Cis, confidence interval.

studies, pathological variables were used as measures of MTX response prediction models, but with varying results. Among patient-related factors, males were connected with a good response to MTX treatment.³⁵ Smoking, alcohol, and high BMI may be risk factors for non-response to MTX in RA patients.³⁶ Whereas in our study, males had a poorer response to treatment. In addition, BMI was at normal levels in both the MTX response and non-response patient groups. Of the disease-related factors, the proportion of Anti-CCP positivity and RF positivity was similar in both groups initially.

Our study and meta-analysis displayed that genetic polymorphisms in MTHFR C677T and MTHFR A1298C were not associated with clinical efficacy response of MTX, and appeared consistent with our results in many studies.^{20,37} In Western Algerian population studies, it was concluded that C677T and A1298C cannot predict clinical response and adverse drug reactions to MTX treatment outcomes.²² Research on 120 patients with RA in the East Bohemian population has not found any correlation between the genotypes C677T and A1298C and ineffectiveness of MTX treatment in the respective genetic statistical models.²⁶ In an early study of the efficacy of MTX with folic acid supplementation, the C677T polymorphism was found to have failed to predicted toxicity or efficacy response to MTX treatment in patients with RA who received folic acid supplementation. At the same time, the

Table 5 Association of MTHFR C677T, MTHFR A1298C and MTRR A66G Gene Polymorphisms with the Outcome of MTX in RA Patients

Genetic Models	Number of Studies	OR (95% CI)	P-value (OR)	I ² (%)	P-value (H)
MTHFR C677T					
CC vs CT+TT (Dominant model)					
Caucasian	2	1.336 (0.868–2.055)	0.188	0	0.795
South Asian	2	1.032 (0.544–1.957)	0.923	0	0.829
Europe	4	0.903 (0.511–1.597)	0.726	52.9	0.095
East Asian	2	1.424 (0.947–2.142)	0.090	0	0.768
Africa	2	0.773 (0.389–1.534)	0.461	3.5	0.309
Overall	12	1.095 (0.81–1.362)	0.412	9.2	0.355
TT vs CC+CT (Recessive model)					
Caucasian	2	0.618 (0.162–2.357)	0.481	63	0.100
South Asian	2	1.344 (0.259–6.987)	0.725	0	0.407
Europe	4	0.497 (0.279–0.886)	0.018	6.6	0.360
East Asian	2	0.894 (0.361–2.218)	0.809	63.7	0.097
Africa	2	1.395 (0.331–5.886)	0.651	21.9	0.258
Overall	12	0.701 (0.466–1.054)	0.087	27.7	0.173
CT vs CC+TT (Codominant model)					
Caucasian	2	0.971 (0.631–1.493)	0.894	0	0.397
South Asian	2	0.929 (0.475–1.818)	0.830	0	0.655
Europe	4	1.492 (1.046–2.126)	0.027	0	0.405
East Asian	2	0.760 (0.517–1.116)	0.161	0	0.625
Africa	2	1.328 (0.405–4.352)	0.640	56	0.132
Overall	12	1.059 (0.844–1.329)	0.621	16	0.287
MTHFR A1298C					
AA vs AC+CC (Dominant model)					
Caucasian	1	1.000 (0.468–2.138)	1.000	-	-
South Asian	2	0.877 (0.470–1.639)	0.681	0	0.921
Africa	2	1.881 (0.558–6.342)	0.308	57	0.127
Europe	2	0.906 (0.268–3.056)	0.873	73.1	0.054
East Asian	2	1.093 (0.583–2.050)	0.781	55.4	0.134
Overall	9	1.023 (0.730–1.435)	0.894	41	0.094
CC vs AA+AC (Recessive model)					
Caucasian	1	4.944 (0.604–40.476)	0.136	-	-
South Asian	2	0.432 (0.201–0.926)	0.031	0	0.517

(Continued)

Table 5 (Continued).

Genetic Models	Number of Studies	OR (95% CI)	P-value (OR)	I ² (%)	P-value (H)
Africa	2	0.756 (0.006–95.889)	0.910	85.1	0.010
Europe	2	0.858 (0.222–3.312)	0.824	61.3	0.108
East Asian	2	0.494 (0.238–1.027)	0.059	0	0.798
Ove55rall	9	0.784 (0.390–1.578)	0.495	55	0.023
AC vs AA+CC (Codominant model)					
Caucasian	1	0.702 (0.327–1.506)	0.634	-	-
South Asian	2	1.981 (1.108–3.543)	0.021	0	0.402
Africa	2	0.498 (0.247–1.005)	0.052	0	0.989
Europe	2	1.493 (0.915–2.435)	0.109	0	0.378
East Asian	2	1.058 (0.685–1.635)	0.799	7.2	0.299
Overall	9	1.075 (0.760–1.520)	0.682	43.8	0.076
MTRR A66G					
AA vs AG+GG (Dominant model)					
South Asian	2	0.821 (0.527–1.278)	0.381	9	0.295
East Asian	1	1.056 (0.553–2.017)	0.869	-	-
Overall	3	0.879 (0.619–1.249)	0.473	0	0.464
GG vs AA+AG (Recessive model)					
South Asian	2	0.952 (0.615–1.476)	0.827	0	0.626
East Asian	1	0.412 (0.067–2.540)	0.340	-	-
Overall	3	0.910 (0.594–1.392)	0.663	0	0.604
AG vs AA+GG (Codominant model)					
South Asian	2	1.171 (0.699–1.962)	0.548	43.5	0.183
East Asian	1	1.065 (0.550–2.061)	0.852	-	-
Overall	3	1.177 (0.850–1.629)	0.327	0	0.389

Notes: P < 0.05 shown in bold.

authors gave the explanation that it is possible that the anti-inflammatory effects are greater than the anti-proliferative properties in the mechanism of action of low-dose MTX for the treatment of RA patients.³⁸ However, Lima et al in a Portuguese cohort of RA patients have found that 677TT was associated with an approximately more than 3-fold increased risk of non-response to MTX when compared to MTHFR 677CC and 677C carriers.³⁰ A recent meta-analysis describing MTX kinetics and efficacy characteristics, it was shown that genotypes 677CT and 677TT carriers had 30% and 65% lower enzyme activity, respectively, when the C677T allele was present, and both genotypes were associated with reduced MTX efficacy and increased toxicity.³⁹ Very few studies have shown that the A1298C SNP is associated with MTX efficacy. One of the studies was Lilya et al who observed that the A allele of the A1298C polymorphism was related to good and moderate response to MTX

Table 6 Summary of the Analyzed Studies and the Distribution of MTHFR A1298C Genotypes

Study	Year	Ethnicity	Genotyping Method	Responders			Nonresponders			Mean Age, Years	Mean Disease Duration (Years)	Sex (Female/Male) (%)	MTX Dose (mg/Week)	Therapy Method	Date of End Point (Week)
				AA	AC	CC	AA	AC	CC						
Soukup T et al ²⁶	2015	Caucasian	Taqman	38	33	9	19	20	1	58.5 ± 12.6	No information	26.7/73.3	11 ± 2.7	Mixed	24
Ghodke-Puranik Y et al ²⁸	2015	South Asian	PCR RFLP	12	29	8	46	66	56	43.8 ± 10.4	5.6 ± 4.9	86/14	7.5–20	MTX	48
Boughrara W et al ²²	2017	Africa	Taqman	16	39	10	10	33	1	48.8 ± 13.44	9.28 ± 8.936	82.3/17.7	12.67 ± 2.33	MTX	24
Salazar J et al ²⁹	2014	Europe	Sanger sequence	24	16	5	6	6	4	55.62 ± 1.30	5.55	81.5/18.5	7.5–25	Mixed	24
Lv S et al ²¹	2018	East Asian	Sanger sequence	68	29	2	47	14	2	52.99 ± 13.81	4	82.4/17.6	8.92 ± 2.26	MTX	12
Berkani LM et al ²³	2017	Africa	Taqman	26	15	0	4	7	2	44.26 ± 14.41	<2	87.04/12.96	15.05 ± 2.4	Mixed	24
Lima A et al ²⁴	2016	Europe	Sequenom iPLEX	48	45	12	78	40	10	51 ± 15.6	0.3–51	84.1/15.9	2.5–25	MTX	12
Wang S et al ²⁰	2020	East Asian	PCR RFLP	112	52	12	66	38	16	54.6 ± 11.6	5.93	70.3/29.7	13.89 ± 1.99	MTX	24
Iqbal MP et al ²⁷	2015	South Asian	PCR-RFLP	19	7	2	16	5	2	42.87 ± 13.5	6.2–7.0	86.6/13.4	15–25	MTX	24

Notes: Mean age was measured using mean±SD; Mean disease duration and MTX dose were measured using mean±SD or range.




Figure 4 Forest plot of the correlation between MTHFR A1298C polymorphism and the efficacy of MTX in RA patients (AC vs AA+CC (codominant model)).
Abbreviations: OR, odds ratio; Cis, confidence interval.

treatment.²³ Although marked differences in the frequency distribution of MTHFR A1298C genotypes were observed in our study, they were not correlated with MTX treatment response.

Furthermore, the results of this study and the meta-analysis showed that the genotype frequency distribution of the MTRR A66G polymorphism was not significantly different between the response and non-response groups and could not predict response to MTX treatment. Consistent with our results, in the study of Chinese RA patients, Lv et al found no significant differences in the frequency distribution of MTRR A66G genotypes, alleles and haplotypes in the MTX response and non-response groups. Similarly, no differences were found in the five genetic statistical models.²¹ Separate studies of RA patients and healthy people groups in Mexican and South Indian Tamil populations also did not find any association between the MTRR A66G polymorphism and response to MTX treatment.^{31,40} In contrast to our findings, López-Rodríguez et al selected 25 relevant SNP sites and studied 956 RA patients from four regional groups, with the result that only MTRR A66G showed close correlation with response to MTX monotherapy.⁴¹ Among Portuguese RA patients, MTRR 66A carriers are associated with an approximately 2-fold increased risk of MTX non-response.²⁴ Chaabane et al suggested that the MTRR A66G polymorphism may produce different MTX treatment responses and toxic effects in RA patients from different ethnic groups.⁴² Our report of the correlation between the MTRR A66G SNP and MTX efficacy offers a possibility for studies in the Zhejiang RA population of China.

Table 7 Summary of the Analyzed Studies and the Distribution of MTRR A66G Genotypes

Study	Year	Ethnicity	Genotyping Method	Responders			Nonresponders			Mean Age, Years	Mean Disease Duration (Years)	Sex (Female/Male) (%)	MTX Dose (mg/Week)	Therapy Method	Date of End Point (Week)
				AA	AG	GG	AA	AG	GG						
Muralidharan N et al ³¹	2018	South Asian	PCR RFLP	54	128	46	33	50	24	42.72 ± 0.55	3.75 ± 0.23	93/7	10–25	MTX	24
Ghodke-Puranik Y et al ²⁸	2015	South Asian	PCR RFLP	13	22	14	41	82	45	43.8 ± 10.4	5.6 ± 4.9	86/14	7.5–20	MTX	48
Lv S et al ²¹	2018	East Asian	Sanger sequence	61	36	2	38	22	3	52.99 ± 13.81	4	82.4/17.6	8.92 ± 2.26	MTX	12

Notes: Mean age was measured using mean±SD; Mean disease duration and MTX dose were measured using mean±SD or range.




Figure 5 Forest plot of the correlation between MTRR A66G polymorphism and the efficacy of MTX in RA patients (AG vs AA+GG (codominant model)).
Abbreviations: OR, odds ratio; Cis, confidence interval.

On the study of the association of the three SNP sites with disease activity, we observed that ESR levels were not significantly different in the C677T polymorphism and were reduced in the 1298AC and 66GG genotypes. DAS28 was lowered in 677TT and 1298AC, TJC and SJC were not different among all genotypes, but neither was statistically different. Instead, each disease activity parameter was significantly lower in the response patient group compared to the baseline period ($p < 0.001$). Kurzawski et al found that ESR levels in the 677TT and 1298CC genotypes and TJC, SJC and DAS28 in patients with the 677TT and 1298AC genotypes were both significantly reduced.⁴³ Similar to our observed DAS28 with genotypes distribution changes. The current findings on the relationship between genetic polymorphisms and disease activity are varied, which may be related to factors such as patient cohorts, MTX dosage and different indicators of disease activity evaluation. Studies of Japanese RA patients reported that average DAS28 levels were markedly lower in patients with the MTHFR 1298AA genotype than in patients with the 1298AC/CC genotype.⁴⁴ González-Mercado et al evaluated the differences between polymorphic genotypes and DAS28, found a modest trend towards increased disease activity in patients heterozygous for A1298C and A66G ($p > 0.05$). There was no considerable difference between MTX monotherapy and combination therapy.⁴⁰

Gender, BMI, smoke status, RF-positive status, and age at onset all affect the response of RA patients to MTX treatment, leading to different levels of treatment response and disease activity.^{45,46} This may also explain the heterogeneous results that occurred when explored SNP genotypes and MTX disease response. The small number of participants in our study makes the results have some limitations. To explore our findings further, we also look forward to confirming our findings in a larger sample size.

Conclusion

This study reported the association of MTHFR C677T, MTHFR A1298C and MTRR A66G polymorphisms with MTX treatment response and disease activity in early RA patients in the Chinese population of Zhejiang. We observed that smoke, alcohol consumption, and males were possible influential factors for MTX non-response. In contrast, the SNP

sites we studied showed no correlation with MTX response and disease activity, and whether they can predict MTX treatment response needs to be repeated in a larger sample size. We also hope that this study may provide a possibility for personalized treatment of MTX in RA patients.

Ethics Statement

This study was approved by the Ethics Committee of Ningbo First Hospital and written informed consent was obtained from all participants. Subject identification information was not disclosed in the study, and patients were identified by numbers to indicate.

Acknowledgments

This research was supported by the Natural Science Foundation of Ningbo (No. 2021J037), the 3315 Innovation Team Foundation of Ningbo (No.2019A-14-C) and Special Research Assistant Program of the Chinese Academy of Sciences.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Lin YJ, Anzaghe M, Schülke S. Update on the pathomechanism, diagnosis, and treatment options for rheumatoid arthritis. *Cells*. 2020;9(4):880.
2. Jekic B, Maksimovic N, Damnjanovic T. Methotrexate pharmacogenetics in the treatment of rheumatoid arthritis. *Pharmacogenomics*. 2019;20(17):1235–1245. doi:10.2217/pgs-2019-0121
3. Smolen JS, Landewé R, Breedveld FC, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2013 update. *Ann Rheum Dis*. 2014;73(3):492–509.
4. Inoue K, Yuasa H. Molecular basis for pharmacokinetics and pharmacodynamics of methotrexate in rheumatoid arthritis therapy. *Drug Metab Pharmacokinet*. 2014;29(1):12–19.
5. Dervieux T, Greenstein N, Kremer J. Pharmacogenomic and metabolic biomarkers in the folate pathway and their association with methotrexate effects during dosage escalation in rheumatoid arthritis. *Arthritis Rheum*. 2006;54(10):3095–3103.
6. Murto T, Kallak TK, Hoas A, et al. Folic acid supplementation and methylenetetrahydrofolate reductase (MTHFR) gene variations in relation to in vitro fertilization pregnancy outcome. *Acta Obstet Gynecol Scand*. 2015;94(1):65–71.
7. Li X, Jiang J, Xu M, et al. Individualized supplementation of folic acid according to polymorphisms of methylenetetrahydrofolate reductase (MTHFR), methionine synthase reductase (MTRR) reduced pregnant complications. *Gynecol Obstet Invest*. 2015;79(2):107–112.
8. Frosst P, Blom HJ, Milos R, et al. A candidate genetic risk factor for vascular disease: a common mutation in methylenetetrahydrofolate reductase. *Nat Genet*. 1995;10(1):111–113.
9. Ashfield-Watt PA, Pullin CH, Whiting JM, et al. Methylenetetrahydrofolate reductase 677C→T genotype modulates homocysteine responses to a folate-rich diet or a low-dose folic acid supplement: a randomized controlled trial. *Am J Clin Nutr*. 2002;76(1):180–186.
10. Yamada K, Chen Z, Rozen R, et al. Effects of common polymorphisms on the properties of recombinant human methylenetetrahydrofolate reductase. *Proc Natl Acad Sci U S A*. 2001;98(26):14853–14858. doi:10.1073/pnas.261469998
11. Froese DS, Huemer M, Suormala T, et al. Mutation update and review of severe methylenetetrahydrofolate reductase deficiency. *Hum Mutat*. 2016;37(5):427–438. doi:10.1002/humu.22970
12. Hekmatdoost A, Vahid F, Yari Z, et al. Methyltetrahydrofolate vs folic acid supplementation in idiopathic recurrent miscarriage with respect to methylenetetrahydrofolate reductase C677T and A1298C polymorphisms: a randomized controlled trial. *PLoS One*. 2015;10(12):e0143569. doi:10.1371/journal.pone.0143569
13. Yang J, Luo G, Chen X. Individualized supplement of folic acid based on the gene polymorphisms of MTHFR/MTRR reduced the incidence of adverse pregnancy outcomes and newborn defects. *Niger J Clin Pract*. 2021;24(8):1150–1158. doi:10.4103/njcp.njcp_381_20
14. Du B, Tian H, Tian D, et al. Genetic polymorphisms of key enzymes in folate metabolism affect the efficacy of folate therapy in patients with hyperhomocysteinaemia. *Br J Nutr*. 2018;119(8):887–895. doi:10.1017/S0007114518000508
15. Arnett FC, Edworthy SM, Bloch DA, et al. The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum*. 1988;31(3):315–324. doi:10.1002/art.1780310302
16. Aletaha D, Neogi T, Silman AJ, et al. Rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Arthritis Rheum*. 2010;62(9):2569–2581. doi:10.1002/art.27584
17. Fransen J, van Riel PL. The disease activity score and the EULAR response criteria. *Rheum Dis Clin North Am*. 2009;35(4):745–757. doi:10.1016/j.rdc.2009.10.001
18. Ye S, Dhillon S, Ke X, et al. An efficient procedure for genotyping single nucleotide polymorphisms. *Nucleic Acids Res*. 2001;29(17):E88. doi:10.1093/nar/29.17.e88
19. Kolan SS, Li G, Grimalizzi F, et al. Identification of SNPs associated with methotrexate treatment outcomes in patients with early rheumatoid arthritis. *Front Pharmacol*. 2022;13:1075603. doi:10.3389/fphar.2022.1075603
20. Wang S, Zuo S, Liu Z, et al. Association of MTHFR and RFC1 gene polymorphisms with methotrexate efficacy and toxicity in Chinese Han patients with rheumatoid arthritis. *J Int Med Res*. 2020;48(2):300060519879588. doi:10.1177/0300060519879588
21. Lv S, Fan H, Li J, et al. Genetic Polymorphisms of TYMS, MTHFR, ATIC, MTR, and MTRR are related to the outcome of methotrexate therapy for rheumatoid arthritis in a Chinese population. *Front Pharmacol*. 2018;9:1390. doi:10.3389/fphar.2018.01390

22. Boughrara W, Benzaoui A, Aberkane M, et al. No correlation between MTHFR c.677 C > T, MTHFR c.1298 A > C, and ABCB1 c.3435 C > T polymorphisms and methotrexate therapeutic outcome of rheumatoid arthritis in West Algerian population. *Inflammation Res.* 2017;66(6):505–513. doi:10.1007/s00011-017-1034-6
23. Berkani LM, Rahal F, Allam I, et al. Association of MTHFR C677T and A1298C gene polymorphisms with methotrexate efficiency and toxicity in Algerian rheumatoid arthritis patients. *Heliyon.* 2017;3(11):e00467. doi:10.1016/j.heliyon.2017.e00467
24. Lima A, Bernardes M, Azevedo R, et al. Moving toward personalized medicine in rheumatoid arthritis: sNPs in methotrexate intracellular pathways are associated with methotrexate therapeutic outcome. *Pharmacogenomics.* 2016;17(15):1649–1674. doi:10.2217/pgs-2016-0067
25. Uribarri M, Ruiz-Larrañaga O, Arteta D, et al. Influence of MTHFR C677T polymorphism on methotrexate monotherapy discontinuation in rheumatoid arthritis patients: results from the GAPAID European project. *Clin Exp Rheumatol.* 2015;33(5):699–705.
26. Soukup T, Dosedel M, Pavěk P, et al. The impact of C677T and A1298C MTHFR polymorphisms on methotrexate therapeutic response in East Bohemian region rheumatoid arthritis patients. *Rheumatol Int.* 2015;35(7):1149–1161. doi:10.1007/s00296-015-3219-z
27. Iqbal MP, Ali AA, Mehboobali N, et al. Short Communication: lack of association between MTHFR gene polymorphisms and response to methotrexate treatment in Pakistani patients with rheumatoid arthritis. *Pak J Pharm Sci.* 2015;28(5):1789–1792.
28. Ghodke-Puranik Y, Puranik AS, Shintre P, et al. Folate metabolic pathway single nucleotide polymorphisms: a predictive pharmacogenetic marker of methotrexate response in Indian (Asian) patients with rheumatoid arthritis. *Pharmacogenomics.* 2015;16(18):2019–2034. doi:10.2217/pgs.15.145
29. Salazar J, Moya P, Altés A, et al. Polymorphisms in genes involved in the mechanism of action of methotrexate: are they associated with outcome in rheumatoid arthritis patients? *Pharmacogenomics.* 2014;15(8):1079–1090. doi:10.2217/pgs.14.67
30. Lima A, Monteiro J, Bernardes M, et al. Prediction of methotrexate clinical response in Portuguese rheumatoid arthritis patients: implication of MTHFR rs1801133 and ATIC rs4673993 polymorphisms. *Biomed Res Int.* 2014;2014:368681.
31. Muralidharan N, Gulati R, Misra DP, et al. Nonassociation of homocysteine gene polymorphisms with treatment outcome in South Indian Tamil Rheumatoid Arthritis patients. *Clin Exp Med.* 2018;18(1):101–107.
32. Weinblatt ME. Methotrexate in rheumatoid arthritis: a quarter century of development. *Trans Am Clin Climatol Assoc.* 2013;124:16–25.
33. Lucas CJ, Dimmitt SB, Martin JH. Optimising low-dose methotrexate for rheumatoid arthritis-A review. *Br J Clin Pharmacol.* 2019;85(10):2228–2234.
34. Maradit-Kremers H, Nicola PJ, Crowson CS, et al. Patient, disease, and therapy-related factors that influence discontinuation of disease-modifying antirheumatic drugs: a population-based incidence cohort of patients with rheumatoid arthritis. *J Rheumatol.* 2006;33(2):248–255.
35. Hoekstra M, van Ede AE, Haagsma CJ, et al. Factors associated with toxicity, final dose, and efficacy of methotrexate in patients with rheumatoid arthritis. *Ann Rheum Dis.* 2003;62(5):423–426.
36. Jansen G, de Rotte M, de Jonge R. Smoking and methotrexate inefficacy in rheumatoid arthritis: what about underlying molecular mechanisms? *J Rheumatol.* 2021;48(10):1495–1497.
37. Swierkot J, Słęzak R. Znaczenie badań farmakogenetycznych w efektywności leczenia metotreksatem chorych na reumatoidalne zapalenie stawów (część 2) [The importance of pharmacogenetic tests in evaluation of the effectiveness of methotrexate treatment in rheumatoid arthritis (part 2)]. *Postepy Hig Med Dosw.* 2011;65:207–215. Polish
38. Aggarwal P, Naik S, Mishra KP, et al. Correlation between methotrexate efficacy & toxicity with C677T polymorphism of the methylenetetrahydrofolate gene in rheumatoid arthritis patients on folate supplementation. *Indian J Med Res.* 2006;124(5):521–526.
39. Torres RP, Santos FP, Branco JC. Methotrexate: implications of pharmacogenetics in the treatment of patients with rheumatoid arthritis. *ARP Rheumatol.* 2022;1(3):225–229.
40. González-Mercado MG, Rivas F, Gallegos-Arreola MP, et al. MTRR A66G, RFC1 G80A, and MTHFR C677T and A1298C polymorphisms and disease activity in Mexicans with rheumatoid arthritis treated with methotrexate. *Genet Test Mol Biomarkers.* 2017;21(11):698–704.
41. López-Rodríguez R, Ferreira-Iglesias A, Lima A, et al. Replication study of polymorphisms associated with response to methotrexate in patients with rheumatoid arthritis. *Sci Rep.* 2018;8(1):7342.
42. Chaabane S, Marzouk S, Akrouit R, et al. Genetic determinants of methotrexate toxicity in tunisian patients with rheumatoid arthritis: a study of polymorphisms involved in the MTX metabolic pathway. *Eur J Drug Metab Pharmacokinet.* 2016;41(4):385–393.
43. Kurzawski M, Pawlik A, Safranow K, et al. 677C>T and 1298A>C MTHFR polymorphisms affect methotrexate treatment outcome in rheumatoid arthritis. *Pharmacogenomics.* 2007;8(11):1551–1559.
44. Kato T, Hamada A, Mori S, et al. Genetic polymorphisms in metabolic and cellular transport pathway of methotrexate impact clinical outcome of methotrexate monotherapy in Japanese patients with rheumatoid arthritis. *Drug Metab Pharmacokinet.* 2012;27(2):192–199.
45. Bansard C, Lequerré T, Daveau M, et al. Can rheumatoid arthritis responsiveness to methotrexate and biologics be predicted? *Rheumatology.* 2009;48(9):1021–1028.
46. Siddiqui A, Totouchian A, Jabar Ali JB, et al. Risk factors associated with non-response to methotrexate in rheumatoid arthritis patients. *Cureus.* 2021;13(9):e18112.

Pharmacogenomics and Personalized Medicine

Dovepress

Publish your work in this journal

Pharmacogenomics and Personalized Medicine is an international, peer-reviewed, open access journal characterizing the influence of genotype on pharmacology leading to the development of personalized treatment programs and individualized drug selection for improved safety, efficacy and sustainability. This journal is indexed on the American Chemical Society's Chemical Abstracts Service (CAS). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/pharmacogenomics-and-personalized-medicine-journal>