ORIGINAL RESEARCH

RETRACTED ARTICLE: LOXLI-ASI Drives The Progression Of Gastric Cancer Via Regulating miR-142-5p/PIK3CA Axis

This article was published in the following Dove Press journal: OncoTargets and Therapy

Ming Li Ou Cai Shiyun Tan

Department of Gastroenterology, Renmin Hospital of Wuhan University, Wuhan, Hubei 430000, People's Republic of China



Correspondence: Ming Li Department of Gastroenterology, Renmin Hospital of Wuhan University, Jiefang Road 238, Wuhan, Hubei Province 430060, People's Republic of China Email liming870221@sina.com



Background: Gastric cancer (GC) is a deadly disease, and wincidence is obecially high in East Asia including China. Recently, some long non-oding NA (lncR/As) have been identified as oncogenes or tumor suppressors. This only aimed to compute the function and mechanism of lncRNA LOXL1-AS1 on the procession of GC.

Methods: RT-PCR was done to measure exp. n levels LOXL1-AS1 and miR-142-5p in GC tissues. The association between part indexes and LOXL1-AS1 logic all lines AC, and BGC823 were used as cell expression was also analyzed. Hum GC models. CCK-8 and colony formation assays w conducted to assess the effect of LOXL1-AS1 on the proliferation of *Concern* lines. Transwel assay was conducted to determine the influence of LOXL1-AS1 cell migration and invasion. Furthermore, luciferase reporter confirm the assay was carried out to lationship of miR-142-5p with LOXL1-AS1. Additionally, Western blot done to tect the regulatory function of LOXL1-AS1 on PIK3CA, a targ viR-142-5p vivo experiment was also performed to validate the AS1 on the growth and metastasis of GC cells. roles and mechan n of D

Results: LOXL1-A blockpression in GC samples was significantly increased, which was corrected were unfavorable pathological indexes. Highly expressed LOXL1-AS1 was closely liked to shater over a survival time and post-progression survival time of the patients. LOC 1.4.61 marked y modulated the malignant phenotypes of GC cells. Additionally, overex, assed LOXL1-AS1 notably reduced the expression of miR-142-5p, but enhanced the expression level of PIK3CA. In vivo experiments further validated that knockdown of OXL1-AS1 inhibited the growth and metastasis of GC cells via regulating miR-142-5p and PharCA.

Conclusion: LOXL1-AS1 was a sponge of tumor suppressor miR-142-5p in GC, enhanced the expression of PIK3CA indirectly and functioned as an oncogenic lncRNA. **Keywords:** LOXL1-AS1, miR-142-5p, PIK3CA, gastric cancer

Introduction

Gastric cancer (GC) is one of the leading deadly cancer worldwide. More than 70% of diagnosed GC cases are patients from developing countries.¹ Although surgery, chemotherapy and targeted drugs offer symptomatic relief and modest improvement in survival for GC patients, the overall survival rate remains unfavorable.² Unfortunately, a majority of patients are diagnosed at an advanced stage, resulting in a high mortality rate. Moreover, in many cases, traditional therapies fail because of relapsing and distant metastasis due to chemoresistance or radioresistance.³ Hence, an in-depth investigation into the underlying mechanism of GC is highly desirable to explore the novel treatment strategies for GC.

OncoTargets and Therapy 2019:12 11345-11357

^{© 2019} Li et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress.com/terms.php you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 42 and 5 d our Terms (https://www.dovepress.com/terms.php).

Long non-coding RNA (lncRNA) is a class of noncoding RNA that is more than 200 nucleotides without protein-coding ability. Mounting studies indicate that lncRNA wields an enormous influence in the gene expression regulation at the transcription level, post-transcriptional level, and translation level.⁴ The role of lncRNAs as oncogenes or tumor suppressors in tumorigenesis and progression has been validated by the accumulating experimental demonstrations.5-7 LOXL1-AS1 is encoded in the complementary strands of LOXL1.8 It conducts a pivotal role in cell stress response and is closely related to Exfoliation Syndrome.9 Besides, high-expressed LOXL1-AS1 facilitates mesenchymal characteristics of glioblastoma via NF-KB pathway¹⁰ LOXL1-AS1 was verified as well to modulate the proliferation of medulloblastoma cells through the activation of PI3K/AKT pathway.¹¹ Nevertheless, the defined role of LOXL1-AS1 in GC cells remains poorly understood.

MicroRNA (miRNA) is a class of short-chain non-coding RNA molecules with a length of 21–25 nucleotides.¹² Abnormally expressed miRNA is a frequent event in the tumorigenesis and progression of diverse cancers. For instance, miR-214-5p inhibits the progression of mesothelioma via regulating the MDM2-p53 pathway;¹³ miR-1 restrains the proliferation of colorectal cancer cells;¹⁴ miR 221 promotes the migration of hepatocellular carcing a cells by targeting PHF2.¹⁵ Recent studies have veale that IncRNAs can interact directly with miRNAs and aulate their expression and activity.^{16,17} In tail, AA ATB has been found to modulate the perferation of lorectal cancer cells via sponging miR-2, c;¹⁸ T1 was verified to target miR-140 to facilitate proliferation f ovarian cancer cells;¹⁹ NEAT1 accelerates the proliferation of laryngeal squamous cell carcine a cel via regulating miR-107/ CDK6 axis.²⁰

The main and of our study acous explore LOXL1-AS1 expression of discribulations in GC cells, as well as to determine the impact of LOXL1-AS1 on malignant phenotypes of G occells and its downstream mechanism. Specifically, we demonstrated that LOXL1-AS1 displayed a high expression in GC tissues, and GC patients with highly expressed LOXL1-AS1 had shorter overall survival time and post-progression survival time. In terms of mechanism, LOXL1-AS1 can enhance the proliferation of GC cells via regulating the miR-142-5p/PIK3CA axis. Collectively, this study confirmed the crucial role of LOXL1-AS1/miR-142-5p/PIK3CA in regulating GC progression, thus providing novel clues for GC therapy.

Materials And Methods

Tissue Collection

All patients involved gave informed consent to the study and signed a written consent form. Tissue collection was conducted in accordance with the Declaration of Helsinki. This study was carried out under the approval of Ethics Review Committee of Renmin Hospital of Wuhan University. GC tissues and matched para-tumorous tissues were procured from 89 cases (49 males and 40 females, 26~70 years old) presented to our hospital from 2014 to 2019. All cancer tissues were pathol 1Uan confirmed as GC and no other malignant the nors were bund. No patients received neoadjuvant the v (chemo erapy or radiotherapy) before the peration. resciled tissues were stored in -196 ° fiquid togen with the following analysis.

Cell Culture

qRT-PCR

Total RNA was isolated from tissues or cells with RNAiso Plus Reagent (Takara, Dalian, China). About 1 μ g total RNA was reversely transcribed to cDNA with SuperScript First Strand cDNA System (Invitrogen, Grand Island, NY, USA). Then, real-time PCR was carried out. The relative expressions were calculated using $2^{-\Delta\Delta CT}$ method. The specific primer sequences are listed in Table 1.

Cell Construction

LOXL1-AS1 sequence lacking poly-A tail was procured from Shanghai GeneChem Co., Ltd. (Shanghai, China) and subcloned into pcDNA3.1 plasmid to establish LOXL1-AS1 overexpression cell line. LOXL1-AS1 shRNA was applied to knock down LOXL1-AS1 expression. Empty vector, pcDNA-LOXL1-AS1 vector or LOXL1-AS1 shRNA was transfected into AGS and BGC823 cells cultured in 6-well plate with Lipofectamine[®] 2000 (Invitrogen, Carlsbad,

Name	Primer Sequences
LOXLI-ASI	Forward: 5'-GATATGTTGGATGATGGA-3' Reverse: 5'-GATATGTTGGATGGATGA-3'
miR-142-5p	Forward: 5'-CAUAAAGUAGAAAGCACUACU-3' Reverse: 5'-CAUAAAGUAGAAAGCACUACU-3'
РІКЗСА	Forward: 5'-CCACGACCATCATCAGGTGAA-3' Reverse: 5'-CCTCACGGAGGCATTCTAAAGT-3'
GAPDH	Forward: 5'-GAAGGTGAAGGTCGGAGTC-3' Reverse: 5'-GAAGATGGTGATGGGATTTC-3'

Table I RT-qPCR Primer Sequences

USA). Twenty-four hours after the transfection, the cells were collected and processed for further analysis.

Cell Proliferation Assay

GC cells in each group were incubated in a 96-well plate for 12 hrs with 5000 cells per well. The culture medium was replaced with a complete medium containing 10 μ l CCK-8 reagent (Dojindo, Tokyo, Japan) at a specified time, and then the cells were incubated at 37 °C for 1 hr. Ultimately, Vmax microspectrophotometer (Beyotime, Shanghai, China) was adopted to measure the absorbance at 450 m

Colony Formation Assay

GC cells with a concentration of 1×10^{37} L we plant 1 in a 6-well plate. After 2 weeks of culture, we culture medium was discarded and the colonic ordere carefully washed with PBS twice. The colonies we then fixed with 10% paraformaldehyde for a min and scined with 0.1% crystal purple for 5 mins. Afterwards, the number of colonies formed has counted and recorded.

Transwell Migra on Ap Invasion Assay mocu ted in 2 well plate equipped with a Cells wer transwerchamber (2 um pore size; BD Biosciences, San A). GC cells were suspended in the upper Jose, CA, cells/well) with a serum-free medium, and chamber (2×1) the lower chamber contained 500 µL RPMI-1640 medium containing 10% FBS. Twenty-four hours later, the cells on the upper surface of the chamber were gently wiped clean with a cotton swab. Subsequently, the cells migrating the lower chamber were fixed using 95% ethanol for 20 mins and then stained by 0.1% crystal violet for 10 mins. Afterwards, the cells were counted under a microscope. In the invasion experiment, except that a layer of matrixgel (BD Biosciences, CA, USA) was coated on the bottom of

the transwell chamber, the other steps were the same as the migration experiment.

Luciferase Reporter Assay

A luciferase reporter vector (pMiR-LOXL1-AS1-WT/ pMiR-LOXL1-AS1-Mout) was constructed using DNA oligonucleotide and pMiR-Reporter vector. HEK-293T cells were co-transfected with pMiR-LOXL1-AS1-WT or pMiR-LOXL1-AS1-Mut and miR-142-5p mimics or negative control (NC miR). Luciferase activity was measured 24 hrs after transfection using the hel-luciferase reporter assay system (Promega, Madie a, WI, UCA).

Western Blot

The cells were lysed with RIPA but pplemented with protease inhibitors. Total totein extract was separated on 12% SDS-PA 2 gels 4 then transferred to nitrocellulose (NC) men res. The met bruce, after blocked by 5% fatfree milk, was in bated with primary antibodies at 4 °C for 12 primary alphodies included anti-PIK3CA (1:1000, hrs ocam, ab40776), anti-E-cadherin (1:1000, Abcam, ab1416), nti-N-cadhei (1:1000, Abcam, ab18203), anti-Vimentin 1000, Ab m, ab92547), anti-ZO-1 (1:1000, Abcam, and GAPDH (1:1000, Abcam, ab8245). Following ab90. the NC membrane was then rinsed with TBST solution and incubated at room temperature for 1 hr with the secondary antibodies (1:2000, Santa Cruz Biotechnology) conjugated to HRP. After that, color rendering was performed using hypersensitive ECL (Hubei Biossci Biotechnology Co, Ltd.). The automatic developing instrument (ChemiDocXRS imaging system) was adopted to detect the signal.

Tumorigenicity Assay

All animal assays were performed under the approval from Animal Care and Use Committee of Renmin Hosptial of Wuhan University. And the experiments were conducted in accordance with UKCCCR Guidelines for the welfare of animals in experimental neoplasia. BALB/c nude mice (female, 4 weeks old) were used in the tumorigenicity assay, and the nursing and use of experimental animals were conducted following the national standard. The mice were randomly divided into the sh-NC group and sh-LOXL1-AS1 group (10 mice in each group). GC cells (1×10^7 cells) injected subcutaneously to the left (sh-LOXL1-AS1 group) and right side (sh-NC group) of nude mice, and the tumor volume was monitored every 7 days. Thirty-five days later, the tumor was resected and weighed from the sacrificed mice. In the end, RT-qPCR was done to determine the expressions of LOXL1-AS1, miR-142-5p and PIK3CA mRNA, and Western blot was conducted to measure PIK3CA expression. In the lung metastasis study, 1×10^7 cells (sh-NC group or sh-LOXL1-AS1 group) were injected into a caudal vein of 10 mice in each group. Two weeks later, mice were killed and lung colonization was quantified through pathological examination.

Statistic Analyses

GraphPad Prism 7.0 was adopted for statistical analysis. Students' *t* test was carried out to analyze the difference in measurement data. Chi-square test was performed to analyze the correlation between LOXL1-AS1 expression and clinicopathological indexes. The survival curve was plotted by Kaplan–Meier method. All data were presented as Mean \pm SD. *P*<0.05 indicated statistical significance.

Results

The Up-Regulation Of LOXLI-ASI In GC Samples Was Related To Patients' Poor Prognosis

First of all, using TCGA data, we analyzed the expression level of LOXL1-AS1 in GC samples, and the results ind cated that LOXL1-AS1 was significantly up-regulated in G tissues compared with normal gastric tissues (1A). JRT-Furthermore, LOXL1-AS1 expression was de included by PCR in collected 89 GC tissues and a scent Consistent with the expression patter in TCC LOXL1-AS1 exhibited a higher expression AC cells as npared with matched adjacent tissues (Ngure N Moreover, compared with normal gastric endelial cell GES all of the GC cell lines showed a sher expession of LOXL1-AS1 aby e results we further investi-(Figure 1C). Based on the betwee LOXY -AS1 expression and gated the relation th the hoof Kaplan–Meier plotter long-term promosis y Highly expressed LOXL1-(http://kmpl com/ to be closely linked to shorter overall AS1 was prov survival time and ost-progression survival time of GC patients (Figure 1D). To sum up, the up-regulation of LOXL1AS1 may conduct a crucial role in GC progression.

The Expression Of LOXLI-ASI Is Correlated With The Pathological Indexes Of GC Patients

Then, we proceeded to analyze the correlation between LOXL1-AS1 expression and the pathological parameters

of the above 89 GC samples (Table 2). Chi-square test signified that the high expression of LOXL1-AS1 in GC cells was closely related to larger tumor (p=0.0165), local lymph node invasion (p=0.0065), distant metastasis (p=0.0084) and higher TNM stage (p=0.0041), but it had no significant correlation with age, gender and tumor differentiation, suggesting that LOXL1-AS1 may facilitate the tumorigenesis of GC.

LOXLI-ASI Modulates The Proliferation, Metastasis And Invasion Concells

The function of LOXL1-AS1 was urther explored after we draw the conclusion that LOXL1-A exhibite a significant up-regulation in GC rs. AGS and CGC 23 cell lines were selected to conjuct LC xL1-ASN overexpression I, respectively (Figure 2A). model and knock wn s detect the prolif-On this basis, K-8 assay eration ability of the above cells. The results demonstrated that LOXL1-AS1 over pression promoted the proliferation of A S cells. On the contrary, the proliferation ability of BG 823 cells we significantly inhibited in the downregulated LOXL1-A group as compared with the sh-NC . In addition, the colony formation ability group (1 cells was further explored. The results demonstrated 01 at the overexpression of LOXL1-AS1 increased the colony number of AGS cells, while compared with BGC823 lls transfected with sh-NC, the colony number of sh-LOXL1-AS1 group decreased significantly (Figure 2C). Next, we monitored the impact of LOXL1AS1 on metastasis by transwell assay. The results showed that overexpression of LOXL1-AS1 significantly facilitated the migration and invasion of AGS cells, while the migration and invasion of BGC823 cells transfected with sh-LOX1-AS1 were sigcompared with shNC nificantly restrained group (Figure 2D). Since our data suggested that LOXL1-AS1 may be implicated in the metastasis of cancer phenotype, Western blot was done to detect the expressions of biomarkers of EMT including Vimentin, N-cadherin, ZO-1 and E-cadherin. As we expected, the overexpression of LOXL1-AS1 can up-regulate the expressions of Vimentin, N-cadherin while down-regulate the expressions of ZO-1 and E-cadherin. On the contrary, knockdown of LOXL1-AS1 can reduce the expressions of Vimentin, N-cadherin but up-regulate the expressions of ZO-1 and E-cadherin (Figure 2E). These data implied that knockdown of LOXL1-AS1 impedes the migration and invasion of GC cells via inhibiting EMT.



Figure I The upregulation of LQ1 claSI in GC cells as associated with the poor prognosis of patients. (A) TCGA data showed the expression level of LOXLI-ASI was up-regulated in GC tissues. (Pi qRT-PCR was used to the termine LOXLI-ASI expressions in GC tissues and adjacent normal tissues. (C) qRT-PCR was performed to measure LOXLI-ASI expressions in normal gastric epithelial cells and 5 kinds of GC cell lines. (D) The overall survival and post-progression survival of GC patients with highly expressed and lowing typessed oXLI-ASI were evaluated by Kaplan–Meier method. **p<0.01, ***p<0.001.

miR-4.2-5p A Key larget Of LOXL1-ASI To elucit terme downstream mechanism of LOXL1-AS1 in Ge we carried out bioinformatics analysis through StarBar database (<u>http://starbase.sysu.edu.cn/</u> index.php). The data showed that LOXL1-AS1 contained the target site of miR-142-5p (Figure 3A). The results of qRT-PCR suggested that miR-142-5p was down-regulated in GC tissues (Figure 3B). We made a hypothesis that the dysregulation of miR-142-5p was due to the overexpression of LOXL1-AS1. To further verify that miR-142-5p is indeed a target for LOXL1-AS1, we detected the expressions of miR-142-5p in GC

cell lines with overexpressed or low-expressed LOXL1-AS1 by qRT-PCR. A decrease in the expression of miR-142-5p was observed in LOXL1-AS1 overexpressed cells whereas an increase in the expression of miR-142-5p was noted in cells with LOXL1-AS1 knocked down (Figure 3C). Additionally, miR-142-5p arrested the luciferase activity of pGL3-LOXL1-AS1 wild type, but had no effect on pGL3-LOXL1-AS1 mutate type (Figure 3D). These results suggested that miR-142-5p could bind to LOXL1-AS1 directly at miRNA recognition sites. Furthermore, the expression of LOXL1-AS1 in GC samples was reversely correlated with miR-142-

Variables	N	High	Low	P-value						
		Expression	Expression							
Age										
<60	38	16	22	0.1185						
≥60	51	30	21							
Gender										
Male	49	23	26	0.3213						
Female	40	23	17							
Tumor size										
<5 cm	25	18	7	0.0165						
≥5 cm	64	28	36							
Peritoneum dissemination										
Absent	33	19	14	0.3933						
Present	56	27	29							
Differentiation										
Well	18	10	8	0.8794						
Moderate	38	20	18							
Poor and others	33	16	17							
Lymph node invasio	n									
Absent	29	21	8	0.0065						
Present	60	25	35							
Distance metastasis										
Absent	37	13	24							
Present	52	33								
TNM stage										
1-11	55	35	20	0.0041						
III-IV	34		23							

Table	еź	2	Correla	ation	Between	Clinicopathological	Parameters		
And LOXLI-ASI Expression In 89 GC Patients									

J. Colectively t se data indicated that 5p (Figure 3 of LOXL1-AS1 and negatively miR-142-5 was a A GC tissues. To further investigate the modulated by LOXL1-AS1 and miR-142-5p, we interaction betwee co-transfected miR-142-5p in GC cells with overexpressed LOXL1-AS1 (Figure 4A and B). It was observed that overexpressed miR-142-5p attenuated the effects of overexpressed LOXL1-AS1 on the malignant phenotypes of GC cells (Figure 4C and D) (p < 0.05). These findings suggested that LOXL1-AS1 accelerates the proliferation, migration and invasion of GC cells via regulating miR-142-5p.

LOXLI-ASI/miR-142-5p Regulates The Expression Of PIK3CA

After confirming that LOXL1-AS1 could modulate miR-142-5p expression, we proceeded to determine the downstream target of miR-142-5p. In non-small cell lung cancer, miR-142-5p exerted the function of tumor suppressor via targeting PIK3CA;²¹ on the other side, PIK3CA has been validated to regulate the progression of GC.²² Based on the above knowledge, we carried out further study and found that in GC cell lines, LOXL1-AS1 knockdown could notably reduce the expression of PIK3CA both at mPLA and rotein level, while overexpressed LOXL1-AS alayed the o osite role (Figure 5A and B). In addition the hole expression of miR-142-5p reduced the expression of PIK3C and counteracted the effect of the upregulation of PL3CA induced by LOXL1--PCR in cated that LOXL1-AS1 (Figure 5C). ather, y related to the expression of PIK3CA AS1 was posit (Figure 5D). In sh LOXL1-AS1 could modulate the expression PIK3CA NGC cells via miR-142-5p.

LC (LI-ASI) Modulates The Progression Of CS Cells In Vivo

etermine whether LOXL1-AS1 affects the growth of nors in vivo, we constructed a stable LOXL1-AS1 lowexpression BGC823 cell with a lentivirus system igure 6A). On this basis, a subcutaneous tumorigenesis model was established. As shown, the tumor formed by BGC823/sh-LOXL-AS1 cells had a smaller volume and weight than that of BGC823/sh-NC cells (Figure 6B–D). To determine the influence of LOXL1-AS1 knockdown on cell metastasis in vivo, BGC823/sh-LOXL1-AS1 and BGC823/sh-NC cells were transplanted into the lateral tail vein of nude mice. Two weeks later, mice were sacrificed and lung metastases were monitored. Consistent with the results of in vitro experiments, compared with BGC823/sh-NC cells, lung metastasis of mice injected with BGC823/sh-LOXL1-AS1 cells was significantly hindered (Figure 6E). Western blot and qRT-PCR were also conducted, respectively, to detect the expression level of PIK3CA in the tumors obtained from subcutaneous tumorigenesis model, and PIK3CA in the sh-LOXL1-AS1 group was found to be significantly lower than that of the sh-NC group (Figure 6E). These results indicated that knockdown LOXL1-AS1 can inhibit the growth of GC cells via regulating PIK3CA in vivo.



Figure 2 LOXLI-ASI can modulate the proliferation, migration and invasion of GC cells. (A) qRT-PCR was done to detect the transfection efficiency of AGS and BGC823. (B) CCK-8 assay was carried out to detect the proliferation of GC cells transfected with pcDNA-LOXLI-ASI or sh-LOXLI-ASI. (C) The cloning ability of cells was detected by colony formation assay. (D) Transwell assay was conducted to monitor the migration and invasion of cells. (E) Western blot and qRT-PCR were used to detect the change of EMT markers after transfection, respectively. *p<0.05, **p<0.01.



Figure 3 miR-142-5p is a target of LOXL1-AS1. (A) StarBase predicted the binding sites of LOXL1-AS1 to miR-142-5p. (B) The expressions of miR-142-5p in GC tissues and adjacent normal to des were detected by qRT-PCR. (C) qRT-PCR was performed to detect the expressions of miR-142-5p in the cell model with over-expressed or knock-down LOXL1-AS1. (D) The targeting relation of LOXL1-AS1 with miR-142-5p was verified by luciferase reporter gene assay. (E) The correlation between the expression levels of LOXL1-AS1 and miR-142-5p. ****p<0.001.

Discussion

The impacts of abnormally expressed LOXL1-AS1 on various malignancies have been well documented in a number of studies. For instance, LOXL1-AS1 regulates the proliferation of medulloblastoma cells.¹¹ Moreover, LOXL1-AS1 can facilitate the proliferation, migration and invasion of osteosarcoma cells.²³ In the current study, up-regulated LOXL1-AS1 was noted in GC tissues and cells, whose high expression was verified to be closely linked to adverse clinical features. To elucidate the role of LOXL1-AS1 in GC, we



Figure 4 miR-142-5p reverse to rarcinogenic fect induced by overexpressed LOXL1-AS1. (A) The expression of LOXL1-AS1 was detected by qRT-PCR after AGS cells with LOXL-AS1 was detected by qRT-PCR after AGS cells with LOXL-AS1 was co-transfected with miR-142-5p. (B) The expression of miR-142-5p was detected by qRT-PCR after AGS cells with LOXL-AS1 was co-transfected with miR-142-5p. (D) The migration of Wich as detected by CCK-8 after AGS cells with LOXL-AS1 were co-transfected with miR-142-5p. (D) The migration and invasion of cells were contransfected by ranswell assay after AGS cells with LOXL-AS1 were co-transfected with miR-142-5p. *p<0.05, **p<0.01, ***p<0.001.

overexpresser and knocked down LOXL1-AS1 to verify its effect, and prover its oncogenic function. Besides, it was further validated that LOXL1-AS1 was associated with EMT markers' expressions in GC cells. These findings confirmed the carcinogenic role of LOXL1-AS1 in the progression of GC.

MiRNA plays a pivotal role in tumor progression. Accumulating evidence have verified that lncRNA can interact directly with miRNA and acts as competitive endogenous RNA (also called "ceRNA") to modulate its expression and activity.²⁴ For instance, LOXL1-AS1, as a ceRNA of miR-324-3p, enhances the progress of cholangiocarcinoma by regulating ABCA1;²⁵ LOXL1-AS1 mediates the cell cycle of prostate cancer cells through regulating miR-541-3p and CCND1;⁸ in addition, LOXL1-AS1 mediates the adriamycin resistance of prostate cancer cells by sponging miR-let-7a-5p.²⁶ It is reported that overexpressed miR-142-5p can arrest the progression of GC, but its specific mechanism remains elusive.²⁷ In the present study, we assumed that LOXL1-



Figure 5 LOXLI-ASI increases to expression of PIK3CA in vitro. (A) qRT-PCR and Western blot were used to detect the expressions of PIK3CA in cell models with overexpressed or knock-down LOXLI-ACID 2) Western but was done to detect the expressions of PIK3CA after GC cells with overexpressed LOXL-ASI was co-transfected with miR-142-5p. (D) are complation to reen the expression levels of LOXLI-ASI and PIK3CA. **p<0.01, ***p<0.001.

AS1 can targe me CNA to play a role in GC. To confirm this hypothesis, presorted to a bioinformatics analysis, which predicts the distence of a binding site between miR-142-5p and LOXL1-AS1. We overexpressed and knocked down LOXL1-AS1 to detect miR-142-5p expressions. The results suggested that LOXL1-AS1 reversely regulated miR-142-5p expression in GC cells. In GC cells, LOXL1-AS1 expression was verified to be reversely correlated with miR-142-5p. In addition, luciferase reporter gene assay was carried out to confirm that LOXL1-AS1 can bind directly to miR-142-5p. The

results indicated that LOXL1-AS1 could directly target miR-142-5p in GC.

Phosphoinositide 3-kinases (PI3Ks) protein family is involved in the regulation of malignant behaviors of cancer cells.^{28–30} PIK3CA is a gene encoding phosphatidylinositol-3 kinase (PI3K) p110 α catalytic subunit. The abnormal up-regulation of PIK3CA expression can promote proliferation, migration and invasion of cancer cells via enhancing the catalytic activity of PI3K and activating PI3K-AKT signaling pathway.³¹ In this study, we attempted to explore the downstream target of miR-



Figure 6 LON the I modulates the growth and metastasis of GC in vivo. (A) qRT-PCR was performed to detect the expressions of LOXLI-ASI and miR-142-5p in stable BGC823 cells when OXL-ASI knocked down. (B) A representative mice carrying tumors (left: sh-LOXLI-ASI group; right: sh-NC group) was shown. (C) The tumor volume of nude mice tum sh-NC or sh-LOXLI-ASI group was measured every 7 days, and the tumor quality was monitored for 35 days. (D) The weight of isolated tumor in sh-NC group or sh-LOXLI-ASI group. (E) Knock-down LOXLI-ASI notably decreased the number of lung metastasis nodules. (F) Western blot and qRT-PCR were performed, respectively, to detect the expression level of PIK3CA in the tumors of the mice from sh-NC group or sh-LOXLI-ASI group. *p<0.05, **p<0.01, ***p<0.001.

142-5p in GC cells. Bioinformatics suggested PIK3CA is a key oncogene that can regulate a variety of cell processes in GC and may be one of the candidate targets of miR-142-5p. Previous studies have confirmed that PIK3CA was a target of miR-142-5p in NSCLC cells.²¹ In this study, we found that the overexpression of miR-

142-5p can suppress the expression of PIK3CA, which was consistent with the previous report. Besides, overexpressed LOXL1-AS1 could notably increase the expression of PIK3CA both at mRNA and protein level, and LOXL1-AS1 is positively correlated with PIK3CA in GC tissues. Interestingly, the overexpression of miR-142-5p can partly offset the promoting effect of LOXL1-AS1 on the expression of PIK3CA. These results signified that LOXL1-AS1 could promote the expression of PIK3CA by targeting miR-142-5p in GC.

This study has certain limitations. Other downstream miRNA of LOXL1-AS1 needs to be identified in the following studies. What' more, in addition to acting as ceRNA, lncRNA also wields its function by regulating mRNA translation or binding to proteins.³² It is quite interesting whether LOXL1-AS1 plays its role via these mechanisms in GC, which also requires further investigation in the future. In summary, our findings highlight a unique lncRNA-regulated mechanism in GC, that is LOXL1-AS1 facilitates the proliferation and metastasis of GC cells by regulating the miR-142-5p/PIK3CA axis.

Data Availability Statement

The data used to support the findings of this study are available from the corresponding author upon request.

Funding

This study is supported by Natural Science Foundation of Hubei Province (2019CFB142).

Disclosure

The authors declare that they have no compending in rest.

References

- Bray F, Ferlay J, Soerjomataram I, Sich BL, Torre LA demal A. Global cancer statistics 2018: GLOBALAN exprates of incidence and mortality worldwide for 36 cancers in 185 converses. *CA Cancer J Clin.* 2018;68(6):394–424. doi:10.3322/caac.21492
- 2. Sheng J, Wang L, Han Y, etc. Dual roles of protein as a template and a sulfur provider: a general approach to netal sulfides for efficient photo-thermal therapy of cancer. St. V. 2008;14:1. doi:10.1002/smll.v14.1
- Duarte HO, Gomman Machae JC, Reis A. Gastric cancer: basic aspects. *Helico acter*. 208;23(Sup. 12:42523. doi:10.1111/hel.2018. 23.issue-S14
- 4. Fatica A, Baroni Januar , oding RNAs: new players in cell differentiation and evelopment. *Nat Rev Genet*. 2014;15(1):7–21. doi:10.1038/nrg30
- 5. Kapranov P, Cheng Dike S, et al. RNA maps reveal new RNA classes and a possible function for pervasive transcription. *Science*. 2007;316(5830):1484–1488. doi:10.1126/science.1138341
- Huarte M, Rinn JL. Large non-coding RNAs: missing links in cancer? *Hum Mol Genet*. 2010;19(R2):R152–61. doi:10.1093/hmg/ddq353
- Zhou XY, Liu H, Ding Z-B, Xi H-P, Wang G-W. lncRNA SNHG16 promotes glioma tumorigenicity through miR-373/EGFR axis by activating PI3K/AKT pathway. *Genomics*. 2019. doi:10.1016/j. ygeno.2019.06.017
- Long B, Li N, Xu X-X, et al. Long noncoding RNA LOXL1-AS1 regulates prostate cancer cell proliferation and cell cycle progression through miR-541-3p and CCND1. *Biochem Biophys Res Commun.* 2018;505(2):561–568. doi:10.1016/j.bbrc.2018.09.160

- Hauser MA, Aboobakar IF, Liu Y, et al. Genetic variants and cellular stressors associated with exfoliation syndrome modulate promoter activity of a lncRNA within the LOXL1 locus. *Hum Mol Genet*. 2015;24(22):6552–6563. doi:10.1093/hmg/ddv347
- Wang H, Li L, Yin L. Silencing lncRNA LOXL1-AS1 attenuates mesenchymal characteristics of glioblastoma via NF-kappaB pathway. *Biochem Biophys Res Commun.* 2018;500(2):518–524. doi:10.1016/j.bbrc.2018.04.133
- Gao R, Zhang R, Zhang C, et al. LncRNA LOXL1-AS1 promotes the proliferation and metastasis of medulloblastoma by activating the PI3K/AKT pathway. *Anal Cell Pathol (Amst)*. 2018;2018:9275685.
- 12. Voinnet O. Origin, biogenesis, and activity of plant microRNAs. *Cell*. 2009;136(4):669–687. doi:10.1016/j.cell.2009.01.046
- Singh A, Bhattacharyya N, Srivastava A, et al. MicroRNA-215-5p treatment suppresses mesothelioma progression view of 2002-p53-signaling axis. *Mol Ther.* 2019;27:1665–1680. doi:10.016/j.ym. 2019.05.020
- Forouzan JZ, Javeri A, Fakhr TM. Theor suppressive effects of the pleiotropically acting miR-195 in colourtal cancer ce a. *EXCLI J*. 2019;18:243–252. doi:10.1717/j.excli201.0166
- Fu Y, Liu M, Li F, et al. Mill 221 promotes in stocy afar carcinoma cells migration via argeting 14HF2. *Ined Res Int.* 2019;2019:4371405, doi: 10.1551/19/4371405
- 16. Cao MX, Jiang Yu, Tang Yu, Liang X, The crosstalk between lncRNA and peroRNA in other mustasis: orchestrating the epithelial-metanel plasticity. *Incotarget*. 2017;8(7):12472– 12483. doi:10.18632/perotarget.13957
- Bayourint S, Sayed A, Loskova Z, et al. Crosstalk between long non-oding RNAs and Mick RNAs in health and disease. *Int J Mol* 2016;17(3):356. doi:10.3390/ijms17030356
- 18. too Z, Zhou H, Wang Y, Chen J, Ou Y. Regulatory effects of h RNA ATB tar ting miR-200c on proliferation and apoptosis of color tal cancer cells. J Cell Biochem. 2019. doi:10.1002/jcb.29180
- Ding Y, Lung Q, Li Y, Wang Y. Amplification of IncRNA PVT1 notes ovarian cancer proliferation by binding to miR-140. *Mamm Genom.* 2019;30:217–225. doi:10.1007/s00335-019-09808-1
- Wang P, Wu T, Zhou H, et al. Long noncoding RNA NEAT1 promotes laryngeal squamous cell cancer through regulating miR-107/CDK6 pathway. *J Exp Clin Cancer Res.* 2016;35:22. doi:10.1186/s13046-016-0444-6
 Wang Z, Liu Z, Fang X, Yang H. MiR-142-5p suppresses tumorigenesis by targeting PIK3CA in non-small cell lung cancer. *Cell Physiol Biochem.* 2017;43(6):2505–2515. doi:10.1159/000484459
- 22. Jang SH, Kim K-J, Oh M-H, et al. Clinicopathological significance of elevated PIK3CA expression in gastric cancer. J Gastric Cancer. 2016;16(2):85–92. doi:10.5230/jgc.2016.16.2.85
- Chen S, Li W, Guo A. LOXL1-AS1 predicts poor prognosis and promotes cell proliferation, migration, and invasion in osteosarcoma. *Biosci Rep.* 2019;39:4. doi:10.1042/BSR20190447
- 24. Salmena L, Poliseno L, Tay Y, Kats L, Pandolfi PP. A ceRNA hypothesis: the Rosetta Stone of a hidden RNA language? *Cell*. 2011;146(3):353–358. doi:10.1016/j.cell.2011.07.014
- 25. Zhang B, Zhou M, Zou L, et al. Long non-coding RNA LOXL1-AS1 acts as a ceRNA for miR-324-3p to contribute to cholangiocarcinoma progression via modulation of ATP-binding cassette transporter A1. *Biochem Biophys Res Commun.* 2019;513(4):827–833. doi:10.1016/j. bbrc.2019.04.089
- Bai T, Liu Y, Li B. LncRNA LOXL1-AS1/miR-let-7a-5p/EGFR-related pathway regulates the doxorubicin resistance of prostate cancer DU-145 cells. *IUBMB Life*. 2019;71:1537–1551. doi:10.1002/iub.v71.10
- 27. Yan J, Yang B, Lin S, Xing R, Lu Y. Downregulation of miR-142-5p promotes tumor metastasis through directly regulating CYR61 expression in gastric cancer. *Gastric Cancer*. 2019;22(2):302–313. doi:10.1007/s10120-018-0872-4
- Wang D, Chen J, Chen H, et al. Leptin regulates proliferation and apoptosis of colorectal carcinoma through PI3K/Akt/mTOR signalling pathway. *J Biosci.* 2012;37(1):91–101. doi:10.1007/s12038-011-9172-4

- Kim EH, Suresh M. Role of PI3K/Akt signaling in memory CD8 T cell differentiation. *Front Immunol.* 2013;4:20. doi:10.3389/ fimmu.2013.00020
- Kou C, Cao X, Qin D, et al. Over-expression of LYRM1 inhibits glucose transport in rat skeletal muscles via attenuated phosphorylation of PI3K (p85) and Akt. *Mol Cell Biochem*. 2011;348(1–2):149– 154. doi:10.1007/s11010-010-0649-5
- 31. Khan N, Jajeh F, Eberhardt EL, et al. Fisetin and 5-fluorouracil: effective combination for PIK3CA-mutant colorectal cancer. *Int J Cancer.* 2019;145:3022–3032. doi:10.1002/ijc.v145.11
- 32. Zhang A, Zhao JC, Kim J, et al. LncRNA HOTAIR enhances the androgen-receptor-mediated transcriptional program and drives castration-resistant prostate cancer. *Cell Rep.* 2015;13(1):209–221. doi:10.1016/j.celrep.2015.08.069

OncoTargets and Therapy

Dovepress

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic

agents and protocols on patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/ testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/oncotargets-and-therapy-journal