

Lipid Metabolism Reprogramming of Immune Cells in Acne: An Update

Shuhui Wu¹, Xi Zhang², Yun Wang¹, Huie Zheng¹, Mingfang Zhu¹

¹Department of Dermatology, Key Laboratory of Vascular Biology and Translational Medicine, Education Department of Hunan Province, The Second Affiliated Hospital of Hunan University of Chinese Medicine, Changsha, Hunan, People's Republic of China; ²Department of Physical Education and Health Promotion, Hunan University of Technology and Business, Changsha, Hunan, People's Republic of China

Correspondence: Mingfang Zhu, Department of Dermatology, Key Laboratory of Vascular Biology and Translational Medicine, Education Department of Hunan Province, The Second Affiliated Hospital of Hunan University of Chinese Medicine, 233 Caie North Road, Kaifu District, Changsha, Hunan, People's Republic of China, Tel/Fax +86 731 8491 7727, Email 320492@hnucm.edu.cn

Abstract: Acne vulgaris is one of the most widespread skin conditions and the main reason for visiting a dermatologist. Inflammatory response and abnormal infiltrations of immune cells are the main pathogenesis of acne. The increased lipid is the prerequisite for the acne, and the perturbation of lipid composition and content is consistent with the severity of acne. Furthermore, the increased lipid production not only contributes to the occurrence and development of acne, but also sensitizes the function of immune cells. The lipid metabolic dysfunction aggravates the severity of local tissue and provides pro-inflammatory-cytokine cues, which indicates the crucial roles of lipid metabolism on immune cells. Recent advances have demonstrated the lipid metabolism reprogramming of various immune cells in acne lesion. The abnormal lipid accumulation, lipolysis, and fatty acid oxidation lead to the activation and differentiation of immune cells, which promotes the pro-inflammatory cytokines production. Thus, this review discusses the emerging role of lipid metabolism reprogramming of immune cells in the progress of acne and aims to constitute food for others' projects involved in acne research.

Keywords: acne vulgaris, inflammatory response, lipid, reprogramming

Introduction

Acne vulgaris (acne) is a chronic inflammatory skin disease of the sebaceous glands characterized by papules, nodulocystic lesions, and inflammation of the pilosebaceous follicles.¹ As the eighth most prevalent disease, acne affects an estimated 9.4% population of the world and 8.1% of China. Acne patients suffer from increased risk of psychiatric comorbidities specifically depression and suicidality.² The pathogenesis of acne is closely related to endocrine, hyperkeratinization, microorganisms and the overcolonization of pilosebaceous units. The upregulated androgens, especially testosterone and dihydrotestosterone, and hyperinsulinemia caused by insulin resistance induce sebocyte activation, leading to the increased sebum production.³ The sebum accumulation creates a new environment that foster microorganisms putatively *Cutibacterium acnes* (*C. acnes*) proliferation.⁴ The ensuring immune infiltration is triggered by the infections of *C. acnes*, accompanied with inflammatory factor secretion. These changes drive pilosebaceous follicles from homeostasis into an inflammatory state, which results in the form of microcomedone. Then the microcomedone evolves into a comedone contributing to the acne.

Inflammatory response throughout acne is the hallmark of acne, and multiplied in large quantities by microorganisms especially *C. acnes*. Abnormal infiltrations of innate immune cells, such as, monocytes, dendritic cells, neutrophils, macrophages, and adaptive immune cells, such as T lymphocytes and B lymphocytes, are thought to be crucial in the pathogenesis of acne.⁵ Besides the effect of inflammation of immune cells, the lipid metabolism reprogramming of immune cells has been implied in the development of acne. Metabolic reprogramming first recognized a century ago is an approach for cells to support the requirements of growth and proliferation by altering the metabolic patterns including glucose, lipid, amino acid metabolism and so on.⁶ Lipid metabolic dysfunction is considered to be the prerequisite in the progress of acne. Skin surface lipids (also called sebum) are comprised of triglycerides, waxy, squalene, free fatty acids and cholesterol.⁷ The emerging insights arising from advanced techniques in metabolic analysis indicated that glycerophospholipids, fatty acyls, and sterol lipids were significant

increased and prenol lipids, saccharolipids were decreased in the cases with acne compared to their normal counterparts.⁸ In addition, the average level of triglycerides, unsaturated free fatty acids and cholesterol showed an increase trend from mild to moderate adolescent acne, which suggested the perturbation of lipid composition and content was consistent with the severity of acne.⁹ The lipid metabolic dysfunction impaired the skin barrier function and provided several pro-inflammatory-cytokine cues. Recent advances have highlighted the lipid metabolic reprogramming in the immune cells, which is critical for the immune response and lipid metabolism. Lipid metabolic reprogramming interfered with immune responses by not only promoting immune cells proliferation and differentiation, but also modulating the immunological activity in acne. Accordingly, the lipid metabolic reprogramming in immune cells promote sebocytes to activate sebum synthesis and secretion, which further aggravate the acne lesions. Therefore, this review mainly discusses the reprogrammed lipid metabolic activities and the possible pathways in immune cells in the progress of acne, with the aim of providing a comprehensive foundation for the development of novel therapies.

Lipid Metabolism Reprogramming of Immune Cells in Acne Neutrophils

Neutrophils, as the most abundant leukocytes in the peripheral blood, are characterized by multi-lobulated or rod-shaped nucleus and granules, that is why neutrophils are also known as polymorphonuclear neutrophils.¹⁰ Neutrophils derived from the myeloid progenitor cells are differentiated and recruited to sites of inflammation after certain triggers.¹¹ The granules of neutrophils are classified into primary, secondary, and tertiary granules, consisting of myeloperoxidase, defensins, proteinase 3, collagenase, gelatinase, lactoferrin, and sialidase.¹² Neutrophils phagocytose, release granular contents and produce the reactive oxygen species (ROS) to eliminate pathogenic agents.^{13,14} In addition to phagocytosis and degranulation, they release neutrophil extracellular traps (NETs) that resemble a net composed largely of decondensed chromatin where various cytoplasmic proteins are attached.^{15,16} NETs play a vital role in capturing and digesting pathogenic agents like bacteria, fungi and viruses at the infection site to prevent severe tissue damage and promote the tissue healing.^{17,18} Whereas excessive NETs alter the homeostasis of immune and further exacerbate the release of pro-inflammatory cytokines, which aggravate the severity of local inflammation and are detrimental to the local tissue.^{19,20}

A notable increased proportions of infiltrated neutrophils were observed around the acne lesions, less than 48 h after their development compared to the healthy individuals.²¹ Consistently, neutrophil gelatinase-associated lipocalin, also named lipocalin 2 (LCN-2), were markedly upregulated in the sebaceous glands and hair follicles of patients with acne.²² LCN-2 initially isolated from neutrophils, is a novel adipocytokine responsible for the transportation of small lipophilic molecules, such as steroids, fatty acids, lipopolysaccharides, etc.²³ As a vital regulator of lipid metabolism, LCN-2 causes a wide spectrum of biological effects, including infections, cell differentiation, apoptosis, liver injury, and even tumorigenesis.^{24–26} The level of LCN-2 was reported to be positive correlation with obesity and hypertriglyceridemia.²³ What is more, the expression of LCN-2 was obviously upregulated in patients with fatty liver diseases or atherosclerotic diseases, which indicated the considerable role of LCN-2 in lipid metabolism.^{27–29} The transcription and secretion of LCN-2 were enhanced after the activation of toll-like receptors (TLRs) combined with the gram-positive bacteria (*C. acnes*) peptides on neutrophils.³⁰ On the one hand, the overexpression of LCN-2 was implicated to promote the production of pro-inflammatory mediators IL-6, IL-12, IL-8 and attenuate the secretion of anti-inflammatory mediators IL-10.^{31,32} The upregulated pro-inflammatory cytokines such as IL-8 and IL-6 were elucidated to be involved in the form of NETs, which further amplified the inflammatory response and contributed to the local damage.^{15,33,34} Moreover, the NETs formation was enhanced in patients with pyogenic arthritis, pyoderma gangrenosum and acne syndrome compared to the healthy individuals.³⁵ Thus, LCN-2 was confirmed to be a sensitive biomarker for inflammation.^{36,37} On the other hand, as a vital regulator of lipid metabolism, LCN-2 deficiency attenuated the expression of Fasn, Srebp2, and Ppar γ gene, LCN-2 deficiency disrupted the mammalian target of rapamycin (mTOR) signaling regulation of lipid metabolism.³⁸

Monocytes/Macrophages

Monocytes, originating from bone marrow, are recruited and differentiated into macrophages with the stimulus of macrophage colony-stimulating factor to in response to homeostasis and inflammatory signals.³⁹ According to the active

state, macrophages are mainly classified as either classically activated M1 macrophages or alternatively activated M2 macrophages.⁴⁰ M1 macrophages predominantly generate the pro-inflammatory cytokines to promote inflammation, contrarily, M2 macrophages produce anti-inflammatory cytokines to dampen inflammation.⁴⁰ Interestingly, the high proportions of infiltrated macrophages were readily detected in the acne lesions,²¹ the number of macrophages especially the M1 macrophages was increased as the acne progressed.⁴¹

Toll-like receptors (TLRs), a kind of pattern recognition receptors, are expressed on macrophages.⁴² Pathogen-associated molecular patterns, such as lipase, hyaluronidase, and proteases produced by *C. acne*, were identified by TLRs on macrophages to initiate the innate immune.⁴³ In a ligand-specific manner, activated TLRs recruited the adapter molecules myeloid differentiation primary response gene 88 (*MYD88*), upregulated *MYD88* activated mTOR signaling pathway by promoting S6K phosphorylation, which induced macrophages metabolic rewiring.⁴⁴ Acetyl-coenzyme A (CoA) carboxylase (ACC) enzymes, the central regulators of lipid metabolism, is the downstream of mTOR signaling pathway, ACC deficiency resulted in less redistribution of lipid species and impaired the pro-inflammatory macrophage activation compared to the control group after LPS stimulus.⁴⁵ Furthermore, activated *MYD88* spontaneously elevated the expression of downstream mitogen-activated protein kinases (MAPK) signaling pathway.⁴⁶ Suppressing the activation of *MYD88* downregulated the phosphorylation of c-Jun N-terminal protein kinase (JNK), extracellular signal-regulated kinase (ERK1/2) and p38, the classical regulators of inflammatory cytokines.⁴⁷ Consequently, pro-inflammatory cytokines like IL-6, IL-8 were reduced by suppressing the *MYD88*/MAPK signaling pathways. In addition, excessive expression of *MYD88* caused a considerable difference on the activation of NOD-like receptor family pyrin domain containing three inflammasomes, which resulted in exaggeration of inflammation.⁴⁸ Recent studies in immunometabolism have shed light on the intricate links between cellular cholesterol metabolism and inflammatory response of macrophage. Accumulating research has displayed that cellular cholesterol metabolism was an essential component of macrophage activation programs, cholesterol metabolism was actively regulated during innate activation of macrophages. Cellular cholesterol levels were temporally changed after TLR activation, accumulated cholesterol was binded to *MYD88*, which was essential for the activation and self-oligomerization of *MYD88*. That said, cellular cholesterol metabolism was integral to innate activation of macrophages and closely related to *MYD88*-dependent inflammatory response in macrophage.⁴⁹ Meanwhile, macrophages took up the intracellular cholesterol and transformed into foam cells when exceeding the scavenging capacity, the foam cell formation was fundamental in the acne progress.⁵⁰

In the regard of M2 macrophages, they seemed to have a negative effect of lipogenesis and positive effect on lipolysis. Through polarization, IL-12 blunted the lipogenesis progress by inhibiting the activity of lipogenic enzymes, such as fatty acid synthase, and the expression of lipolytic enzymes was confirmed to be increased in the macrophages.⁵¹ Through the lipolysis in lipid droplets, triacylglycerols were disintegrated into diacylglycerols and monoacylglycerols, accompanied by fatty acid release, and significantly increased amounts of three out of four monoacylglycerols were found in M2 macrophages, which indicated the positive effect of lipolysis.⁵² What is more, neuropeptide Y prevented 7KC-induced foam cell formation in resting macrophages and shifted macrophages toward the anti-inflammatory M2-like phenotype to exert anti-inflammatory functions.⁵³

Mast Cells

Mast cells are well known in allergic inflammatory diseases and are considered to be one of the “first responders” to innate inflammation. Infiltration and degranulation of mast cells were observed in acne lesions.⁵⁴ Mast cells are rich in lipid rafts, highly ordered membrane microdomains enriched in cholesterol, glycosphingolipids, and certain proteins, lipid modification is tightly linked to the degranulation, endocytosis of mast cells.⁵⁵ Lipid reprogramming exerted a dependent role in mast cell activation. The phosphorylation of immunoreceptor tyrosine-based activation motif activated the mitogen promoter protein (MAK) kinase signaling pathway to decompose membrane phospholipid choline (PC) to produce arachidonic acid, then the arachidonic acid was the essential substance for the synthesis of arachidonic acids such as prostaglandin D2 (PGD2), leukotrienes, and platelet-activating factor, which induced the inflammatory response.⁵⁶ Importantly, research reported that supplementing with PC reduced the alteration of lipid profile and attenuated IgE-induced immune responses in mast cells, which suggested the importance of lipid metabolism in mast cells biological activity.⁵⁷ Surprisingly, the production of PGD2 was initiated by hydrolysis of membrane phospholipids

by phospholipase A2 (PLA2), an enzyme for cleaving cell membrane phosphodiacylglycerides, loss of PLA2 impaired the maturation of mast cells and failed to facilitate acute anaphylactic response.⁵⁸ In addition, palmitoylethanolamide, a pleiotropic endogenous lipid mediator, downmodulated mast cells activation and degranulation by stimulating the activity of diacylglycerol lipase α and β .⁵⁹

Dendritic Cells

Dendritic cells are the classical antigen-presenting cells to transport the peptide MHC class I complexes to the cell surface. Lipid metabolism has a great impact on the activation and inflammatory response of dendritic cells. When infected, genes related to lipid metabolism, including cholesterol and fatty acid biosynthesis-related genes were uniformly upregulated and resulted in the lipid accumulation in dendritic cells.⁶⁰ Meanwhile, the risk of infection was significantly elevated when dendritic cells underwent abnormal lipid peroxidation and greater lipid uptake, which was beneficial for the immune escape. Recent advance highlighted that electrophilic oxidatively truncated lipids covalently bind to heat shock protein 70 impaired the translocation of peptide MHC class I complexes and prevented the cross-presentation.⁶¹ Except electrophilic oxidatively truncated lipids, the fatty acid oxidation was active in dendritic cells in high-fat diet-induced obesity, which indicated that alteration of the lipid reprogram might help improve dendritic cells function in obese individuals.⁶²

T Lymphocytes

T lymphocytes, composed of T helper cells (Th cells), regulatory T cells (Tregs), cytotoxic T cells, memory T cells, natural killer cells and gamma delta ($\gamma\delta$) T cells, are essential for the innate immune response as well as adaptive immune response. Acne was characterized by various degrees of T cell infiltration. For instance, a Th1 response instead of Th2 response predominated in acne lesions,⁶³ and a higher infiltration of Th17 cells in acne lesions was detected relative to nonlesional skin areas of acne patients.²¹ Thus, some researches pointed out that acne was immune-mediated chronic inflammatory skin disease, especially characterized by Th1/Th17-mediated immune response, which suggested the pivotal role of Th1/Th17 in acne.^{63,64} On the contrary, Tregs were substantially lower, relative to nonlesional sites of acne patients and skin of healthy individuals.²¹

In comparison to the control group, acetylcoenzyme A carboxylase 1 (ACC1), a key enzyme in lipid homeostasis was positively linked with Th1/Th17 cells, Jiang et al recommended the correlation of ACC1 with Th17 and Th1 cells as the potential biomarker of acute ischemic stroke and implied the effect of Th1/Th17 cells in dyslipidemia.⁶⁵ The monophosphoryl lipid A had an negative effect on the Th1 cytokines and enhanced the production of the regulatory cytokine IL-10,⁶⁶ the pro-resolving lipid mediator lipoxin A4 dampened the activation of Th1/Th17 cells and reduced the inflammatory mediators.⁶⁷ In turn, Th1 cells upregulated interferon-gamma response signaling and antigen presentation pathways and downregulated lipid metabolism. While the lipid metabolism appeared to display a mix effect on T cells. Experiments have showed that ACC1-associated de novo fatty acid synthesis mediated the Th17 cell differentiation⁶⁸ and long-chain fatty acids enhanced Th1 and Th17 cell differentiation and proliferation via the P38-MAPK pathway, exacerbating the inflammatory disease in vivo.⁶⁹ Fatty acid-binding protein (FABP5) is responsible for fatty acid uptake, and it is highly expressed in Treg cells. In vitro studies show that the inhibition of FABP5 in naïve T cells suppresses Treg cell differentiation. The inhibition of *FABP5* activates cyclic GMP-AMP synthase (cGAS) stimulator of interferon gene- (STING-) dependent type I IFN signaling, leading to IL-10 production and Treg cell activation.⁷⁰ Sphingomyelin and glycosylphosphatidylinositol, abnormal in acne skin, are components of lipid rafts, which are closely to the activation and differentiation of naïve T cells.⁷¹ Ceramides, one of sphingosine-based lipid molecules, were confirmed to capable of upregulating Tregs and maintaining the Treg function, which ceramides were significantly lower in acne lesions.⁷² Moreover, the upregulated lipid synthesis but not fatty acid β -oxidation induced the IFN- γ production accompanying with higher activation of Th1 subsets, while inhibiting fatty acid β -oxidation led to less IL-17 production by pro-inflammatory Th17 cells.^{73,74}

B Lymphocytes

In acne, B lymphocytes were responsible for the long-standing scar, B-cell infiltrations, which were not found in early papules of acne, were involved in 23% of all acne scar specimen.^{75,76} Despite secreting antibodies is the hallmark of B cells, it is increasingly appreciated that B cells dysfunction contribute to the inflammatory cytokines production. DHA, a n-3 LCPUFAs, was incorporated into the B cell membrane to alter lipid microdomain clustering and enhanced the level of IgA.⁷⁷ In vivo, inhibiting B-cell-activating factor signaling, which caused a difference in promoting differentiation of the B-cell population, led to lipid reprogramming in liver. Lipid content was notably enriched in BAFF-R^{-/-} group, genes related to lipid metabolism such as fatty acid-binding protein, ACC, FAS were significantly upregulated in BAFF-R^{-/-} group.⁷⁸ Lipid G-protein coupled receptor *GPR55*, expressed by splenic plasma cells, was involved in plasma cells maturation. Lack of *GPR55* hyperactivated B cells, disturbed PC maturation and supported a possible relevance for complications in human atherosclerosis pathophysiology.⁷⁹ In conclusion, lipid metabolism interacts with B cells function.

Conclusion

Acne has proven challenging to treat due to its chronicity, a tendency to relapse, and inherent recalcitrance. To date, topical antibiotics, retinoids, hormonal therapy are still the first line and mainstay for the treatment of acne. Retinoids are the first approach used worldwide for the treatment of acne.⁸⁰ All the lines and recommendations of dermatologists and skin experts support the use of these substances, especially for acne vulgaris. Furthermore, lasers and light-based modalities like intense pulsed light, photodynamic therapy, and picosecond lasers are used in the treatment of refractory acne.⁸¹ However, research has shown that the effect of topical therapy, possibly leading to the damage of skin barrier, rebound postinflammatory hyperpigmentation and recurrence, is not obvious, which limits the widespread application.^{82–84} Considering the long duration of acne treatment, the visibility of the lesions, and the struggle with various side effects, researchers have paid more attention to TCM for fewer side effects and definite anti-inflammatory effect.⁸⁵ Infiltration of immunocytes and inflammatory response are the hallmark of acne, moreover, lipid levels, as well as abnormal metabolism of immunocytes also contribute to acne development and occurrence. Immunocytes, such as neutrophils, macrophages, T cells, B cells, and dendritic cells regulate their own phenotypes or functions via metabolic changes due to metabolite accumulation. When lipid accumulates, macrophages can be transformed from pro-inflammatory M1 to anti-inflammatory M2. In addition, pro-inflammatory cytokines were produced in the neutrophils when abnormal lipid peroxidation and greater lipid uptake happened. Thus, lipid metabolism reprogramming of immunocytes appears to contribute to the acne development and aggravate the inflammatory response. Furthermore, the immune response rate can be improved by regulating immune cell metabolism and improving the micro-environmental immune status. Inhibiting the lipid metabolism reprogramming can effectively decreasing the secretion of pro-inflammatory cytokines and improve the progress of acne. Therefore, further study on the cometabolic targets of acne and immune cells is of great importance for achieving satisfactory treatment outcomes that shorten the duration of acne treatment.

Data Sharing Statement

Data openly available in a public repository.

Funding

This work was supported by National Natural Science Foundation of China (8217437) and Scientific Research Project of Hunan Provincial Health Commission (202202124772).

Disclosure

The authors report no competing interests in this work.

References

1. Kostecka M, Kostecka J, Szwed-Guła O, Jackowska I, Kostecka-Jarecka J. The impact of common acne on the well-being of young people aged 15–35 years and the influence of nutrition knowledge and diet on acne development. *Nutrients*. 2022;14(24):5293. doi:10.3390/nu14245293
2. Kohn AH, Pourali SP, Rajkumar JR, Hekmatjah J, Armstrong AW. Mental health outcomes and their association to race and ethnicity in acne patients: a population-based study. *J Am Acad Dermatol*. 2022;87(1):140–142. doi:10.1016/j.jaad.2021.06.866

3. Bakry OA, El Shazly RM, El Farargy SM, Kotb D. Role of hormones and blood lipids in the pathogenesis of acne vulgaris in non-obese, non-hirsute females. *Indian Dermatol Online J.* 2014;5(5):S9–S16. doi:10.4103/2229-5178.144506
4. Zouboulis CC, Jourdan E, Picardo M. Acne is an inflammatory disease and alterations of sebum composition initiate acne lesions. *J Eur Acad Dermatol Venereol.* 2014;28(5):527–532. doi:10.1111/jdv.12298
5. Tangheiti EA. The role of inflammation in the pathology of acne. *J Clin Aesthet Dermatol.* 2013;6(9):27–35.
6. Faubert B, Solmonson A, DeBerardinis RJ. Metabolic reprogramming and cancer progression. *Science.* 2020;368(6487). doi:10.1126/science.aaw5473
7. Wertz PW. Epidermal lipids. *Semin Dermatol.* 1992;11(2):106–113.
8. Zhou M, Gan Y, He C, Chen Z, Jia Y. Lipidomics reveals skin surface lipid abnormality in acne in young men. *Br J Dermatol.* 2018;179(3):732–740. doi:10.1111/bjd.16655
9. Zhou M, Yang M, Zheng Y, et al. Skin surface lipidomics revealed the correlation between lipidomic profile and grade in adolescent acne. *J Cosmet Dermatol.* 2020;19(12):3349–3356. doi:10.1111/jocd.13374
10. Scieszka D, Lin YH, Li W, Choudhury S, Yu Y, Freire M. NETome: a model to decode the human genome and proteome of neutrophil extracellular traps. *Sci Data.* 2022;9(1):702. doi:10.1038/s41597-022-01798-1
11. Almaraz-Arreortua A, Sosa-Luis SA, Ríos-Ríos WJ, et al. Morphological and compositional analysis of neutrophil extracellular traps induced by microbial and chemical stimuli. *J Vis Exp.* 2022; 189. doi:10.3791/64522
12. Vorobjeva N, Dagil Y, Pashenkov M, Pinegin B, Chernyak B. Protein kinase C isoforms mediate the formation of neutrophil extracellular traps. *Int Immunopharmacol.* 2022;114:109448. doi:10.1016/j.intimp.2022.109448
13. Chen H, Xu X, Tang Q, et al. (+)-Borneol inhibits the generation of reactive oxygen species and neutrophil extracellular traps induced by phorbol-12-myristate-13-acetate. *Front Pharmacol.* 2022;13:1023450. doi:10.3389/fphar.2022.1023450
14. Kobporchai P, Reamtong O, Phuphisut O, Malaitong P, Adisakwattana P. Serine protease inhibitor derived from *Trichinella spiralis* (TsSERP) inhibits neutrophil elastase and impairs human neutrophil functions. *Front Cell Infect Microbiol.* 2022;12:919835. doi:10.3389/fcimb.2022.919835
15. Pérez-Figueroa E, Álvarez-Carrasco P, Ortega E. Crosslinking of membrane CD13 in human neutrophils mediates phagocytosis and production of reactive oxygen species, neutrophil extracellular traps and proinflammatory cytokines. *Front Immunol.* 2022;13:994496. doi:10.3389/fimmu.2022.994496
16. Hirschfeld J, Chicca JJ, Moonen C, et al. Characterization, quantification, and visualization of neutrophil extracellular traps. *Methods Mol Biol.* 2023;2588:451–472.
17. Artigas-Jerónimo S, González-García A, de la Fuente J, et al. Low NETosis induced in anaplasma phagocytophilum-infected cells. *Vaccines.* 2022;10(10):1756. doi:10.3390/vaccines10101756
18. Saito S, Cao DY, Okuno A, et al. Creatine supplementation enhances immunological function of neutrophils by increasing cellular adenosine triphosphate. *Biosci Microbiota Food Health.* 2022;41(4):185–194. doi:10.12938/bmhf.2022-018
19. Yao H, Cao G, Liu Z, et al. Inhibition of netosis with PAD inhibitor attenuates endotoxin shock induced systemic inflammation. *Int J Mol Sci.* 2022;23(21):13264. doi:10.3390/ijms232113264
20. Yang S, Feng Y, Chen L, et al. Disulfiram accelerates diabetic foot ulcer healing by blocking NET formation via suppressing the NLRP3/Caspase-1/GSDMD pathway. *Transl Res.* 2022;254:115–127.
21. Yang L, Shou YH, Yang YS, Xu JH, Fang D. Elucidating the immune infiltration in acne and its comparison with rosacea by integrated bioinformatics analysis. *PLoS One.* 2021;16(3):e0248650. doi:10.1371/journal.pone.0248650
22. Watanabe K, Yoshino T, Takahashi M, et al. Clinical significance of neutrophil gelatinase-associated lipocalin and galectin-7 in tape-stripped stratum corneum of acne vulgaris. *J Dermatol.* 2018;45(5):618–621. doi:10.1111/1346-8138.14261
23. Jaber SA, Cohen A, D'Souza C, et al. Lipocalin-2: structure, function, distribution and role in metabolic disorders. *Biomed Pharmacother.* 2021;142:112002. doi:10.1016/j.biopha.2021.112002
24. Morales-Valencia J, Lau L, Martí-Nin T, Ozerdem U, David G. Therapy-induced senescence promotes breast cancer cells plasticity by inducing Lipocalin-2 expression. *Oncogene.* 2022;41(38):4361–4370. doi:10.1038/s41388-022-02433-4
25. Huang Y, Zhang N, Xie C, et al. Lipocalin-2 in neutrophils induces ferroptosis in septic cardiac dysfunction via increasing labile iron pool of cardiomyocytes. *Front Cardiovasc Med.* 2022;9:922534. doi:10.3389/fcvm.2022.922534
26. Li Q, Ru X, Yang Y, et al. Lipocalin-2-mediated insufficient oligodendrocyte progenitor cell remyelination for white matter injury after subarachnoid hemorrhage via SLC22A17 receptor/early growth response protein 1 signaling. *Neurosci Bull.* 2022;38(12):1457–1475. doi:10.1007/s12264-022-00906-w
27. Kim HY, Yoo YH. Recombinant FGF21 attenuates polychlorinated biphenyl-induced NAFLD/NASH by modulating hepatic lipocalin-2 expression. *Int J Mol Sci.* 2022;23(16):8899.
28. Shin HJ, Jin Z, An HS, et al. Lipocalin-2 deficiency reduces hepatic and hippocampal triggering receptor expressed on myeloid cells-2 expressions in high-fat diet/streptozotocin-induced diabetic mice. *Brain Sci.* 2022;12(7):878. doi:10.3390/brainsci12070878
29. Gan J, Zheng Y, Yu Q, et al. Serum lipocalin-2 levels are increased and independently associated with early-stage renal damage and carotid atherosclerotic plaque in patients with T2DM. *Front Endocrinol.* 2022;13:855616. doi:10.3389/fendo.2022.855616
30. Betten R, Scharner B, Probst S, et al. Tonicity inversely modulates lipocalin-2 (Lcn2/24p3/NGAL) receptor (SLC22A17) and Lcn2 expression via Wnt/β-catenin signaling in renal inner medullary collecting duct cells: implications for cell fate and bacterial infection. *Cell Commun Signal.* 2018;16(1):74. doi:10.1186/s12964-018-0285-3
31. Ghosh S, Shang P, Yazdankhah M, et al. Activating the AKT2-nuclear factor-κB-lipocalin-2 axis elicits an inflammatory response in age-related macular degeneration. *J Pathol.* 2017;241(5):583–588. doi:10.1002/path.4870
32. Eilenberg W, Stojkovic S, Piechota-Polanczyk A, et al. Neutrophil Gelatinase-Associated Lipocalin (NGAL) is associated with symptomatic carotid atherosclerosis and drives pro-inflammatory state in vitro. *Eur J Vasc Endovasc Surg.* 2016;51(5):623–631. doi:10.1016/j.ejvs.2016.01.009
33. K L, M D S S, H M S. Action of BjussuMP-II, a snake venom metalloproteinase isolated from Bothrops jararacussu venom, on human neutrophils. *Toxicon.* 2022;222:106992. doi:10.1016/j.toxicon.2022.106992
34. Bonneau S, Landry C, Bégin S et al. Correlation between Neutrophil Extracellular Traps (NETs) expression and primary graft dysfunction following human lung transplantation. *Cells.* 2022;11(21):3420. doi:10.3390/cells11213420
35. Mistry P, Carmona-Rivera C, Ombrello AK, et al. Dysregulated neutrophil responses and neutrophil extracellular trap formation and degradation in PAPA syndrome. *Ann Rheum Dis.* 2018;77(12):1825–1833. doi:10.1136/annrheumdis-2018-213746

36. Deng X, Wang JW, Wu Q, Pan L, Mou T, Du CY. Lipocalin 2 as a useful biomarker for risk stratification in patients with acute cholangitis: a single-center prospective and observational study. *Clin Chim Acta*. 2022;533:22–30. doi:10.1016/j.cca.2022.05.022
37. Yadav SK, Ito N, Mindur JE, et al. Fecal Lcn-2 level is a sensitive biological indicator for gut dysbiosis and intestinal inflammation in multiple sclerosis. *Front Immunol*. 2022;13:1015372. doi:10.3389/fimmu.2022.1015372
38. Deis J, Lin TY, Bushman T, Chen X. Lipocalin 2 deficiency alters prostaglandin biosynthesis and mTOR signaling regulation of thermogenesis and lipid metabolism in adipocytes. *Cells*. 2022;11(9):1535. doi:10.3390/cells11091535
39. Moroni F, Ammirati E, Norata GD, Magnoni M, Camici PG. The role of monocytes and macrophages in human atherosclerosis, plaque neoangiogenesis, and atherothrombosis. *Mediators Inflamm*. 2019;2019:7434376. doi:10.1155/2019/7434376
40. Gordon S. Alternative activation of macrophages. *Nat Rev Immunol*. 2003;3(1):23–35. doi:10.1038/nri978
41. Ozlu E, Karadag AS, Ozkanli S, et al. Comparison of TLR-2, TLR-4, and antimicrobial peptide levels in different lesions of acne vulgaris. *Cutan Ocul Toxicol*. 2016;35(4):300–309. doi:10.3109/15569527.2015.1120742
42. Kawasaki T, Kawai T. Toll-like receptor signaling pathways. *Front Immunol*. 2014;5:461. doi:10.3389/fimmu.2014.00461
43. Harder J, Tsuruta D, Murakami M, Kurokawa I. What is the role of antimicrobial peptides (AMP) in acne vulgaris. *Exp Dermatol*. 2013;22(6):386–391. doi:10.1111/exd.12159
44. Owen AM, Luan L, Burelbach KR, et al. MyD88-dependent signaling drives toll-like receptor-induced trained immunity in macrophages. *Front Immunol*. 2022;13:1044662. doi:10.3389/fimmu.2022.1044662
45. Yeudall S, Upchurch CM, Seegren PV, et al. Macrophage acetyl-CoA carboxylase regulates acute inflammation through control of glucose and lipid metabolism. *Sci Adv*. 2022;8(47):eabq1984. doi:10.1126/sciadv.abq1984
46. Chuang LT, Huang WC, Hou YC, et al. Suppressive effect of two cucurbitane-type triterpenoids from momordica charantia on cutibacterium acnes-induced inflammatory responses in human THP-1 monocytic cell and mouse models. *Molecules*. 2021;26(3):579. doi:10.3390/molecules26030579
47. Fang F, Xie Z, Quan J, Wei X, Wang L, Yang L. Baicalin suppresses Propionibacterium acnes-induced skin inflammation by downregulating the NF- κ B/MAPK signaling pathway and inhibiting activation of NLRP3 inflammasome. *Braz J Med Biol Res*. 2020;53(12):e9949. doi:10.1590/1414-431x20209949
48. Chen Y, Cao P, Xiao Z, Ruan Z. m(6)A methyltransferase METTL3 relieves cognitive impairment of hyperuricemia mice via inactivating MyD88/NF- κ B pathway mediated NLRP3-ASC-Caspase1 inflammasome. *Int Immunopharmacol*. 2022;113:109375. doi:10.1016/j.intimp.2022.109375
49. Hayakawa S, Tamura A, Nikiforov N, et al. Activated cholesterol metabolism is integral for innate macrophage responses by amplifying Myd88 signaling. *JCI Insight*. 2022;7(22). doi:10.1172/jci.insight.138539
50. Tang L, Kuang C, Shan D, et al. The ethanol extract of Edgeworthia gardneri (Wall.) Meisn attenuates macrophage foam cell formation and atherogenesis in ApoE(-/-) mice. *Front Cardiovasc Med*. 2022;9:1023438. doi:10.3389/fcvm.2022.1023438
51. Feng J, Li L, Ou Z, et al. IL-25 stimulates M2 macrophage polarization and thereby promotes mitochondrial respiratory capacity and lipolysis in adipose tissues against obesity. *Cell Mol Immunol*. 2018;15(5):493–505. doi:10.1038/cmi.2016.71
52. Huang SC, Everts B, Ivanova Y, et al. Cell-intrinsic lysosomal lipolysis is essential for alternative activation of macrophages. *Nat Immunol*. 2014;15(9):846–855. doi:10.1038/ni.2956
53. Profumo E, Maggi E, Aresse M, et al. Neuropeptide Y promotes human M2 macrophage polarization and enhances p62/SQSTM1-dependent autophagy and NRF2 activation. *Int J Mol Sci*. 2022;23(21):13009. doi:10.3390/ijms232113009
54. Elsaie ML, Aly DG. The immunogenetics of acne. *Adv Exp Med Biol*. 2022;1367:137–154.
55. Freitas Filho EG, Jaca L, Baeza LC, et al. Proteomic analysis of lipid rafts from RBL-2H3 mast cells. *Int J Mol Sci*. 2019;20(16):3904. doi:10.3390/ijms20163904
56. Zhang W, Xu L, Zhu L, Liu Y, Yang S, Zhao M. Lipid droplets, the central hub integrating cell metabolism and the immune system. *Front Physiol*. 2021;12:746749. doi:10.3389/fphys.2021.746749
57. Li J, Li L, Liu R, et al. Integrative lipidomic features identify plasma lipid signatures in chronic urticaria. *Front Immunol*. 2022;13:933312. doi:10.3389/fimmu.2022.933312
58. Taketomi Y, Endo Y, Higashi T, et al. Mast Cell-specific deletion of group III secreted phospholipase A(2) impairs mast cell maturation and functions. *Cells*. 2021;10(7):1691. doi:10.3390/cells10071691
59. Petrosino S, Schiano Moriello A, Verde R, et al. Palmitoylethanolamide counteracts substance P-induced mast cell activation in vitro by stimulating diacylglycerol lipase activity. *J Neuroinflammation*. 2019;16(1):274. doi:10.1186/s12974-019-1671-5
60. Branche E, Wang YT, Viramontes KM, et al. SREBP2-dependent lipid gene transcription enhances the infection of human dendritic cells by Zika virus. *Nat Commun*. 2022;13(1):5341. doi:10.1038/s41467-022-33041-1
61. Veglia F, Tyurin VA, Mohammadyani D, et al. Lipid bodies containing oxidatively truncated lipids block antigen cross-presentation by dendritic cells in cancer. *Nat Commun*. 2017;8(1):2122. doi:10.1038/s41467-017-02186-9
62. Chen IC, Awasthi D, Hsu CL, et al. High-fat diet-induced obesity alters dendritic cell homeostasis by enhancing mitochondrial fatty acid oxidation. *J Immunol*. 2022;209(1):69–76. doi:10.4049/jimmunol.2100567
63. Sardana K, Verma G. Propionibacterium acnes and the Th1/Th17 axis, implications in acne pathogenesis and treatment. *Indian J Dermatol*. 2017;62(4):392–394. doi:10.4103/ijdr.IJD_483_16
64. Antiga E, Verdelli A, Bonciani D, Bonciolini V, Caproni M, Fabbri P. Acne: a new model of immune-mediated chronic inflammatory skin disease. *G Ital Dermatol Venereol*. 2015;150(2):247–254.
65. Jiang J, Meng S, Li L, Duan X, Xu H, Li S. Correlation of acetyl-coenzyme A carboxylase 1 with Th17 and Th1 cells, serving as a potential prognostic biomarker for acute ischemic stroke patients. *J Clin Lab Anal*. 2022;36(10):e24607. doi:10.1002/jcla.24607
66. Komai-Koma M, Ji Y, Cao H, Liu Z, McSharry C, Xu D. Monophosphoryl lipid A directly regulates Th1 cytokine production in human CD4(+) T-cells through Toll-like receptor 2 and 4. *Immunobiology*. 2021;226(5):152132. doi:10.1016/j.imbio.2021.152132
67. Derada Troleiti C, Enzmänn V, Chiurchiu V, et al. Pro-resolving lipid mediator lipoxin A(4) attenuates neuro-inflammation by modulating T cell responses and modifies the spinal cord lipidome. *Cell Rep*. 2021;35(9):109201. doi:10.1016/j.celrep.2021.109201
68. Berod L, Friedrich C, Nandan A, et al. De novo fatty acid synthesis controls the fate between regulatory T and T helper 17 cells. *Nat Med*. 2014;20(11):1327–1333. doi:10.1038/nm.3704
69. Kim W, Barhoumi R, McMurray DN, Chapkin RS. Dietary fish oil and DHA down-regulate antigen-activated CD4 + T-cells while promoting the formation of liquid-ordered mesodomains. *Br J Nutr*. 2014;111(2):254–260. doi:10.1017/S0007114513002444

70. Cai F, Jin S, Chen G, Dobrzyn A. The effect of lipid metabolism on CD4(+) T cells. *Mediators Inflamm.* 2021;2021:6634532. doi:10.1155/2021/6634532
71. Bai A, Guo Y. Acid sphingomyelinase mediates human CD4(+) T-cell signaling: potential roles in T-cell responses and diseases. *Cell Death Dis.* 2017;8(7):e2963. doi:10.1038/cddis.2017.360
72. Kue CS, Lim HX, Jung MY, Hong HJ, Cho D, Kim TS. C6-ceramide in combination with transforming growth factor- β enhances Treg cell differentiation and stable FoxP3 expression in vitro and in vivo. *Immunobiology.* 2013;218(7):952–959. doi:10.1016/j.imbio.2012.11.003
73. Nicholas DA, Proctor EA, Agrawal M, et al. Fatty acid metabolites combine with reduced β oxidation to activate Th17 inflammation in human type 2 diabetes. *Cell Metab.* 2019;30(3):447–461.e5. doi:10.1016/j.cmet.2019.07.004
74. Iwata S, Zhang M, Hao H, et al. Enhanced fatty acid synthesis leads to subset imbalance and IFN- γ overproduction in T helper 1 cells. *Front Immunol.* 2020;11:593103. doi:10.3389/fimmu.2020.593103
75. Carlavan I, Bertino B, Rivier M, et al. Atrophic scar formation in patients with acne involves long-acting immune responses with plasma cells and alteration of sebaceous glands. *Br J Dermatol.* 2018;179(4):906–917. doi:10.1111/bjd.16680
76. Chancheewa B, Asawanonda P, Noppakun N, Kumtornrut C. Myofibroblasts, B Cells, and Mast Cells in Different Types of Long-Standing Acne Scars. *Skin Appendage Disord.* 2022;8(6):469–475. doi:10.1159/000524566
77. Gurzell EA, Teague H, Harris M, Clinthorne J, Shaikh SR, Fenton JI. DHA-enriched fish oil targets B cell lipid microdomains and enhances ex vivo and in vivo B cell function. *J Leukoc Biol.* 2013;93(4):463–470. doi:10.1189/jlb.0812394
78. Kawasaki K, Abe M, Tada F, et al. Blockade of B-cell-activating factor signaling enhances hepatic steatosis induced by a high-fat diet and improves insulin sensitivity. *Lab Invest.* 2013;93(3):311–321. doi:10.1038/labinvest.2012.176
79. Guillamat-Prats R, Hering D, Derle A, et al. GPR55 in B cells limits atherosclerosis development and regulates plasma cell maturation. *Nat Cardiovasc Res.* 2022;1(11):1056–1071. doi:10.1038/s44161-022-00155-0
80. Layton AM, Ravenscroft J. Adolescent acne vulgaris: current and emerging treatments. *Lancet Child Adolesc Health.* 2022;7(2):136–144. doi:10.1016/S2352-4642(22)00314-5
81. Abdelwahab AA, Omar G, Hamdino M. A combined subcision approach with either fractional CO(2) laser (10,600 nm) or cross-linked hyaluronic acid versus subcision alone in atrophic post-acne scar treatment. *Lasers Med Sci.* 2022;38(1):20. doi:10.1007/s10103-022-03677-y
82. Başak PY, Cetin ES, Gürses I, Özseven AG. The effects of systemic isotretinoin and antibiotic therapy on the microbial floras in patients with acne vulgaris. *J Eur Acad Dermatol Venereol.* 2013;27(3):332–336. doi:10.1111/j.1468-3083.2011.04397.x
83. Leheta TM, Abdel Hay RM, El Gareem YF. Deep peeling using phenol versus percutaneous collagen induction combined with trichloroacetic acid 20% in atrophic post-acne scars; a randomized controlled trial. *J Dermatolog Treat.* 2014;25(2):130–136. doi:10.3109/09546634.2012.674192
84. Leheta TM, Abdel Hay RM, Hegazy RA, El Gareem YF. Do combined alternating sessions of 1540 nm nonablative fractional laser and percutaneous collagen induction with trichloroacetic acid 20% show better results than each individual modality in the treatment of atrophic acne scars? A randomized controlled trial. *J Dermatolog Treat.* 2014;25(2):137–141. doi:10.3109/09546634.2012.698249
85. Baroni A, Ruocco E, Russo T, et al. The use of traditional Chinese medicine in some dermatologic diseases: part I--Acne, psoriasis, and atopic dermatitis. *Skinmed.* 2015;13(1):32–38; quiz 39.

Clinical, Cosmetic and Investigational Dermatology

Dovepress

Publish your work in this journal

Clinical, Cosmetic and Investigational Dermatology is an international, peer-reviewed, open access, online journal that focuses on the latest clinical and experimental research in all aspects of skin disease and cosmetic interventions. This journal is indexed on CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-cosmetic-and-investigational-dermatology-journal>