Hepatic Medicine: Evidence and Research downloaded from https://www.dovepress.com/ For personal use only.

## Erratum

Xiao N, Shi S, Zhuang H. A meta-analysis that compares the use of either peginterferon-α2a or peginterferon-α2b plus ribavirin for HCV infection. *Hepatic Medicine: Evidence and Research.* 2010;2:99–109.

The following errors have been published in this work:

Abstract results: The confidence intervals for early virological response (EVR) should be 1.06–1.19, not 1.60–1.19.

Introduction: The sentence beginning, 'Such parameters include rapid response rate...' should read, 'Such parameters include rapid response rate (RVR; HCV-RNA negative at week 4 of therapy),...'

Viral Response: The first sentence should read, 'Compared with PEG-IFN- $\alpha$ 2b plus ribavirin, PEG-IFN- $\alpha$ 2a plus ribavirin led to a higher EVR (RR = 1.12, 95% CI: 1.06–1.19, n = 6 trials; Figure 2A), a higher ETR (RR = 1.21, 95% CI: 1.14–1.28, n = 4 trials; Figure 2B), and a higher level of SVR (RR = 1.10, 95% CI: 1.02–1.18, n = 5 trials; Figure 2C).'

Adverse events: The last sentence of the first paragraph should read, 'Therefore, compared with the PEG-IFN- $\alpha$ 2b group, the risk ratio of the discontinuation of treatment for safety reasons in the PEG-IFN- $\alpha$ 2a group was 0.85 (95% CI: 0.52 to 1.38, n = 6; Figure 4).'

Discussion: The first sentence should be replaced with the following two sentences, 'In this meta-analysis, we demonstrated the regimen of PEG-IFN- $\alpha$ 2a plus ribavirin led to slightly better rates of EVR, ETR, and SVR compared with the use of PEG-IFN- $\alpha$ 2b plus ribavirin in patients with HCV infections.'

Discussion: In the second sentence of the sixth paragraph, two, not three trials reported adequate allocation sequence generation.

Discussion: The second-to-last sentence of the fourthto-last paragraph should read, 'When assessing the SVR, it is not so obvious.

Discussion: The first sentence of the third-to-last paragraph should read, 'Adverse events were observed less frequently among patients treated with PEG-IFN- $\alpha$ 2a, though the numbers of patients that suffered from anemia, depression and rash were not significantly different in this treatment group'.

The legend for Table 1 should read: 'The basic characteristics of seven trials included in this study (1)'.

The legend for Table 2 should read: 'The basic characteristics of seven trials included in this study (2)'.

In Figure 1, the numbers 5, 12 and 469 should be replaced with 6, 13 and 468, respectively.

## Hepatic Medicine: Evidence and Research

## Publish your work in this journal

Hepatic Medicine: Evidence and Research is an international, peerreviewed, open access journal covering all aspects of adult and pediatric hepatology in the clinic and laboratory including the following topics: Pathology, pathophysiology of hepatic disease; Investigation and treatment of hepatic disease; Pharmacology of drugs used for the treatment of hepatic disease. Issues of patient safety and quality of care will also be considered. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: http://www.dovepress.com/hepatic-medicine-evidence-and-research-journal