

A Systematic review on Prevalence, Serotypes and Antibiotic resistance of *Salmonella* in Ethiopia, 2010–2022

Atsebaha Gebrekidan Kahsay¹, Tsehay Asmelash Dejene¹, Enquebaher Kassaye²

¹Department of Medical Microbiology and Immunology, Mekelle University, Mekelle, Tigray, Ethiopia; ²Department of Food Safety and Veterinary Microbiology, Mekelle University, Mekelle, Tigray, Ethiopia

Correspondence: Atsebaha Gebrekidan Kahsay, Department of Medical Microbiology and Immunology, Mekelle University, P. O. Box: 1871, Mekelle, Tigray, Ethiopia, Email atsebaha.gebrekidan@mu.edu.et

Background: In Ethiopia, salmonellosis is one of the most common zoonotic and foodborne illnesses. Ethiopia continues to be at risk for its fast-expanding medication resistance. For the development of preventative and control methods, summarized knowledge regarding salmonellosis is necessary. Determining a thorough evaluation of the prevalence, serotypes, and antibiotic resistance of *Salmonella* in humans and animals from January 1, 2010, to December 30, 2022, in Ethiopia was our goal.

Methods: To find *Salmonella* related articles that published in English, we used the Google Scholar and PubMed search engines. Three researchers conducted the eligible studies using the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist, making sure to include the necessary keywords. If studies were duplicates, incomplete publications, or reported without an antimicrobial test were excluded. Excel 2013 was used to calculate frequencies and tabulate data.

Results: There were a total of 43 investigations from food handlers, diarrhoeic patients, and animals. The prevalence rates ranged from 1% to 10% and 1% to 13% among food handlers and diarrhoea patients, respectively. The highest prevalence was among pigs (41.6%). *S. Anatum* in animals and *S. Typhimurium* in people were the predominant serotypes. Amoxicillin and ampicillin were claimed to be 100% resistant in human studies. The highest recorded resistances for ceftriaxone and ciprofloxacin were 16.7% and 100%, respectively. Animal studies revealed that *Salmonella* resistances to ampicillin, streptomycin and tetracycline were 100%, 90%, 86.4%, respectively. *S. Kentucky* showed complete resistance to tetracycline, ampicillin, gentamicin, ciprofloxacin, and streptomycin.

Conclusion: The prevalence of *Salmonella* among asymptomatic food handlers, diarrheal patients and animals were high in Ethiopia. *S. Typhimurium* that have the zoonotic importance was presented predominantly in human study. High levels of resistances were showed to tetracycline, ampicillin and streptomycin in animal studies. Salmonellosis prevention and control techniques should be strengthened.

Keywords: antibiotic resistance, prevalence, systematic review, serotypes, *Salmonella*, Ethiopia

Introduction

Salmonellosis continues to be a serious global public health issue, particularly in developing nations.^{1,2} One-fourth of 550 million diarrheic people worldwide are thought to contract *Salmonella* each year as a result of eating contaminated food.³ Human typhoid fever is caused by *Salmonella* Typhi and *Salmonella* Paratyphi.⁴ Some *Salmonella* are specific to animal species although others have wide variety of animals, in addition to humans, such as *Salmonella* Typhimurium and *Salmonella* Enteritidis.⁵

Red meat, carcasses, slaughterhouse equipment, and utensils can all get contaminated by the mesenteric lymph nodes and faeces of sick cattle, swine, sheep, or reservoir animals.^{6–10} As a result, eating raw or undercooked meat or eggs results in human infection.¹¹ *Salmonella* infections can also be carried by human and animal faeces, which can contaminate crops.¹² In addition, hospitalization has been implicated as a source of invasive non-typhoidal

Salmonella.¹³ Salmonella infections can lead to outbreaks of human salmonellosis and a variety of clinical symptoms, such as mild gastroenteritis, bacteremia, and extra-intestinal localized infections affecting numerous organs.¹

One of the present and expanding global risks to public health is Salmonella drug resistance. According to a study from Mexico, non-typhoidal Salmonella had a resistance range of 16.9% to 40.3% to the antibiotics trimethoprim-sulfamethoxazole, amoxicillin-clavulanic acid, chloramphenicol, and tetracycline.¹⁴ Similar to this, the *Salmonella* serotypes found in food and people in Italy were resistant to tetracycline (48%) and ampicillin (45%),¹⁵ and from China, the NTS resistance pattern to ceftriaxone was 37%.¹⁶ Additionally, 36,000 and 33,000 cases of salmonella infections resistant to ciprofloxacin and ceftriaxone, respectively, were reported annually from the United States.¹⁷ Salmonella serotypes resistant to ceftriaxone have also been reported in Kenya.¹⁸

According to the prior data, an article on *Salmonella* in Ethiopia had been published since 1985,¹⁹ and in 1994, Salmonella Newport had been discovered in suspected food poisoning cases among students and food handlers in Gondar, Ethiopia.²⁰ From various regions in Ethiopia, a lot of papers about *Salmonella* have been published. These researches looked at the frequency of different Salmonella species, serotypes, and medication resistance patterns in people, animals, and vegetables. Numerous *Salmonella* isolates with high levels of antibiotic resistance have been discovered.^{21–24}

Ethiopia had vast livestock animals in Africa, and living alongside people is prevalent there.²⁵ This may cause the spread of zoonotic illnesses like non-typhoidal salmonella²⁶ as individuals in Ethiopia do not fully comprehend how zoonoses, such as *Salmonella* are spreading.²⁷ As a result, large number of vulnerable individuals like malnourished children may be exposed to the invasive form of non-typhoidal *Salmonella*.^{28,29}

Despite the fragmented local studies about *Salmonella* that have been reported from various regions of Ethiopia, concise information from a systematic review is more helpful for scientific users to identify gaps for additional studies and for policymakers to develop prevention and control strategies based on the scientific information provided. Thus, we sought to conduct a systematic evaluation of the prevalence, serotypes, and antibiotic resistance of Salmonella in people, animals, and their products in Ethiopia from January 1, 2010, to December 30, 2022.

Methods

Literature Search

To find English-language publications concerning *Salmonella* that had already been published, we used the Google Scholar and PubMed search engines. Keywords like “prevalence”, “incidence”, “Salmonella”, “Salmonella serotypes”, “food handlers”, “diarrheic patients”, “antibiotic susceptibility”, “antimicrobial susceptibility”, “antibiotic resistance”, “antimicrobial resistance”, “animals”, “humans”, and “Ethiopia” were used to search the included studies.

The Inclusion and Exclusion Standards

Three researchers independently conducted the eligible studies using the PRISMA³⁰ checklist to make sure all pertinent data was included. If a study met the requirements listed below, it was qualified for the systematic review: *Salmonella* resistance and prevalence in animals and humans are mentioned in the purpose, which is also stated in the design, sample size, description of the microbiological procedures, and number of isolates. The study was published in English. Studies were disregarded if they were duplicates, partial papers, or reported Salmonella without an antimicrobial test.

Obtaining and Analyzing Data

From each examined paper, the names of the authors, the publication year, the kind of media, the study setting, the sampling techniques, the study populations, the type of specimens, the sample size, the number of positive isolates, the antimicrobial susceptibility tests, and the list of serotypes (if any) were taken. Using Excel 2013, frequencies and percentages were examined.

Results

Between January 1, 2010, and December 30, 2022, the search engines Google Scholar and PubMed turned up a total of 207 published papers; 100 of those articles were disregarded from additional examination owing to duplication. After reviewing the 107 research titles and abstracts, it was determined that 52 did not meet the requirements for inclusion.




Figure 1 Flow Diagram for Selection of Eligible Studies. Adapted from Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.* 2009;6(7):e1000097. Creative Commons.³⁰

Only 55 articles that met the requirements for eligibility were given a second evaluation, and 12 of them were disqualified for various reasons. Figure 1 shows that our systematic review only comprised 43 papers.

Characteristics of Studies on *Salmonella* among Asymptomatic Food Handlers and Diarrhoea Patients

Of the 43 eligible studies that were conducted, 28 were asymptomatic food handlers from the community and diarrhoeic patients from public health facilities. While 13 types of research involved asymptomatic food handlers working in university and community cafeterias, 15 studies involved children and adults with diarrhoea. The only clinical specimens reported in any of the investigations for *Salmonella* isolation were faeces. The most frequently employed media were Xylose Lysine Deoxycholate, Deoxycholate Citrate Agar, MacConkey Agar, and *Salmonella* Shigella Agar. According to Table 1, convenience sampling was used in 19 of the investigations, while simple random sampling was used in the other nine.

Salmonella Prevalence in Asymptomatic Food Handlers and Diarrheal Patients

Salmonella prevalence in asymptomatic food handlers (n=13) ranges from 1% to 10%, but it ranges from 1% to 13% in diarrhoeic patients (n=15). More than three-fourths of the reports among asymptomatic food workers had prevalence rates that were less than 5%. On the other hand, in more than two-thirds of the cases, including diarrheal patients, the prevalence rates were 5% or higher as shown in Table 2.

Characteristics of Research on *Salmonella* in Animal-Related Samples

The specimens used to isolate *Salmonella* came from cattle, poultry, pigs, goats, and sheep. These included the contents of the caecum, mesenteric lymph nodes, liver, tongue, hide swabs, rumen, faeces, and cecum, as well as the contents of market eggs and market eggshells. All studies involving animals used the following essential pre-enrichment, enrichment, and selective media: buffered peptone water, brilliant green phenol red lactose sucrose, xylose lysine deoxycholate, MacConkey agar, brilliant green agar, xylose lysine territory 4, selenite cysteine broth, Rappaport-Vassiliadis, and Tetrathionate broth.

Salmonella is present in all of the animal-related specimens, with prevalence ranging from 3.1% in poultry to 29.1% in pig mesenteric lymph nodes. Table 3 lists the results of the investigations that examined the tissue and faeces of pigs, cattle, poultry, sheep, and goats.

Table 1 A Systematic Review of Characteristics of Asymptomatic Food Handlers and Public Health Facility Diarrheic Patients in Ethiopia, January 1, 2010, to December 30, 2022

Authors Name with Reference	Transport & Enrichment Media	Selective and Differential Media	Study Setting	Sampling Method	Study Population	Specimen
Food handlers						
Gebreyesus et al, 2014 ³¹	SFB	XLD	University	Convenience	Food handlers	Stool
Getie et al, 2019 ³²	CTM and SCB	XLD and SSA	Community	SRS	Food handlers	Stool
Abera et al, 2010 ³³	SFB	SSA	Community	SRS	Food handlers	Stool
Solomon et al, 2018 ³⁴	SFB	XLD and BGA	Community	SRS	Food handlers	Stool
Awol et al, 2019 ³⁵	CTM	XLD and MAC	Community	SRS	Food handlers	Stool
Yesigat et al, 2020 ³⁶	CTM and SFB	XLD and MAC	Community	SRS	Food handlers	Stool
Aklilu et al, 2015 ³⁷	SFB	XLD	University	SRS	Food handlers	Stool
Kifelew et al, 2014 ³⁸	SCB	RVB, TTB and XLD	University	SRS	Food handlers	Stool
Getnet et al, 2014 ³⁹	SFB	XLD	University	Convenience	Food handlers	Stool
Mengist et al, 2018 ⁴⁰	RVB	TTB, XLD, BGA	University	Convenience	Food handlers	Stool
Tadesse et al, 2019 ⁴¹	CTM and SFB	XLD and DCA	Community	SRS	Street Food vendor	Stool
Mama and Alemu, 2016 ⁴²	SFB	XLD and MAC	University	Convenience	Food handlers	Stool
Diriba et al, 2020 ⁴³	CTM and SFB	XLD and DCA	University	SRS	Food handlers	Stool
Diarrheic patients						
Tosisa et al, 2020 ⁴⁴	CTM and SFB	XLD, MAC	Healthfacility	Convenience	Diarrheic children	Stool
Abebe et al, 2018 ⁴⁵	CTM and SFB	XLD, DCA	Healthfacility	Convenience	Diarrheic children	Stool
Getamesay et al, 2014 ⁴⁶	CTM and SFB	XLD, DCA	Healthfacility	Convenience	Diarrheic children	Stool
Mamuye et al, 2015 ⁴⁷	CTM	XLD, DCA, SSA	Healthfacility	Convenience	Diarrheic children	Stool
Mulu et al, 2017 ⁴⁸	SFB	MAC, XLD	Healthfacility	Convenience	Diarrheic patients	Stool
Kebede et al, 2017 ⁴⁹	SFB	XLD, MAC	Healthfacility	Convenience	Diarrheic patients	Stool
Amsalu et al, 2021 ⁵⁰	CTM	MAC	Healthfacility	Convenience	Diarrheic patients	Stool
Beyene and Tasew, 2014 ⁵¹	CTM and SFB	XLD, DCA	Healthfacility	Convenience	Diarrheic children	Stool
Eguala et al, 2015 ⁵²	RVB,	XLT 4	Healthfacility	Convenience	Diarrheic patients	Stool
Gebregeziabher, 2018 ⁵³	SFB	MAC, XLD	Healthfacility	Convenience	Diarrheic children	Stool
Terfassa, 2018 ⁵⁴	SFB	XLD, SSA	Healthfacility	Convenience	Diarrheic patients	Stool
Asefa et al, 2019 ⁵⁵	SCB	HEA, SSA	Healthfacility	Convenience	Diarrheic children	Stool
Lamboro et al, 2016 ⁵⁶	SFB	XLD	Healthfacility	Convenience	Diarrheic patients	Stool
Reda et al, 2011 ⁵⁷	SFB	CTM, XLD, DCA	Healthfacility	Convenience	Diarrheic patients	Stool
Ameya et al, 2018 ⁵⁸	Not use	XLD, MAC	Healthfacility	Convenience	Diarrheic children	Stool

Abbreviations: CTM, Cary Blair Transport media; DCA, Deoxycholate Citrate Agar; XLD, xylose Lysine Deoxycholate Agar; Selenite F broth; nr not reported; MAC, MacConkey agar; SSA, Salmonella Shigella agar; RVB, Rappaport-Vassiliadis; TTB, Tetrathionate broth; BGA, brilliant green; SRS, Simple Random sampling.

Table 2 A Systematic Review of the Prevalence of *Salmonella* Among Asymptomatic Food Handlers and Public Health Facilities Diarrheic Patients in Ethiopia from January 1, 2010, to December 30, 2022

Authors name with Reference	Sample size	Isolates	Percent
Food handlers			
Gebreyesus et al, 2014 ³¹	307	3	1
Getie et al, 2019 ³²	257	3	1
Abera et al, 2010 ³³	384	6	1.6
Solomon et al, 2018 ³⁴	387	35	2
Awol et al, 2019 ³⁵	236	5	2
Yesigat et al, 2020 ³⁶	243	6	2
Aklilu et al, 2015 ³⁷	172	6	3
Kifelew et al, 2014 ³⁸	423	13	3
Getnet et al, 2014 ³⁹	233	8	3.4
Mengist et al, 2018 ⁴⁰	220	8	4
Tadesse et al, 2019 ⁴¹	218	13	6
Mama and Alemu, 2016 ⁴²	345	24	7
Diriba et al, 2020 ⁴³	220	21	10
Diarrheic patients			
Tosisa et al, 2020 ⁴⁴	239	3	1
Abebe et al, 2018 ⁴⁵	204	2	1
Getamesay et al, 2014 ⁴⁶	158	4	3
Mamuye et al, 2015 ⁴⁷	253	10	4
Mulu et al, 2017 ⁴⁸	575	24	4
Kebede et al, 2017 ⁴⁹	215	11	5
Amsalu et al, 2021 ⁵⁰	150	8	5.3
Beyene and Tasew, 2014 ⁵¹	260	16	6
Egualé et al, 2015 ⁵²	957	59	6.2
Gebreegziabher, 2018 ⁵³	260	19	7
Terfassa, 2018 ⁵⁴	422	30	7
Asefa et al, 2019 ⁵⁵	422	29	7
Lamboro et al, 2016 ⁵⁶	176	19	11
Reda et al, 2011 ⁵⁷	244	28	11
Ameya et al, 2018 ⁵⁸	167	21	13

Salmonella Antibiotic Resistance Among Food Handlers and Diarrheal Patients

A minimum of nine antibiotics were used to assess the resistance of *Salmonella* isolates. Amoxicillin, ampicillin, tetracycline, trimethoprim-sulfamethoxazole, chloramphenicol, nalidixic acid, gentamicin, ciprofloxacin, and ceftriaxone

Table 3 A Systematic Review of the Prevalence of *Salmonella* in Animals Associated Specimens in Ethiopia from January 1, 2010, to December 30, 2022

Authors Name with Reference	Pre-Enrichment	Enrichment & Selective Media	BT	SAT	Participant	Type of Samples	Sample Size	Isolates	%
Egualé et al, 2016 ²²	BPW	RVB, BPLS, XLD	Yes	Yes	Pig	Caecal content, MLN	556	162	29.1
Dagnew et al, 2020 ⁵⁹	BPW	RVB, XLD	Yes	Yes	Poultry	Cloacal and fecal content	718	22	3.1
Worku et al, 2022 ⁶⁰	BPW	MAC, XLD	Yes	Nr	Cattle	Meat and swab	556	23	4.1
Tadesse et al, 2019 ⁶¹	BPW	SCB, BGA, XLD	Yes	Nr	Egg	Market egg and eggshell	166	7	4.2
Egualé T, 2018 ⁶²	BPW	RVB, BGA, XLD	Yes	Nr	Poultry	Faecal	549	26	4.7
Alemu and Zewde, 2012 ⁶³	BPW	SCB, XLD	Yes	Nr	Cattle	MLN, Intestinal, carcass	558	26	4.7
Mustefa and Gebreedhin, 2018 ⁶⁴	BPW	RVB, TTB, XLT4	Yes	Nr	Cattle	Faecal, MLN	300	17	4.7
Alemu et al, 2022 ⁶⁵	BPW	RVB, BPLS, XLD	Yes	Yes	Goats and sheep	Carcass & skin swabs	345	21	6
Wabeto et al, 2017 ⁶⁶	BPW	RVB, BGA, XLD	Yes	Nr	Cattle	Carcass	896	56	6.3
Addis et al, 2011 ⁶⁷	BPW	RVB, TTB, XLT4	Yes	Nr	Cow	Fecal and milk	390	21	7
Takele et al, 2018 ⁶⁸	BPW	RVB, MAC, XLD	Yes	Nr	Cattle	Caecal and fecal	390	33	8.9
Mohammed and Dubie, 2022 ⁶⁹	BPW	RV	Yes	Nr	Poultry	Cloacal swab	168	21	12.5
Sibhat et al, 2011 ⁷⁰	BPW	RVB, BPLS, XLD	Yes	Yes	Cattle	Hide, rumen, caecal, MLN	400	64	16
Usmael et al, 2022 ⁷¹	BPW	SCB, BGA, XLD	Yes	Nr	Dog	Rectal swab samples	415	26	6.3
Belachew et al, 2021 ⁷²	BPW	RV, SSA, XLD, BGA	Yes	Nr	Chicken	Cloacal & cecum	289	71	24.6

Abbreviations: Nr, Not reported; BPW, Buffered Peptone Water; BPLS, brilliant green phenol red lactose sucrose; XLD, Xylose Lysine Deoxycholate; MAC, MacConkey; BGA, Brilliant Green Agar; XLT4, Xylose Lysine Tergitol 4; SCB, Selenite Cysteine Broth; RVB, Rappaport-Vassiliadis; TTB, Tetrathionate broth; BT, Biochemical tests; SAT, Slide Agglutination Test; MLN, Mesenteric Lymph Node.

were on the lists of antibiotics taken from the studies that qualified. In six investigations for amoxicillin (two from food handlers and four from diarrhoea patients) and eight for ampicillin (four from food handlers and four from diarrhoea patients), all of the isolates displayed 100% resistance. Seven investigations (three in food handlers and four in diarrhoea patients) and eight (two in food handlers and six in diarrhoea patients), respectively, identified resistant isolates for ciprofloxacin and ceftriaxone. Although diarrhoeic patients appear to have relatively significant resistance, Table 4 shows that the highest reported ciprofloxacin and ceftriaxone resistance rates were 16.7% and 100%, respectively. Among all the asymptomatic food handlers and diarrheal patients tested for *Salmonella*, only one study was found eligible for serotyping. The leading serotype that reported from that study was *S. Typhimurium*, whereas single isolate of *S. Concord* showed 100% resistance to amoxicillin, ampicillin, tetracycline, gentamicin, sulfamethoxazole trimethoprim and ciprofloxacin as indicated in Table 4.

Table 4 A Systematic Review of Antibiotic Resistance of *Salmonella* Among Asymptomatic Food Handlers and Public Health Facility Diarrheic Patients in Ethiopia from January 1, 2010 to December 30, 2022

Authors name, Number of isolates and Reference	Antibiotics Resistance, n (%)								
	AMO	AMP	T	SXT	C	NA	CN	CIP	CRO
Food handlers									
Gebreyesus et al (n=3) ³¹	ND	2 (66.7)	2 (66.7)	ND	3 (100)	ND	0	0	ND
Getie et al (n=3) ³²	1 (33.3)	0	2 (66.7)	1 (33.3)	0	0	1 (33.3)	0	0
Abera et al (n=6) ³³	ND	6 (100)	4 (66.7)	5 (83.3)	2 (33.3)	ND	2 (33.3)	ND	ND
Solomon et al (n=35) ³⁴	26 (74.3)	30 (85.7)	ND	24 (68.6)	9 (25.7)	15 (42.9)	ND	5 (14.3)	10 (28.6)
Awol et al (n=5) ³⁵	Nr	3 (60)	4 (80)	2 (40)	2 (40)	1 (20)	ND	0	0
Yesigat et al (n=6) ³⁶	ND	6 (100)	5 (83.3)	1 (16.7)	0	ND	ND	0	ND
Aklilu et al (n=6) ³⁷	6 (100)	6 (100)	0	ND	ND	ND	0	0	ND
Kifelew et al (n=13) ³⁸	6 (46.2)	7 (53.8)	6 (42.3)	2 (15.4)	ND	2 (15.4)	0	ND	0
Getnet et al (n=8) ³⁹	ND	7 (87.5)	1 (12.5)	1 (12.5)	1 (12.5)	Nr	0	1 (12.5)	0
Mengist et al (n=8) ⁴⁰	ND	8 (100)	5 (83.3)	3 (37.5)	2 (25)	ND	0	0	ND
Tadesse et al (n=13) ⁴¹	12 (92.3)	12 (92.3)	10 (77)	2 (15.4)	1 (7.7)	2 (15.4)	2 (15.4)	0	0
Mama and Alemu (n=24) ⁴²	24 (100)	ND	ND	0	0	ND	0	ND	0
Diriba et al (n=21) ⁴³	15 (71.4)	17 (81)	ND	10 (47.6)	15 (71)	ND	ND	5 (23.8)	3 (100)
Diarrheic patients									
Tosisa et al (n=3) ⁴⁴	0	3 (100)	2 (66.7)	2 (66.7)	1 (33.3)	0	0	0	ND
Abebe et al (n=2) ⁴⁵	ND	2 (100)	ND	1 (50)	0	0	2 (100)	0	0
Getamesay et al (n=4) ⁴⁶	0	0	0	0	0	1 (25)	0	0	4 (100)
Mamuye et al (n=10) ⁴⁷	8 (80)	8 (80)	ND	6 (60)	4 (40)	2 (20)	0	0	0
Mulu et al (n=24) ⁴⁸	8 (33.3)	22 (91.7)	0	8 (33.3)	4 (16.7)	ND	ND	4 (16.7)	9 (37.5)
Kebede et al (n=11) ⁴⁹	ND	ND	7 (63.7)	8 (72.7)	8 (72.7)	1 (9)	1 (9)	1 (9)	1 (9)
Amsalu et al (n=8) ⁵⁰	ND	ND	7 (87.5)	7 (87.5)	4 (50)	3 (37.5)	2 (25)	1 (12.5)	2 (25)
Beyene and Tasew (n=16) ⁵¹	16 (100)	10 (62.5)	ND	5 (31.3)	3 (18.8)	2 (12.5)	0	0	0
Egual et al, 2015 ⁵²									
• S. Typhimurium (n=27)	4 (14.8)	6 (22.2)	3 (11.1)	1 (3.7)	1 (3.7)	0 (0)	0 (0)	0 (0)	1 (3.7)
• S. Newport (n=2)	1 (50)	1 (50)	0 (0)	1 (50)	0 (0)	1 (50)	0 (0)	0 (0)	0 (0)
• S. Virchow (n=21)	1 (4.8)	1 (4.8)	3 (14.3)	0 (0)	0 (0)	10 (47.6)	0 (0)	0 (0)	0 (0)
• S. Kottbus (n=7)	0 (0)	0 (0)	1 (14.3)	0 (0)	1 (14.3)	0 (0)	0 (0)	1 (14.3)	0 (0)
• S. Enteritidis (n=1)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
• S. Braenderup (n=1)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
• S. Concord (n=1)	1 (100)	1 (100)	1 (100)	1 (100)	0 (0)	0 (0)	1 (100)	1 (100)	0 (0)
• S. Miami (n=3)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)

(Continued)

Table 4 (Continued).

Authors name, Number of isolates and Reference	Antibiotics Resistance, n (%)								
	AMO	AMP	T	SXT	C	NA	CN	CIP	CRO
• S. Kentucky (n=2)	2 (100)	2 (100)	2 (100)	2 (100)	0 (0)	2 (100)	2 (100)	2 (100)	0 (0)
Gebregziabher et al (n=19) ⁵³	ND	17 (89.5)	17 (89.5)	11 (57.9)	15 (78.9)	6 (31.6)	3 (15.8)	0	2 (10.5)
Terfassa et al (n=30) ⁵⁴	27 (90)	ND	ND	ND	2 (6.6)	1 (3.3)	3 (10)	2 (6.7)	1 (3.3)
Asefa et al (n=29) ⁵⁵	29 (100)	ND	23 (79)	ND	9 (31)	ND	ND	0	0
Lamboro et al (n=19) ⁵⁶	ND	19 (100)	9 (47.6)	1 (5.3)	1 (5.3)	5 (26.3)	1 (5.3)	0	ND
Reda et al (28) ⁵⁷	28 (100)	28 (100)	20 (71.4)	ND	18 (64.3)	ND	1 (3.6)	ND	ND
Ameya et al (21) ⁵⁸	21 (100)	ND	ND	8 (38.1)	9 (42.9)	ND	4 (19)	0	ND

Abbreviations: AMP, Ampicillin; AMO, Amoxicillin; SXT, sulfamethoxazole-trimethoprim; NA, Nalidixic acid; CRO, Ceftriaxone; CIP, Ciprofloxacin; C, Chloramphenicol; CN, Gentamicin; T, Tetracycline; ND, Not Done.

Animal *Salmonella* with Antibiotic Resistance

As shown in Table 5, all *Salmonella* isolates from investigations involving animal-related samples were resistant to ampicillin, 90% to streptomycin, 66.7% to chloramphenicol, 86.4% to tetracycline, 35.2% to ciprofloxacin, 29.4% to gentamicin, and 23.2% to ceftriaxone. When it comes to *salmonella* serotypes' resistance to antibiotics, the ampicillin,

Table 5 A Systematic Review of Antibiotic Resistance of *Salmonella* in Species and Serotype Level in Animals Related Specimens in Ethiopia from January 1, 2010 to December 30, 2022

Authors Name, Number of Isolates and Reference	Antibiotics Resistance, n (%)						
	AMP	GEN	CIP	C	S	T	CRO
Egualle et al, 2016 ²²							
• S. Kentucky (n=5)	5 (100)	5 (100)	5 (100)	0 (0)	5 (100)	5 (100)	0 (0)
• S. Virchow (n=5)	3 (60)	1 (20)	0 (0)	0 (0)	2 (40)	1 (20)	0 (0)
• S. Saintpaul (n=6)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
• S. Typhimurium (n=7)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
• S. Dublin (n=3)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Dagnew et al, 2020 (n=22) ⁵⁹	0 (0)	0 (0)	0 (0)	0 (0)	20 (90.1)	19 (86.4)	0 (0)
Worku et al, 2022 (n=23) ⁶⁰	ND	0 (0)	ND	ND	ND	6 (26)	0 (0)
Egualle 2018 ⁶²							
• S. Saintpaul (n=20)	9 (45)	0 (0)	0 (0)	10 (50)	18 (90)	4 (20)	5 (25)
• S. Typhimurium (n=3)	0 (0)	0 (0)	0 (0)	0 (0)	3 (100)	1 (33.3)	1 (33.3)
• S. Kentucky (n=2)	2 (100)	2 (100)	2 (100)	1 (50)	2 (100)	2 (100)	0 (0)
Alemu and Zewde, 2012 (n=21) ⁶³	1 (4.8)	0 (0)	1 (4.8)	0 (0)	ND	2 (9.5)	0 (0)
Mustefa et al, (n=17) ⁶⁴	17 (100)	5 (29.4)	6 (35.2)	0 (0)	7 (41.1)	ND	0 (0)
Wabeto et al, 2017 (n=56) ⁶⁶	26 (46.4)	7 (12.5)	4 (7.1)	29 (51.8)	37 (66)	47 (83.9)	13 (23.2)

(Continued)

Table 5 (Continued).

Authors Name, Number of Isolates and Reference	Antibiotics Resistance, n (%)						
	AMP	GEN	CIP	C	S	T	CRO
Addis et al, 2011 (n=21) ⁶⁷	21 (100)	4 (19)	0 (0)	2 (9.5)	16 (76.2)	7 (33.3)	ND
Takele et al, 2018 (n=42) ⁶⁸	13 (31)	3 (7)	0 (0)	3 (7)	10 (23.8)	10 (23.8)	0 (0)
Mohammed and Gebremedhin (n=21) ⁶⁹	7 (33.3)	(4.8)	(4.8)	7 (33.3)	13 (61.9)	15 (71)	ND
Sibhat et al, 2011 ⁷⁰							
• S. Typhimurium (n=1)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
• S. Newport (n=13)	0 (0)	0 (0)	0 (0)	0 (0)	4 (66.7)	0 (0)	0 (0)
• S. Anatum (n=39)	2 (66.7)	0 (0)	0 (0)	0 (0)	3 (100)	2 (66.7)	0 (0)
Usmael et al, 2022 ⁷¹	11 (41.7)	0 (0)	ND	ND	ND	9 (21.2)	ND
Belachew et al, 20 (n=71) ⁷²	38 (53.5)	7 (10)	ND	47 (66.7)	43 (60.6)	56 (79)	ND

Abbreviations: AMP, Ampicillin; GEN, Gentamycin; CIP, ciprofloxacin; CAF, Chloramphenicol; S, streptomycin; TTC, tetracycline; CRO, ceftriaxone; ND, Not done.

gentamicin, ciprofloxacin, streptomycin, and tetracycline resistance in *S. Kentucky* was 100% although the leading isolates of salmonella serotypes reported from the researches were *S. Anatum*, *S. Saintpaul*, *S. Newport*, and *S. Typhimurium*.

Discussion

One of the main contributors to foodborne and zoonotic diseases in underdeveloped nations is *Salmonella*.^{1,2} It is one of Ethiopia's top seven priority zoonotic diseases⁷³ and a significant source of foodborne pathogens.⁷⁴

Salmonella prevalence among asymptomatic food handlers ranges from 1% to 10%, which was consistent with research from Pakistan (9.1%).⁷⁵ Similar to this, the prevalence of *Salmonella* among individuals who have diarrhoea ranges from 1% to 13%. However, one instance at a hospital suggests that there are 30 other cases in the community who are unable to visit hospitals for a variety of reasons.⁷⁶

The prevalence rates of *Salmonella* in animal-related sources range from 3.1% in poultry to 29.4% in pigs, followed by cloacal and cecal content of chicken (24.6%) which is consistent with studies from Burkina Faso,^{77,78} Italy,⁷⁹ Kenya,⁸⁰ South Africa,⁸¹ and Uganda,⁸² Vietnam,⁸³ China,⁸⁴ and Louisiana.⁸⁵

S. Typhimurium were reported in all of the legible studies for serotyping which was also reported from Gambia,⁸⁶ Mali,⁸⁷ India,⁸⁸ Mexico,⁸⁹ and China.⁹⁰ According to reports from sub-Saharan African nations, *S. Typhimurium* and *S. Enteritidis* are invasive forms of NTS, especially among susceptible people, such as those with HIV, malnourished children, and in malaria areas.^{77,91} It is possible that the existence of such *Salmonella* germs in cattle, poultry, pigs, and other animal sources is regarded as a potential source of contamination in humans and maybe the main risk factor for *Salmonella* outbreaks in humans.⁸

Six human studies found that every isolate of *Salmonella* tested had a 100% amoxicillin resistance rate, which is consistent with an Ethiopian study.²⁴ Ciprofloxacin-resistant isolates were detected in seven human studies which is similar to studies from China,⁹² and Mexico.⁹³

Salmonella isolates from two investigations had 100% ampicillin resistance, which was higher than the prior study in Japan.⁶ However, the streptomycin resistance patterns were consistent with earlier Japanese studies.⁶ For ciprofloxacin, gentamicin, and ceftriaxone, similar animal specimens showed at least 25% resistance. *S. Kentucky* showed 100% resistance to ampicillin, gentamicin, ciprofloxacin, streptomycin, and tetracycline out of the known serotypes from our systematic evaluation.

The results of our systematic review are very beneficial in helping people understand important information concerning *Salmonella*. We were unable to include *Salmonella* in food, fruits, or vegetables because of limited data. Additionally, no studies that were included in this systematic review performed molecular characterization of *Salmonella* and its resistance genes or invasive non-typhoidal salmonella among malnourished children because of no studies related.

Conclusions

The prevalence of *Salmonella* in animals, diarrhoeal patients, and asymptomatic food handlers was high. The prevalence of *Salmonella* among asymptomatic food handlers and diarrheal patients was nearly similar. *S. Typhimurium* that has the zoonotic nature was recovered from human and animal studies. Studies on animals revealed high levels of resistance to tetracycline, ampicillin, and streptomycin. Circulating of *Salmonella* in the community in Ethiopia is a homework for academic and non-academic researchers to overcome fear of future outbreaks. National and international organizations should work on strengthening the prevention and control of salmonellosis. Due to Ethiopia's high prevalence of underweight children and the prevalence of those who are susceptible to the invasive form of NTS, invasive non-typhoidal salmonella among malnourished children should be examined.

Disclosure

The authors report no conflicts of interest in this work.

References

- Lo HY, Lai FP, Yang YJ. Changes in epidemiology and antimicrobial susceptibility of nontyphoidal *Salmonella* in children in southern Taiwan, 1997–2016. *J Microbiol Immunol Infect*. 2020;53(4):585–591. doi:10.1016/j.jmii.2018.06.004
- Stanaway J, Parisi A, Sarkar K, et al. The global burden of non-typhoidal salmonella invasive disease: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet Infect Dis*. 2019;19(12):1312–1324. doi:10.1016/S1473-3099(19)30418-9
- World Health Organization. *Salmonella* (non-typhoidal); 2018. Available from: <https://www.who.int/news-room/fact-sheets/detail/salmonella>. Accessed October 4, 2023.
- Ikhimiukor O, Oaikhen A, Afolayan AO, et al. Genomic characterization of invasive typhoidal and non-typhoidal *Salmonella* in southwestern Nigeria. Genomic characterization of invasive typhoidal and non-typhoidal *Salmonella* in southwestern Nigeria. *PLoS Negl Trop Dis*. 2022;16(8):e0010716. doi:10.1371/journal.pntd.0010716
- Hawwas HA, Aboueisha AM, Fadel HM, ElMahallawy HS. *Salmonella* serovars in sheep and goats and their probable zoonotic potential to humans in Suez Canal Area. *Egypt Cta Veterinaria Scandinavica*. 2022;64(1):17. doi:10.1186/s13028-022-00637-y
- Duc VM, Nakamoto Y, Fujiwara A, et al. Prevalence of *Salmonella* in broiler chickens in Kagoshima, Japan in 2009 to 2012 and the relationship between serovars changing and antimicrobial resistance. *BMC Vet Res*. 2019;15(1):108. doi:10.1186/s12917-019-1836-6
- Mannion C, June F, McLernon J, et al. The role of transport, lairage and slaughter processes in the dissemination of *Salmonella* spp. in pigs in Ireland. *Food Res Int*. 2012;45(2):871–879. doi:10.1016/j.foodres.2011.02.001
- Hugas M, Beloeil PA. Controlling *Salmonella* along the food chain in the European Union - progress over the last ten years. *Euro Surveill*. 2014;19(19):20804. doi:10.2807/1560-7917.ES2014.19.19.20804
- Boschi-Pinto C, Young M, Black RE. The Child Health Epidemiology Reference Group reviews the effectiveness of interventions to reduce maternal, neonatal and child mortality. *Int J Epidemiol*. 2010;39(Supplement 1):3–6. doi:10.1093/ije/dyq018
- Reza HS, Zohreh M, Amir G. Prevalence and antimicrobial resistance of *Salmonella* serotypes isolated from retail chicken meat and giblets in Iran. *J Infect Dev Countr*. 2015;9(05):463–469. doi:10.3855/jide.5945
- Jacob ST, Moore CC, Banura P, et al. Severe sepsis in two Ugandan hospitals: a prospective observational study of management and outcomes in a predominantly HIV-1 infected population. *PLoS One*. 2009;4(11):7782. doi:10.1371/journal.pone.0007782
- Kowalska B. Fresh vegetables and fruit as a source of *Salmonella* bacteria. *Ann Agric Environ Med*. 2023;30(1):9–14. doi:10.2644/aaem/156765
- Oxman JM, Boadla LR, Sells N. An isolated case of nosocomial acquisition of invasive non-typhoidal *Salmonella*. *IDCases*. 2023;33:01816. doi:10.1016/j.idcr.2023.e01816
- Delgado-Suarez EJ, Guterrez TP, Ruiz-Lo'pez FA, et al. Genomic surveillance of antimicrobial resistance shows cattle and poultry are a moderate source of multi-drug resistant nontyphoidal *Salmonella* in Mexico. *PLoS One*. 2021;16(5):e0243681. doi:10.1371/journal.pone.0243681
- Capuano F, Mancusi A, Capparelli R, et al. Characterization of Drug Resistance and Virulotypes of *Salmonella* Strains Isolated from Food and Humans. *Foodborne Pathog Dis*. 2013;10(11):963–968. doi:10.1089/fpd.2013.1511
- Ke Y, Lu W, Liu W, et al. Non-typhoidal *Salmonella* infections among children in a tertiary hospital in Ningbo, Zhejiang, China, 2012–2019. *PLoS Negl Trop Dis*. 2020;14(10):e0008732. doi:10.1371/journal.pntd.0008732
- Centers for Disease Control and Prevention. Antibiotic Resistance Threats In the United States; 2013.
- Mutai WC, Muigai AW, Waikiki P, Kariuki S. Multi-drug resistant *Salmonella* enterica serovars Typhi isolates with reduced susceptibility to ciprofloxacin in Kenya. *BMC Microbiol*. 2018;18(1):187. doi:10.1186/s12866-018-1332-3
- Gebre-Yohannes A. *Salmonella* from Ethiopia: prevalent species and their susceptibility to drugs. *Ethiopia Med J*. 1985;23(3):97–102.
- Aseffa A, Mengistu G, Tiruneh M. *Salmonella* Newport: outbreak of food poisoning among college students due to contaminated undercooked eggs. *Ethiopia Med J*. 1994;32:1–6.

21. Guchi B, Ashenafi M. Microbial Load, Prevalence and Antibigrams of Salmonella and Shigella in Lettuce and Green Peppers. *Ethiop J Health Sci.* 2010;20:42–48.
22. Eguale T, Engidawork E, Gebreyes WA, et al. Fecal prevalence, serotype distribution and antimicrobial resistance of Salmonellae in dairy cattle in central Ethiopia. *BMC Microbiol.* 2016;16(1):20. doi:10.1186/s12866-016-0638-2
23. Mitiku H, Weldegebreal F, Marami D, Teklemariam Z. Nontyphoidal Salmonella bacteremia in antiretroviral therapy-naïve HIV-infected individuals at three public hospitals in eastern Ethiopia: prevalence, antimicrobial susceptibility patterns, and associated factors. *HIV/AIDS.* 2019;2019:11 23–29.
24. Dessale M, Mengistu G, Mengist HM. Prevalence, antimicrobial resistance pattern, and associated factors of Salmonella and Shigella among under five diarrheic children attending public health facilities in Debre Markos town, Northwest Ethiopia. *Front Public Health.* 2023;11. doi:10.3389/fpubh.2023.1114223
25. Food and Agriculture Organization of the United Nations. *The Future of Livestock in Ethiopia. Opportunities and Challenges in the Face of Uncertainty.* Rome: Food and Agriculture Organization of the United Nations; 2019.
26. Hoelzer K, Andrea Switt AIM, Wiedmann M. Animal contact as a source of human non-typhoidal salmonellosis. *Vet Res.* 2011;42(1):34. doi:10.1186/1297-9716-42-34
27. Alebie A, Tewachew T. Household practice related to zoonotic diseases transmission in rural community of Gondar Zuria District. *Vet Med.* 2021;12:109–115.
28. Mtove G, Amos B, von Seidlein L, et al. Invasive salmonellosis among children admitted to a rural Tanzania hospital and a comparison with previous studies. *PLoS One.* 2010;5(2):e9244. doi:10.1371/journal.pone.0009244
29. Abebe HF, Taffere GR, Fisseha MA, Bezabih AM. Risk factors and spatial distributions of underweight among children under-five in urban and rural communities in Tigray, Northern Ethiopia: using ordinal logistic regression analysis. *Nutr Dietary Suppl.* 2022;14:21–37. doi:10.2147/NDS.S371773
30. Moher D, Liberati A, Tetzlaff J, Altman DG; The PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.* 2009;6(7):e1000097. doi:10.1371/journal.pmed.1000097
31. Gebreyesus A, Adane K, Negash L, et al. Prevalence of Salmonella typhi and intestinal parasites among food handlers in Mekelle University student cafeteria, Mekelle, Ethiopia. *Food Control.* 2014;44:45–48. doi:10.1016/j.foodcont.2014.03.040
32. Getie M, Abebe W, Tessema B. Prevalence of enteric bacteria and their antimicrobial susceptibility patterns among food handlers in Gondar town, Northwest Ethiopia. *Antimicrob Resist Infect Control.* 2019;8(1):111. doi:10.1186/s13756-019-0566-7
33. Abera B, Biadegelgen F, Bezabih B. Prevalence of Salmonella typhi and intestinal parasites among food handlers in Bahir Dar Town, Northwest Ethiopia. *Ethiop J Health Dev.* 2010;24:46–51.
34. Solomon F, Wada FW, Anjulo AA, et al. Burden of intestinal pathogens and associated factors among asymptomatic food handlers in South Ethiopia: emphasis on salmonellosis. *BMC Res Notes.* 2018;11(1):502. doi:10.1186/s13104-018-3610-4
35. Awol N, Nigusse D, Ali M. Prevalence and antimicrobial susceptibility profile of Salmonella and Shigella among food handlers working in food establishment at Hawassa city, Southern Ethiopia. *BMC Res Notes.* 2019;12(1):712. doi:10.1186/s13104-019-4725-y
36. Yesigat T, Jemal M, Birhan W. Prevalence and associated risk factors of Salmonella, Shigella, and intestinal parasites among food handlers in Motta Town, North West Ethiopia. *Can J Infect Dis Med Microbiol.* 2020;2020:1–11. doi:10.1155/2020/6425946
37. Aklilu AK, Dessalegn M, Tarekegn N, et al. Prevalence of intestinal parasites, Salmonella and Shigella among apparently healthy food handlers of Addis Ababa University student's cafeteria, Addis Ababa, Ethiopia. *BMC Res Notes.* 2015;8(1):17. doi:10.1186/s13104-014-0967-x
38. Kiflew GL, Wondafrash N, Feleke A. Identification of drug-resistant Salmonella from food handlers at the University of Gondar, Ethiopia. *BMC Res Notes.* 2014;7(1):545. doi:10.1186/1756-0500-7-545
39. Getnet F, Gebre-Selassie S, Alemayehu H, et al. Prevalence and antimicrobial resistance of salmonella isolated from food handlers in Addis Ababa University Students' Cafeteria, Ethiopia. *Afr J Basic Appl Sci.* 2014;6(6):210–216.
40. Mengist A, Mengistu G, Reta A. Prevalence and antimicrobial susceptibility pattern of Salmonella and Shigella among food handlers in catering establishments at Debre Markos University, Northwest Ethiopia. *Int J Infect Dis.* 2018;75:74–79. doi:10.1016/j.ijid.2018.08.008
41. Tadesse G, Mitiku H, Teklemariam Z, Marami D. Salmonella and shigella among asymptomatic street food vendors in the Dire Dawa city, Eastern Ethiopia: prevalence, antimicrobial susceptibility pattern, and associated factors. *Environ Health Insights.* 2019;13:1–8. doi:10.1177/1178630219853581
42. Mama M, Alemu G. Prevalence, antimicrobial susceptibility patterns and associated risk factors of Shigella and Salmonella among food handlers in Arba Minch University, South Ethiopia. *BMC Infect Dis.* 2016;16(1):686. doi:10.1186/s12879-016-2035-8
43. Diriba K, Awulachew E, Ashuro Z. Prevalence and antimicrobial resistance pattern of Salmonella, Shigella, and intestinal parasites and associated factor among Food Handlers in Dilla University Student Cafeteria, Dilla, Ethiopia. *Int J Microbiol Res.* 2020;2020:1–10. doi:10.1155/2020/3150539
44. Tosisa W, Mihret A, Ararsa A, et al. Prevalence and antimicrobial susceptibility of Salmonella and Shigella species isolated from diarrheic children in Ambo town. *BMC Pediatr.* 2020;20(91). doi:10.1186/s12887-020-1970-0
45. Abebe W, Earsido A, Taye S, et al. Prevalence and antibiotic susceptibility patterns of Shigella and Salmonella among children aged below five years with Diarrhoea attending Nigist Eleni Mohammed Memorial Hospital, South Ethiopia. *BMC Pediatr.* 2018;18(1):241. doi:10.1186/s12887-018-1221-9
46. Getamesay M, Getent B, Ahmed Z. Prevalence of Shigella, Salmonella and Campylobacter species and their susceptibility patterns among under-five children with diarrhoea in Hawassa town, south Ethiopia. *Ethiop J Health Sci.* 2014;24(2):101–108. doi:10.4314/ejhs.v24i2.1
47. Mamuye Y, Metaferia G, Birhanu A, et al. Isolation and antibiotic susceptibility patterns of Shigella and Salmonella among under 5 children with acute diarrhoea: a cross-sectional study at selected public health facilities in Addis Ababa, Ethiopia. *Clin Microbiol.* 2015;4(01). doi:10.4172/2327-5073.1000186
48. Mulu W, Abera B, Yimer M, et al. Bacterial agents and antibiotic resistance profiles of infections from different sites that occurred among patients at Debre Markos Referral Hospital, Ethiopia: a cross-sectional study. *BMC Res Notes.* 2017;10(1):254. doi:10.1186/s13104-017-2584-y
49. Kebede A, Solomon A, Tchalew TS. The common enteric bacterial pathogens and their antimicrobial susceptibility patterns among HIV-infected individuals attending the antiretroviral therapy clinic of Hawassa University Hospital, southern Ethiopia. *Antimicrob Resist Infect Control.* 2017;6(128). doi:10.1186/s13756-017-0288-7

50. Amsalu TG, Adem Siraj Y, Adem Siraj Y. Salmonella Typhi and Salmonella Paratyphi prevalence, antimicrobial susceptibility profile and factors associated with enteric fever infection in Bahir Dar, Ethiopia. *Sci Rep*. 2021;11(1):7359. doi:10.1038/s41598-021-86743-9
51. Beyene G, Tasew H. Prevalence of intestinal parasite, Shigella and Salmonella species among diarrheal children in Jimma health centre, Jimma southwest Ethiopia: a cross-sectional study. *Ann Clin Microbiol Antimicrobiol*. 2014;13(10):1–7.
52. Eguale T, Gebreyes WA, Asrat D, et al. Non-typhoidal Salmonella serotypes, antimicrobial resistance and co-infection with parasites among patients with diarrhoea and other gastrointestinal complaints in Addis Ababa, Ethiopia. *BMC Infect Dis*. 2015;15(1):497. doi:10.1186/s12879-015-1235-y
53. Gebreegziabher G, Asrat D, W/Amanuel Y, Hagos T. Isolation and antimicrobial susceptibility profile of Shigella and Salmonella Species from children with acute diarrhoea in Mekelle Hospital and Semen Health Center, Ethiopia. *Ethiopia J Health Sci*. 2018;28(2):197–206.
54. Terfassa A, Jida M. Prevalence and Antibiotics Susceptibility Pattern of Salmonella and Shigella Species among Diarrheal Patients Attending Nekemte Referral Hospital, Oromia, Ethiopia. *Int J Microbiol*. 2018;2018:9214689. doi:10.1155/2018/9214689
55. Assefa A, Girma M. Prevalence and antimicrobial susceptibility patterns of Salmonella and Shigella isolates among children aged below five years with diarrhoea attending Robe General Hospital and Goba Referral Hospital, South East Ethiopia. *Trop Dis Travel Med Vaccines*. 2019;5:19. doi:10.1186/s40794-019-0096-6
56. Lamboro T, Ketema T, Bacha K. Prevalence and antimicrobial resistance in Salmonella and Shigella species isolated from outpatients, Jimma University Specialized Hospital, Southwest Ethiopia. *Can J Infect Dis Med Microb*. 2016;2016:4210760.
57. Reda A, Seyoum B, Andualem YG, et al. Antibiotic susceptibility patterns of Salmonella and Shigella isolates in Harar, Eastern Ethiopia. *J Infect Dis Immun*. 2011;3(8):134–139.
58. Ameyu G, Tsalla T, Getu F, Getu E. Antimicrobial susceptibility pattern, and associated factors of Salmonella and Shigella infections among under-five children in Arba Minch, South Ethiopia. *Ann Clin Microbiol Antimicrob*. 2018;17(1):1. doi:10.1186/s12941-018-0253-1
59. Dagnew B, Alemayehu H, Medhin G, Eguale T. Prevalence and antimicrobial susceptibility of Salmonella in poultry farms and in-contact humans in Adama and Modjo towns, Ethiopia. *Microbiol Open*. 2020;9(8):e1067. doi:10.1002/mbo3.1067
60. Worku WD, Menjetta T, Menjetta T. High prevalence and antimicrobial susceptibility pattern of salmonella species and extended-spectrum beta-lactamase producing Escherichia coli from raw cattle meat at butcher houses in Hawassa city, Sidama regional state, Ethiopia. *PLoS One*. 2022;17(1):e0262308. doi:10.1371/journal.pone.0262308
61. Taddese D, Tolosa T, Deresa B, et al. Antibigrams and risk factors of Salmonella isolates from laying hens and eggs in Jimma Town, South Western Ethiopia. *BMC Res Notes*. 2019;12(1):472. doi:10.1186/s13104-019-4516-5
62. Eguale T. Non-typhoidal Salmonella serovars in poultry farms in central Ethiopia: prevalence and antimicrobial resistance. *BMC Vet Res*. 2018;14(1):217. doi:10.1186/s12917-018-1539-4
63. Alemu S, Zewde BM. Prevalence and antimicrobial resistance profiles of Salmonella enterica serovars isolated from slaughtered cattle in Bahir Dar, Ethiopia. *Trop Anim Health Prod*. 2012;44(3):595–600. doi:10.1007/s11250-011-9941-y
64. Mustefa BA, Gebremedhin EZ. Carriage and antimicrobial resistance of non-typhoidal Salmonella in cattle slaughtered in Ambo municipality abattoir, West Shewa zone, Oromia, Ethiopia - a point prevalence survey. *Ethiop Vet J*. 2018;22(2):94. doi:10.4314/evj.v22i2.8
65. Alemu A, Regassa F, Kebede N, et al. The magnitude and Antimicrobial Susceptibility Profile of Salmonella Recovered from Export Abattoirs Located in East Shewa, Ethiopia. *Infect Drug Resist*. 2022;15:1353–1365. doi:10.2147/IDR.S348773
66. Wabeto W, Abraham Y, Anjulo AA. Detection and identification of antimicrobial-resistant Salmonella in raw beef at Wolaita Sodo municipal abattoir, Southern Ethiopia. *J Health Popul Nutr*. 2017;36(1):52. doi:10.1186/s41043-017-0131-z
67. Addis Z, Kebede N, Sisay Z, et al. Prevalence and antimicrobial resistance of Salmonella isolated from lactating cows and in contact humans in dairy farms of Addis Ababa: a cross-sectional study. *BMC Infect Dis*. 2011;11(1):222. doi:10.1186/1471-2334-11-222
68. Takele S, Woldemichael K, Gashaw M, et al. Prevalence and drug susceptibility pattern of Salmonella isolates from apparently healthy slaughter cattle and personnel working at the Jimma municipal abattoir, southwest Ethiopia. *Trop Dis Travel Med Vaccines*. 2018;4(1):13. doi:10.1186/s40794-018-0072-6
69. Mohammed Y, Dubie T. Isolation, identification and antimicrobial susceptibility profile of Salmonella isolated from poultry farms in Addis Ababa, Ethiopia. *Vet Med Sci*. 2022;8(3):1166–1173. doi:10.1002/vms3.762
70. Sibhat B, Zewde BM, Zerihun A, et al. Salmonella serovars and antimicrobial resistance profiles in beef cattle, slaughterhouse personnel and slaughterhouse environment in Ethiopia. *Zoonoses Public Health*. 2011;58(2):102–109. doi:10.1111/j.1863-2378.2009.01305.x
71. Usmael B, Abraha B, Alemu S, et al. Isolation, antimicrobial susceptibility patterns, and risk factors assessment of nontyphoidal Salmonella from apparently healthy and diarrheic dogs. *BMC Vet Res*. 2022;18(1):37. doi:10.1186/s12917-021-03135-x
72. Belachew TM, Roba E, Asefa YT, et al. Prevalence and Antimicrobial-Susceptibility Profiles of Salmonella in Smallhold Broiler Supply Chains in Central Ethiopia. *Infect Drug Resist*. 2021;14:4047–4055. doi:10.2147/IDR.S331249
73. Centers for Disease Control and Prevention, Ethiopia Public Health Institute and Ethiopian Ministry of Agriculture. Zoonotic Disease Prioritization for Inter-sectoral Engagement in Ethiopia. Addis Ababa; 2015.
74. Asfaw T, Genetu D, Shenkute D, et al. Foodborne pathogens and antimicrobial resistance in Ethiopia: an urgent call for action on “One Health”. *Infect Drug Resist*. 2022;15:5265–5274. doi:10.2147/IDR.S375043
75. Siddiqui TR, Bibi S, Mustufa MA, et al. High prevalence of typhoidal Salmonella enterica serovars excreting food handlers in Karachi-Pakistan: a probable factor for regional typhoid endemicity. *J Health Popul Nutr*. 2015;33(1):27. doi:10.1186/s41043-015-0037-6
76. Centers for Diseases Control and Prevention. Food safety: Salmonella and food. Available from: <https://www.cdc.gov/foodsafety/communication/salmonella-food.html>. Accessed June 5, 2023.
77. Post AS, Diallo SN, Guiraud I, et al. Supporting evidence for a human reservoir of invasive non Typhoidal Salmonella from household samples in Burkina Faso. *PLoS Negl Trop Dis*. 2019;13(10):e0007782. doi:10.1371/journal.pntd.0007782
78. Kagambèga A, Lienemann T, Aulu L, et al. Prevalence and characterization of Salmonella enterica from the faeces of cattle, poultry, swine and hedgehogs in Burkina Faso and their comparison to human Salmonella isolates. *BMC Microbiol*. 2013;13(1):253. doi:10.1186/1471-2180-13-253
79. Busani L, Cigliano A, Taioli E, et al. Prevalence of Salmonella enterica and Listeria monocytogenes contamination in foods of animal origin in Italy. *J Food Prot*. 2005;68(8):1729–1733. doi:10.4315/0362-028X-68.8.1729
80. Langata LM, Maingi JM, Musonye HA, et al. Antimicrobial resistance genes in Salmonella and Escherichia coli isolates from chicken droppings in Nairobi, Kenya. *BMC Res Notes*. 2019;12(1):1–6. doi:10.1186/s13104-019-4068-8

81. Magwedere K, Rauff D, De Klerk G, et al. Incidence of nontyphoidal Salmonella in food-producing animals, animal feed, and the Associated Environment in South Africa, 2012–2014. *Clin Infect Dis*. 2015;61(suppl 4):283–289. doi:10.1093/cid/civ663
82. Odoch T, Wasteson Y, Abée-Lund T, et al. Prevalence, antimicrobial susceptibility and risk factors associated with non-typhoidal Salmonella on Ugandan layer hen farms. *BMC Vet Res*. 2017;13(1):365. doi:10.1186/s12917-017-1291-1
83. Quynh HL, Fries R, Padungtod PP, et al. Prevalence of Salmonella in Retail Chicken Meat in Hanoi, Vietnam. *Ann N Y Acad Sci*. 2006;1081(1):257–261. doi:10.1196/annals.1373.032
84. Baowei Y, Meili X, Xin W, et al. Prevalence of Salmonella on raw poultry at retail markets in China. *J Food Prot*. 2011;74(10):1724–1728. doi:10.4315/0362-028X.JFP-11-215
85. Lestari SI, Feifei H, Wang FEI, Beilei G. Prevalence and antimicrobial resistance of Salmonella serovars in conventional and organic chickens from Louisiana retail stores. *J Food Prot*. 2009;72(6):1165–1172. doi:10.4315/0362-028X-72.6.1165
86. Dione M, Ikumapayi UN, Saha D, et al. Clonal Differences between Non-Typhoidal Salmonella (NTS) Recovered from Children and Animals Living in Close Contact in The Gambia. *PLoS Negl Trop Dis*. 2011;5(5):e1148. doi:10.1371/journal.pntd.0001148
87. Sharon M, Tennant SD, Levy H, et al. Identification by PCR of Non-typhoidal Salmonella enterica Serovars Associated with Invasive Infections among Febrile Patients in Mali. *PLoS Negl Trop Dis*. 2010;4(3):e621.
88. Saravanan S, Purushothaman V, Murthy TR, et al. Molecular epidemiology of Nontyphoidal Salmonella in poultry and poultry products in India: implications for human health. *Indian J Microbiol*. 2015;55(3):319–326. doi:10.1007/s12088-015-0530-z
89. Quiroz-Santiago C, Rodas-Suárez OR, Vázquez QCR, et al. Prevalence of Salmonella in vegetables from Mexico. *J Food Prot*. 2009;72(6):1279–1282. doi:10.4315/0362-028X-72.6.1279
90. Liang Z, Ke B, Deng X, et al. Serotypes, seasonal trends, and antibiotic resistance of non-typhoidal Salmonella from human patients in Guangdong Province, China, 2009–2012. *BMC Infect Dis*. 2015;15(1):53. doi:10.1186/s12879-015-0784-4
91. Oneko M, Kariuki S, Muturi-Kioi V, et al. Emergence of community-acquired, multidrug-resistant invasive nontyphoidal Salmonella Disease in Rural Western Kenya, 2009–2013. *Clin Infect Dis*. 2015;61(suppl 4):310–316. doi:10.1093/cid/civ674
92. Gao F, Huang Z, Xiong Z, et al. Prevalence, serotype, and antimicrobial resistance profiles of children infected with Salmonella in Guangzhou, southern China, 2016–2021. *Front Pediatric*. 2023;11:1077158. doi:10.3389/fped.2023.1077158
93. Delgado-Sua' rez EJ, Palo's-Guite'rrez T, Rui'za Lo'pez FA, et al. Genomic surveillance of antimicrobial resistance shows cattle and poultry are a moderate source of multi-drug resistant nontyphoidal Salmonella in Mexico. *PLoS One*. 2020;16(5):e0243681. doi:10.1371/journal.pone.0243681

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>