ORIGINAL RESEARCH **RETRACTED ARTICLE: LncRNA HOTAIRMI** Inhibits the Proliferation and Invasion of Lung Adenocarcinoma Cells via the miR-498/WWOX Axis

This article was published in the following Dove Press journal: Cancer Management and Research

Tian-jun Chen^{1,*} Fei Gao^{1,2,*} Tian Yang¹ Hong Li¹ Yang Li¹ Hui Ren¹ Ming-wei Chen¹

¹Department of Respiratory and Critical Care Medicine, The First Affiliated Hospital of Xi'an Jiaotong University, Xi'an, Shaanxi 710061, People's Republic of China; ²Ultrasound Department, Huashan Central Hospital of Xi'an, Xi'an, Shaanxi 710043, People's Republic of China

*These authors contributed equally to this work



Correspondence: Ming-wei Chen Department of Respiratory and Critical Care Medicine, The First Affiliated Hospital of Xi'an Jiaotong University, No. 277, Yanta West Road, Xi'an, Shaanxi, People's Republic of China Tel +86 29-85323888 Email wosyishen66@163.com



Background: Lung adenocarcinoma (ADC) is a *p* for form of lun, r, which is a main ar cause of global cancer-related death in male and smale presents. LncRAAs are implicated in sms of the LncRNA HOTAIRM1 in tumor development. However, the functions and me ADC are not known.

Materials and Methods: Here, the down ulated HOLARM1 in ADC was selected by TCGA analysis. Subsequently, <u>aRT-PCR</u>, CCA EdU, cell apoptosis, cell cycle and cell invasion assays were utilized for evaluating the ress of HOTAIRM1 in ADC. Finally, we explored the mechanism of HOTAIRM1 ADC.

Results: HOTAIRM1 expression was conditionably decreased in ADC tissues. The knockdown of HOTAIRM1 promoth the cell cycle, growth, and invasion of ADC. Moreover, HOTAIRM1 content bound min-498 to regulate the expression of WWOX.

IRM . ressed the proliferation and invasion of ADC cells via the Conclusion: HO modulation of mike WWOX axis. This finding suggested that it might be clinically piomarker for ADC. Furthermore, the findings suggest LncRNA HOTAIRM1 valv Je as a candid therapet is target in ADC.

s: HOTALKM1, lung adenocarcinoma, miR-498, WWOX, cell proliferation Key

troduction

Lung cancer, one of the most commonly diagnosed malignant tumors, is a major cause of cancer-related deaths around the world.^{1,2} Lung cancer can be categorized into five histological types: adenocarcinoma, squamous cell carcinoma, large cell carcinoma, small cell carcinoma, and bronchoalveolar carcinoma.^{3,4} Among them, lung adenocarcinoma (ADC) is the most prevalent and has high metastatic and invasion likelihood.⁵ The 5-year overall survival rate of patients with lung cancer is about 18.1%; this rate is dependent on tumor histology, cancer stage, general health status of the host and other factors.⁶ Currently, no biomarker is available for the early diagnosis and prognosis of ADC. Therefore, it is important to find new targets and biomarkers for its early diagnosis.

Long noncoding RNAs (LncRNAs) are a new class of host factors that has been attracting many attentions in recent years. As the most common subtype of noncoding RNAs, LncRNAs have over 200 nucleotides. In addition, they also participate in many physiopathologic processes, including epigenetic gene expression regulation, genomic imprinting, chromatin organization, immunoregulation, cell development and differentiation, oncogenesis and viral pathogenesis.⁷⁻¹⁰ For

CO 000 COLOR then et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress.com/terms.php you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (https://www.dovepress.com/terms.php).

instance, LncRNA ODIR1 suppresses osteogenic differentiation of hUC-MSCs via the FBXO25/H2BK120ub/ H3K4me3/OSX axis.¹¹ LncRNAs participate in the development of colorectal cancer and are a new target biomolecule.¹² LncRNA LCAT1 sponges miR-4715-5p in lung cancer as a ceRNA to regulate RAC1 functions.¹³

HOXA transcript antisense RNA, myeloid-specific 1 (HOTAIRM1), an LncRNA between the HOXA1 and HOXA2 loci, was first determined as a myeloid-specific regulator of HOXA gene family, which modulates the transcription of target genes by chromosome remodeling during myeloid cell maturation and differentiation.^{14,15} HOTAIRM1 serves as a tumor inhibitor in head and neck, colorectal, and hepatocellular carcinoma;^{16–18} however, its roles in ADC are unclear. WW domain-containing oxidoreductase which is known as WWOX, FOR and WOX1 initially considered as a tumor suppressor protein.¹⁹

We identified that HOTAIRM1 expression is decreased in ADC tissues and that this downregulation enhances the growth and invasion of ADC cells. Moreover, HOTAIRM1 may exert its actions via sponging of miR-498. Overall, our results suggest that HOTAIRM1 might have a crucial role in ADC pathogenesis.

Materials and Methods Sample Collection from ADC Patient and Culture of ADC Cell Line.

Forty-six pairs of tissue samples were collected from ADC patients of the First Affiliated H sphere of Xi'an scrotong University. Ethical approvale for this state x was obtained from the First Affiliater Hospital of Xian Jiaotong University's Ethics Committee and written informed consents were obtained from the participants. All populationrelated studies were performed in the World Medical Association perclaration of Helsenki.

The ADC all Vies From and PC-9 were procured from American Type alture Collection (ATCC, Rockville, MD, USA) and cultured the humid 5% CO₂ atmosphere at 37°C in RPMI-1640 medium containing 10% fetal bovine serum (GibcoTM, Thermo Fisher Scientific, CA).

Real-Time PCR Assay, Cell Transfection, Lentivirus Production and Transduction

RNAs were detached from cells with the Trizol reagent (Invitrogen, Life Technologies). GoScript[™] Reverse Transcription System (Promega, Madison, WI, USA) was

utilized for converting total RNAs to cDNAs, while GoTaq[®] qRT-PCR Master mixes (Promega, Madison, WI, USA) and C1000 Touch[™] Real-Time Thermal Cycler (Bio-Rad Laboratories, Hercules, CA, USA) were used for all the qRT-PCR reactions. The following primers were used in this study: GAPDH Forward: 5'-AATGGG CAGCCGTTAGGAAA-3'; Reverse: 5'-TGAAGGGGTC AT TGATGGCA-3', HOTAIRM1 Forward: 5'- GCGGTT CTCTTGGCTCAAAA -3'; Reverse: 5'- CCCCGTCCG AAAGCATTAAG -3', miR-498 Forward: 5'- ACACTCC AGCTGGGCUUUUUGCGGGGGGGGCCC-3'; Reverse: 5'- CTCAACTGGTGTCGTGGAC CGGCA TTCAGTT GAGAAAGTTCG -3', WWOX Forward: 5'- CTCCCA GGGATGTTTTGTGC -3' Revers 5'- CAGCCAAC CAGGTTTTTACT -3' I experiment v carried out in triplicate. $2^{-\Delta\Delta t}$ method was applied for measuring the expression of target genes.

Ruibo (Guo, thou, China, we used for synthesizing miRNA-NC, miR-4, anti-miR-NC and anti-miR-498 for transference. The plashid WWOX was commercially acquired from Santa Cruz. Subsequently, oligonucleotide transfection was conducted with Lipofectamine[®] 2000 reage. (Invitrogun, Thermo Fisher Scientific) following its protoc

TAIRM1-specific oligonucleotides encoding short airpin RNAs (shRNAs) were subcloned into GV102 Genechem, Shanghai, China). Neomycin or puromycin mvitrogen) was added for the screening of stable cell lines. Sh-NC was applied as the control.

Cell Counting Kit-8 (CCK8) Assay and EdU Assay

After seeding preconditioned cells into a 96-well plate, cells were incubated for 2 h with CCK-8 reagent (DingGuo Bio) at 37°C. We measured the absorbance with a microplate reader (BioTek) at 450 nm.

During EdU assay, Cell-Light EdU DNA Cell Proliferation Kit (RiboBio, Guangzhou, China) was used. Cells were incubated using EdU for 2 h, fixed in 4% paraformaldehyde, dyed using Apollo Dye Solution and mounted using Hoechst 33342. The mounted samples were then photographed and EdU positive cells were counted.

Transwell Assay, Cell Cycle and Apoptosis Assay

Transwell[®] chambers were coated with Matrigel[™] (BD biosciences) for 30 min at 37 °C. Cells treated with

mitomycin were seeded into the upper chambers and 500 μ L complete medium was added to the lower chambers. After 48 h, the cells on the lower chambers were washed with PBS, fixed with 4% paraformaldehyde, and stained with crystal violet solution. Images were captured under a microscope. Each group was analyzed in triplicate.

After isolation by trypsinization, cells were rinsed twice using precooled PBS and then fixed in 70% ethanol at -20° C overnight. Subsequently, fixed cells suspended in 50 µg/mL of propidium iodide (KeyGen BioTECH) and 100 µg/mL of RNaseA (KeyGen BioTECH), cells were incubated for 40 min at room temperature and in the absence of light. Flow cytometry was used for determining the cycle of filtered cells.

In the apoptosis assay, cells were washed using PBS and stained with Annexin V-FITC Apoptosis Detection Kit (Affymetrix eBioscience) according to manufacturer instructions. Flow cytometer (BD Biosciences, BD FACSTM Flow Cytometer) was used for cell analysis.

Western Blot Detection

RIPA lysis buffer (Thermo Fisher Scientific) was used for protein extraction, while BCA Protein assay kit (Beyotime, Nantong, China) was used for the assay of protein concentrations. After protein separation through electrophoresis, proteins were transferred onto PVDF membranes and blocked using 5% defatted milk. Privary antibodies were incubated on polyvinylidene flooride (PODF) membranes overnight at 4°C, and hor radish pero dase (HRP)conjugated secondary untibodite (Beyotine, Nantong, China) were applied or 1 h of room experature to allow the blocking of provins. Prospectrum 600 Imaging System



Figure I HOTAIRMI was decreased in ADC and related to unfavorable overall survival. (A and B) Expression profile of HOTAIRMI in TCGA ADC samples. (C) HOTAIRMI expression was remarkably reduced in ADC tissues in comparison with no-tumor lung tissues. (D) High expression of HOTAIRMI was related to better overall survival, consistent with the survival results of (E) TCGA data analysis. Data were shown as the mean ± SEM of three experiments. ****P<0.001.

(UVP, CA, USA) was used for imaging. The primary antibody WWOX was obtained from Abcam (1:1000; Lot No. ab238144) and GAPDH (1:500; Lot No. ab9484).

RNA-Binding Protein Immunoprecipitation (RIP) Assay

EZ-Magna RIP Kit from Millipore (Billerica, MA, USA) was applied for RIP assay. Cell lysis was conducted using RIP lysis buffer containing RNase and proteinase inhibitor (Millipore). Subsequently, RIP lysates were treated using RIP buffer containing magnetic beads post-conjugation with nonspecific mouse IgG antibody or human anti-Ago2 antibody (Millipore). After digesting proteinase K into the immunoprecipitate, qRT-PCR and gel-staining analyses were conducted for detecting HOTAIRM1 enrichment. The concentrations of RNAs were detected using NanoDrop spectrophotometer, with purified RNAs used for qRT-PCR analysis.

Luciferase Reporter Assay

3'-UTR of WWOX and HOTAIRM1 was amplified and cloned into the downstream the firefly luciferase gene in pGL3 vector (Promega) respectively. This vector was called wild-type (WT) 3'-UTR accordingly. Quick change site-directed mutagenesis kit (Stratagene, Cedar Creek, USA) was used for the mutation of miR-10° coding sites in WWOX and HOTAIRM1 3'-UTR clous, this vector was called mutant (MUT) 3'-UTP. ALS clous were transfected with mut-3'-UTR or wto -UTR also miR-NC or miR-498. Dual Luciferase reporter assay lessem (Promega) was applied for 48 h post consfection. All groups were separately doermined in triplices.

Immunofluorescen Staini g

After fixation and scaling, house were incubated in the antibody of WWO'. (Invitrogen, Carlsbad, California, USA) at 4° C and h. After washing, cells were incubated in FITC-labeled a G (H+L) (Beyotime, Nantong, China) for 60 min. DAPI was used for nuclei staining. The protocol followed was described in detail in our previous study.⁵

Mouse Model

A total of 6 female BALB/c-nu/nu mice (age, 5 weeks; weight, 16–22 g) were purchased from the Model Animal Research Center of Nanjing University. The mice were housed in a sterile room under a 12-h light/dark cycle at

~23°C and 50% humidity, with ad libitum access to food and water. 2×10⁶ H1650 (sh-NC or sh-HOTAIRM1) were loaded into the flanks of BALB/c-nu/nu mice by subcutaneous inoculation. The mice were euthanized after harboring tumors for four weeks. Animals were maintained and tested according to the US National Institute of Health Guidelines for Use of Experimental Animals. Ethical approval was obtained from the Ethnic Committee for Experimental Animals in Xi'an Jiaotong University. Animal experiments took place in SPF Animal Laboratory at Xi'an Jiaotong University

Statistical Analysis

Fisher's exact test was ar ned for ide offyin any difference between categorie t variables. For normally distributed data with equal variables the difference was evaluated by two-tailed Statent's *t*-test two group comparisons) or one-way ANOVA colowed by the Bonferroni post hoc test (multigroup, comparisons). For non-normally distributed data of data with unequal variances, the difference was evaluated by a nonparametric Mann–Whitney *U*-test (two group comparisons)) or the Kruskal–Wallis test followed by the Bondene post hoc test (multigroup comparisons).

able I The Correlation of HOTAIRMI Expression with	Clinical
arameters in Patients with Lung Adenocarcinoma	

Clinicopathological Features	Number of Cases	HOTAIRMI Expression		p value
		Low (n=23)	High (n=23)	
Gender				0.2362
Male	25	10	15	
Female	21	13	8	
Age				0.7683
<60	24	13	11	
≥60	22	10	12	
Tumor size				0.0377*
≤5	24	8	16	
>5	22	15	7	
TNM stages				0.0058*
1/11	28	9	19	
III/IV	18	14	4	
Lymph node				0.0072*
metastasis				
Absent	24	7	17	
Present	22	16	6	

Note: *P< 0.05.

Data are presented as the mean \pm SD. P<0.05 was considered to indicate a statistically significant difference.

Results HOTAIRMI Expression Was Decreased in ADC Tissues

The analysis was conducted via the TCGA database and the downregulated HOTAIRM1 in ADC was selected as the topic of interest (Figure 1A and B). qRT-PCR assay was conducted for determining HOTAIRM1 expression in ADC tissues and corresponding normal tissues, which uncovered that HOTAIRM1 expression was remarkably reduced in ADC tissues compared with those in normal lung tissues (Figure 1C).

Low Expression of HOTAIRMI Was Associated with Unfavorable Prognosis of ADC Patients

The association between HOTAIRM1 expression and patients' clinicopathological features are shown in Table 1. Forty-six ADC patients were stratified equally into low expression group and high expression group with median HOTAIRM1 expression as the patoff. No opparent association existed between HOTAP W1 expression and gender



Figure 2 HOTAIRM1 down-regulation promoted the proliferation and cell cycle of ADC cells. (A) HOTAIRM1 expression in H1650 and PC-9 cells post transduction with shRNA vector (sh-NC) or HOTAIRM1 shRNA vector (sh-HOTAIRM1) control. (B and C) HOTAIRM1 down-regulation notably increased cell proliferation in CCK-8 and EdU assays. (D) sh-HOTAIRM1 cells presented with significantly high percentage in S phase. Data were shown as the mean ± SEM of three experiments. *P<0.05, **P<0.01, ***P<0.001.



Figure 3 HOTAIRM1 down-regulation promoted ADC cell invasion and inhibited cell apoptor (A) Down-regulated HOTAIRM1 decreased the percentage of apoptosis in ADC cells. (B) Down-regulated HOTAIRM1 increased the ADC cell invasion. Data were show as the mean ± EM of three experiments. *P<0.05, **P<0.01.

(P=0.2362) or age (P = 0.7683). However, an overt association was observed between the knockdown of Varue RM1 expression and clinical stage (P=0.0058), and lymph node metastasis (P=0.0072) and tumor size (P=0.007). Additionally, the knockdown of HOT aRM1 we linked to worse overall survival of ADC cannots, as induced by Kaplan-Meier analysis (Figure 1D). This essult was consistent with the TCGA database analysis (Figure 1E).

HOTAIRMI Down Regulation Promoted ADC in vitro

and PC-9 cells, which proved to be We established H16 stable for transition with sh-HOTAIRM1 (Figure 2A). HOTAIRM1 dow regulation notably enhanced the cell proliferation of cells as shown observed in the CCK-8 assay (Figure 2B). During the EdU assay, cells incorporating EdU were considerably higher in the sh-HOTAIRM1 group than in the sh-NC group (Figure 2C). Additionally, based on the flow cytometry analysis, less cells in G1 phase and more cells in S phase were observed in sh-HOTAIRM1 cells (Figure 2D). Cell apoptosis was analyzed to identify whether LncRNA HOTAIRM1 had any effect on the apoptosis, which suggested that downed tated HOTAIRM1 reduced cell apoptosis in ADC cells (Figure 3A). Following that, Transwell avasion Assay showed that inhibition of HOTAIRM1 increased ADC cell invasion (Figure 3B).

HOTAIRM1 Functioned as a ceRNA in ADC by Sponging miR-498

miRNAs capable of complementary base paring with HOTAIRM1 were searched by using online software program starbase v2.0 (http://starbase.sysu.edu.cn/ mirLncRNA.php). miR-498 was selected from miRNAs that promoted the cell proliferation and invasion in many cancers.^{20,21} Figure 4A depicted the binding site of miR-498 on HOTAIRM1. A mutated miR-498 binding site of HOTAIRM1 (HOTAIRM1-MUT) and predicted miR-498 binding site of HOTAIRM1 (HOTAIRM1-WT) were cloned into reporter plasmids. Luciferase activity was notably reduced by co-transfecting miR-498 and HOTAIRM1-WT. On the other hand, luciferase activity remained unchanged after co-transfection with miR-498 and HOTAIRM1-MUT (Figure 4A). miRNA is distributed in the cytoplasm, which is a component of the RNA-induced silencing complex (RISC) containing Ago2. Ago2 is



Figure 4 Reciprocal replacion between LncRNA HOTAIRMI and miR-498. (A) Binding sites of miR-498 on HOTAIRMI and dual-luciferase reporter assay for the validation of the interaction between HOTAIRMI and miR-498. (B) RIP assay was performed to reveal the combination between HOTAIRMI and miR-498. (C) Up-regulated miR-498 was seen to C tissue up qRT-PD assay. (D) miR-498 expression was reduced by anti-miR-496 and increased by miR-496 mimics. (E) miR-498 negatively regulated HOT aRMI expression. The 498 expression was increased by sh-HOTAIRMI. Data were shown as the mean ± SEM of three experiments. **P<0.01, ***P<0.001

required for h. 2NA-mediated gene silencing. In this study, we analyzed whether HOTAIRM1 and miR-498 contained the same RISC and affected RIP assay similarly in H1650 and PC-9 cells. It was observed that HOTAIRM1 and miR-498 were enhanced in Ago2 immunoprecipitate compared to IgG immunoprecipitate controls (Figure 4B). Moreover, miR-498 expression was higher in ADC tissues in comparison with adjacent normal lung tissues (Figure 4C). To further verify the correlation between HOTAIRM1 and miR-498, miR-498 expression in ADC cells was changed

by transfecting with miR-498 overexpressed plasmids and anti-miR-498 (Figure 4D). Whether the levels of miR-498 was affected, HOTAIRM1 expression levels was determined. As displayed in Figure 4E, lower HOTAIRM1 expression was observed in miR-498 treated cells, as opposed to HOTAIRM1 expression in anti-miR-498 treated cells. Then, we determined whether miR-498 expression was regulated by HOTAIRM1. Results showed that sh-HOTAIRM1 could significantly increase the level of miR-498 in H1650 and PC-9 cells (Figure 4F). In summary, our



Figure 5 WWOX was the direct target of miR-498. (A) Brang sequence between $(R-498 \text{ and } WWOX \text{ and } dual-luciferase reporter assay for the validation of the interaction between miR-498 and WWOX. (B) WWOX provin expression (DC tissues. (C and D) miR-498 repressed WWOX expression in ADC cells. (E and F) Decreased HOTAIRMI inhibited the expression of WV X, we are miR-498 treasment contributed to partial increase in WWOX expression level. Data were shown as the mean <math>\pm$ SEM of three experiments. *P<0.05, ****0.01.

findings include that miR-498 diverse bound to HOTAIRM1 had negative regulatory effects on HOTAIRM1 expression

WWOX V as the Direct arget of miR-498

MicroRNA exerted peir biological functions mainly by means of remating their downstream targets.^{22,23} WWOX was sended from the predicted target genes of miR-498 for further studies. Dual-luciferase reporter assay revealed that miR-496 mimics efficiently decreased the luciferase activity of WWOX-WT instead of that of WWOX-MUT (Figure 5A). Furthermore, WWOX expression was also observed in ADC tissues. Western blot assay uncovered that WWOX expression was reduced in ADC tissues compared with those in adjacent normal lung tissues (Figure 5B). Furthermore, miR-498 repressed WWOX expression in H1650 and PC-9 cells at both RNA and protein levels (Figure 5C and D). On the other hand, WWOX level was decreased by sh-HOTAIRM1 and recovered by anti-miR-498 treatment in sh-HOTAIRM1 cells (Figure 5E and F). In all, our data uncovered that the WWOX was a downstream target of miR-498 and under the regulation of HOTAIRM1.

HOTAIRMI Down-Regulation Increased Tumor Growth in vivo

In vivo experiment was utilized for verifying the roles of HOTAIRM1 in tumorigenesis. In Figure 6A, the down-regulation of HOTAIRM1 promoted tumor growth. Xenograft tumors from sh-HOTAIRM1 group had higher volume and weight of xenograft tumors than those from the sh-NC group (Figure 6B and C). HOTAIRM1 expression was then determined in excised tumor tissues, which



Figure 6 HOTAIRM1 inhibition increased tumor growth in vivo. (A) Xenograft tumors (=3 in each group). (B and C) Measurement and calculation of tumor volume and tumor weight. (D) HOTAIRM1 expression was determined in xenograft tumors. (E) K takdown of HOTA M1 notably restrained WWOX in tumors in comparison with the negative control group. *P<0.05, **P<0.01.

revealed the downregulation of HOTAIRM1 expression in the sh-HOTAIRM1 group relative to that of the shell' group (Figure 6D). Tumor section staining was perform 1 for WWOX expression, which was decreased in she HOTAIRM1 group (Figure 6E). It summary, these mudings supported the growth inhibition effect to HOTAIRM1 in vivo.

HOTAIRMI Bugulated ADC Cell Proliferation, Avanon, Apoptosis and Cycle Through in R-465/WWOX Axis In order to verse the roles of HOTAIRM1/miR-498/ WWOX wisein tung adenocarcinoma, we performed a "rescue exteriment" in H1650 and PC-9 cells. H1650 (sh-HOTAIRM1 and PC-9 (sh-HOTAIRM1) cells which were stable in transfection with sh-HOTAIRM1 were stratified into three groups for transfection with NC, anti-miR -498, and WWOX overexpressed plasmids. CCK-8 and EdU experiments indicated that the proliferation of antimiR-498 group and WWOX group was notably decreased (Figure 7A and B). Additionally, in both anti-miR-498 and WWOX groups, the invasion of H1650 (sh-HOTAIRM1) and PC-9 (sh-HOTAIRM1) cells was inhibited (Figure 7C) and the G1 arrest and apoptosis of ADC cells were promover (Figure 7D and E).

Discussion

ADC, as the most prevalent non-small cell lung cancer, causes over 500,000 deaths per year around the world.²⁴ At present, there is an urgent demand for effective new therapies for ADC; however, efforts should first be made to deepen the understanding of underlying mechanisms of the development and progression of ADC. Increase evidence shows that LncRNAs may modulate multiple tumorigenesis processes, especially ADC.^{25–27} In this study, through TCGA data analysis of ADC related LncRNAs, downregulated HOTAIRM1 was selected as the topic of interest and was subsequently verified in human tissues.

During functional analysis, HOTAIRM1 inhibition promoted the proliferation and invasion of H1650 and PC-9 cells. Furthermore, downregulated HOTAIRM1 contributed to the development from the G1 phase into the S phase of H1650 and PC-9 cells and prevented them from apoptosis. In addition, the reduction of HOTAIRM1 expression resulted in worse overall survival of ADC patients. These data identified HOTAIRM1 as a suppressor gene in ADC. Moreover, HOTAIRM1 may be a candidate prognostic marker for ADC.



Figure 7 HOTAIRM1 regulated bC cell procention and invasion via miR-498/WWOX axis. (A) CCK-8 assay and (B) EdU assay revealed that anti-miR-498 and WWOX overexpression promoted cell procentic on H1650 and PC-9 cells after stable transfection with sh-HOTAIRM1. (C) Cell invasion assay in H1650 (sh-HOTAIRM1) and PC-9 (sh-HOTAIRM1) cells. (P) Cell cycles as in H1650 (sh-HOTAIRM1) and PC-9 (sh-HOTAIRM1) cells. (P) Cell cycles as in H1650 (sh-HOTAIRM1) and PC-9 (sh-HOTAIRM1) cells. (E) Cell apoptosis assay in H1650 (sh-HOTAIRM1) and PC-9 (sh-HOTAIRM1) cells. D2 were nown as mean SEM of three experiments. *P<0.05, **P<0.01.

Human W VO gene is distributed on chromosomal site FRA16D or 16q. FRA16D, which is a focus of genomic instability and is preciposed to breakage and germline and somatic copy number variation. Thus, WWOX is a commonly deleted target in various cancers,^{28,29} such as breast cancer,³⁰ non-small cell lung cancer,³¹ bladder cancer,³² prostate cancer.³³ In the present study, WWOX was reduced in ADC tissues. In addition, stable downregulation of HOTAIRM1 expression significantly promoted the proliferative and invasive capabilities of ADC cells, but overexpression WWOX reversed these influences.

Conclusion

In conclusion, HOTAIRM1 is downregulated in ADC and this downregulation enhances the proliferation and invasion of ADC cells via the miR-498/WWOX axis (Supplementary Figure 1).

Abbreviations

ADC, lung adenocarcinoma; LncRNAs, long noncoding RNAs; HOTAIRM1, HOXA transcript antisense RNA, myeloid-specific 1; CCK8, Cell counting kit-8; RIP, RNA-binding protein immunoprecipitation.

Data Sharing Statement

The datasets used and/or analyzed during the present study are available from the corresponding author upon reasonable request.

Funding

This work was supported by the Key Research and Development Project of Shaanxi Province [2019KW-037].

Disclosure

The authors report no conflicts of interest in this work.

References

- Yuan S, Liu Q, Hu Z, et al. Long non-coding RNA MUC5B-AS1 promotes metastasis through mutually regulating MUC5B expression in lung adenocarcinoma. *Cell Death Dis.* 2018;9(5):450. doi:10.1038/ s41419-018-0472-6
- Shen QM, Wang HY, Xu S. MARCH9 suppresses lung adenocarcinoma progression by downregulating ICAM-1. *Cell Physiol Biochem.* 2018;50(1):92–107. doi:10.1159/000493961
- 3. Yang Y, Liu L, Zhang Y, et al. MiR-503 targets PI3K p85 and IKK-beta and suppresses progression of non-small cell lung cancer. *Int J Cancer*. 2014;135(7):1531–1542. doi:10.1002/ijc.28799
- Duruisseaux M, Esteller M. Lung cancer epigenetics: from knowledge to applications. *Semin Cancer Biol.* 2018;51:116–128. doi:10.1016/j.semcancer.2017.09.005
- 5. Hao X, Han F, Ma B, et al. SOX30 is a key regulator of desmontance gene suppressing tumor growth and metastasis in ung adenocarcinoma. J Exp Clin Cancer Res. 2018;37(1) U doi:10.1186/s13046-018-0778-3
- Alam H, Li N, Dhar SS, et al. HP1gamma prodotes lupp adenocal cinoma by downregulating the transcriptor-repression explator. NCOR2 and ZBTB7A. *Cancer Procession*, 23, 377–43, 3834–3548. doi:10.1158/0008-5472.CAN-17-357
- 7. Qu D, Sun WW, Li L, et al. Longer Lodding RNA W LAT1 releases epigenetic silencing of HIV-1 epilicatic by displacing the polycomb repressive complex 2 from binding to the LTR promoter. *Nucleic Acids Res.* 2019;47(6);20:5–3027. doi:10.10.104/mar/gkz117
- 8. Yong H, Wu G, Chete, et al. In RNA MALAM accelerates skeletal muscle cell apopto is and collammatory response in sepsis by decreasing BRCA1 portion by opruiting EZH2. *Mol Ther Nucleic Acid* (2019)19:19. 108. doi: 0.1016/j.omtn.2019.10.028
- Liao K, Jun Y, G W, et a Bloring IncRNA MALAT1/miR-199a/ ZHX1 axis inhibit glioblasto, a proliferation and progression. *Mol Ther. Netc.* 4 (2019) 288–399. doi:10.1016/j.omtn.2019.09.005
- Xu H, J. Lev, Xu X, et al. Inducible degradation of lncRNA Sros1 promotes V-gamma-mediated activation of innate immune responses by ubilizing Stat1 mRNA. *Nat Immunol.* 2019;20 (12):1621–1630. doi:10.1038/s41590-019-0542-7
- He S, Yang S, Zhang Y, et al. LncRNA ODIR1 inhibits osteogenic differentiation of hUC-MSCs through the FBXO25/H2BK120ub/ H3K4me3/OSX axis. *Cell Death Dis.* 2019;10(12):947. doi:10. 1038/s41419-019-2148-2
- Sun W, Ren S, Li R, Zhang Q, Song H. LncRNA, a novel target biomolecule, is involved in the progression of colorectal cancer. *Am J Cancer Res.* 2019;9(11):2515–2530.
- Yang J, Qiu Q, Qian X, et al. Long noncoding RNA LCAT1 functions as a ceRNA to regulate RAC1 function by sponging miR-4715-5p in lung cancer. *Mol Cancer*. 2019;18(1):171. doi:10.1186/s12943-019-1107-y

al Long non-coding RNA_HOTAIRM1

Chen et al

- Liang Q, Li X, Guan G, et al. Long non-coding RNA, HOTAIRM1, promotes glioma malignancy by forming a ceRNA network. *Aging* (*Albany NY*). 2019;11(17):6805–6838. doi:10.18632/aging.102205
- Zhang X, Lian Z, Padden C, et al. A myelopoiesis-associated regulatory intergenic noncoding RNA transcript within the human HOXA cluster. *Blood*. 2009;113(11):2526–2534. doi:10.1182/blood-2008-06-162164
- Zheng M, Liu X, Zhou Q, Liu G. HOTAIRM1 competed endogenously with miR-148a to regulate DLGAP1 in head and neck tumor cells. *Cancer Med.* 2018;7:3143–3156.
- Wan L, Kong J, Tang J, et al. HOTAIRM1 as a potential biomarker for diagnosis of colorectal cancer functions the role in the tumour suppressor. J Cell Mol Med. 2016;20(11):2036–2044. doi:10.1111/ jcmm.12892
- Zhang Y, Mi L, Xuan Y, et al. LncPiper 10TAIRM1 inhibits the progression of hepatocellular carcine and by innerting the Wnt signaling pathway. *Eur Rev Med Physicacol Sci.* 2018 2(15):4861–4868. doi:10.26355/eurrev 201808 150.
- McBride KM, Kil H, My , et al. Veox deletio in mouse B cells leads to genomic instal any, neoplastic vestor ration, and monoclonal gammopathies *cont Once* 2019;9:51 vaoi:10.3389/fonc.2019. 00517
- Chai C, Wuy, Wang ucisenstat Lo, Leng RP. MicroRNA-498 promotes the liferation and signation by targeting the tumor suppressor sor PTLA upbreast cance wells. *Carcinogenesis*. 2018;39(9): 1185–1196. doi: 1093/carcin/bgy092
- 21. Pro XM, Liu XI, et al. MicroRNA-498 promotes proliferation, migration, and vasion of prostate cancer cells and decreases radiation sensitivity by targeting PTEN. *Kaohsiung J Med Sci.* 2019;35(11): 59–671. doi:10.1002/kjm2.12108
- Roberts JT, Borchert GM. Computational prediction of microRNA taget genes, target prediction databases, and web resources. *Methods Mol Biol.* 2017;1617:109–122.
- 24. Pu X, Wang J, Li W, et al. COPB2 promotes cell proliferation and tumorigenesis through up-regulating YAP1 expression in lung adenocarcinoma cells. *Biomed Pharmacother*. 2018;103:373–380. doi:10.1016/j.biopha.2018.04.006
- 25. Zhou J, Xiao H, Yang X, et al. Long noncoding RNA CASC9.5 promotes the proliferation and metastasis of lung adenocarcinoma. *Sci Rep.* 2018;8(1):37. doi:10.1038/s41598-017-18280-3
- 26. Zhang M, Gao C, Yang Y, et al. Long noncoding RNA CRNDE/ PRC2 participated in the radiotherapy resistance of human lung adenocarcinoma through targeting p21 expression. *Oncol Res.* 2018;26(8):1245–1255. doi:10.3727/096504017X14944585873668
- 27. Peng Z, Wang J, Shan B, et al. The long noncoding RNA LINC00312 induces lung adenocarcinoma migration and vasculogenic mimicry through directly binding YBX1. *Mol Cancer*. 2018;17(1):167. doi:10. 1186/s12943-018-0920-z
- Saigo C, Kito Y, Takeuchi T. Cancerous protein network that inhibits the tumor suppressor function of WW domain-containing oxidoreductase (WWOX) by aberrantly expressed molecules. *Front Oncol.* 2018;8:350. doi:10.3389/fonc.2018.00350
- 29. Tochigi Y, Takamatsu Y, Nakane J, Nakai R, Katayama K, Suzuki H. Loss of Wwox causes defective development of cerebral cortex with hypomyelination in a rat model of lethal dwarfism with epilepsy. *Int J Mol Sci.* 2019;20(14):3596. doi:10.3390/ijms20143596
- Pospiech K, Pluciennik E, Bednarek AK. WWOX tumor suppressor gene in breast cancer, a historical perspective and future directions. *Front Oncol.* 2018;8:345. doi:10.3389/fonc.2018.00345
- 31. Wang XH, Gan CZ, Xie JY. Inhibition of miR-24 suppresses malignancy of human non-small cell lung cancer cells by targeting WWOX in vitro and in vivo. *Thorac Cancer*. 2018;9(12):1583–1593. doi:10. 1111/1759-7714.12824

- 32. Li G, Sun L, Mu Z, Huang Y, Fu C, Hu B. Ectopic WWOX expression inhibits growth of 5637 bladder cancer cell in vitro and in vivo. *Cell Biochem Biophys.* 2015;73(2):417–425. doi:10.1007/s12013-015-0654-0
- 33. Lin JT, Li HY, Chang NS, Lin CH, Chen YC, Lu PJ. WWOX suppresses prostate cancer cell progression through cyclin D1-mediated cell cycle arrest in the G1 phase. *Cell Cycle*. 2015;14 (3):408–416. doi:10.4161/15384101.2014.977103

Cancer Management and Research

Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/cancer-management-and-research-journal

Dovepress