ORIGINAL RESEARCH

Current Psychotic-Like Experiences Among Chinese College Students: Prevalence, Correlates, and Its Relationship with Suicidal Ideation

Bingna Xu¹, Chunping Chen¹, Dongfang Wang^{2,3}

¹Institute of Education, Xiamen University, Xiamen, People's Republic of China; ²School of Psychology, South China Normal University, Guangzhou, People's Republic of China; ³Centre for Studies of Psychological Applications, Guangdong Key Laboratory of Mental Health and Cognitive Science, Ministry of Education Key Laboratory of Brain Cognition and Educational Science, South China Normal University, Guangzhou, People's Republic of China

Correspondence: Chunping Chen, Institute of Education, Xiamen University, Siming Road, Xiamen, 361005, People's Republic of China, Email 894906014@qq.com; Dongfang Wang, School of Psychology, South China Normal University, Shipai Road, Guangzhou, 510631, People's Republic of China, Email wdfpsyc@126.com; wangdf@m.scnu.edu.cn

Background: Current psychotic-like experiences (PLEs) are prevalent among adolescents. However, epidemiological data on the prevalence rate of current PLEs among college students and its clinical correlates are scarce and limited. To address this study gap, this study aimed to investigate the prevalence and correlates of current PLEs among Chinese college students.

Methods: A cross-sectional study was designed with a sample of 18,578 college students in China. Each participant completed a survey on social-demographic characteristics variables, PLEs, sleep disturbance, depression, anxiety, and suicidal ideation. Logistic regression analyses were conducted to explore the influential factors for PLEs among all participants and suicidal ideation among those with PLEs.

Results: Among Chinese college students, 40.3% of the participants reported having at least one PLE in the past month, while 7.5% had frequent PLEs. Males, age, residence location in town, left-behind experiences, poor family economic status, chronic physical illness, family history of mental disorder, BMI \geq 28 [obesity], sleep disturbance, depression, anxiety, and suicidal ideation were independent correlates for frequent PLEs. Furthermore, the prevalence of suicidal ideation among participants with frequent PLEs was 32.1%. Lower grades, sleep disturbance, depression, and anxiety were independent predictors of suicidal ideation among college students who experience PLEs frequently.

Conclusion: Current PLEs are widespread among Chinese college students. Suicidal ideation is largely prevalent among individuals who suffer frequent PLEs, suggesting that special measures and attention should be provided to these students based on relevant factors to prevent their suicidal ideation and behavior.

Keywords: psychotic-like experiences, suicidal ideation, correlates, college students

Introduction

The prevalence of mental disorders among children and adolescents has significantly increased in recent decades.¹ The investigation of the underlying factors contributing to mental disorders has garnered considerable attention. Psychotic-like experiences (PLEs), for instance, delusions and hallucinations, are universally defined as the resemblance of positive symptoms of psychosis in the absence of a full-blown psychotic disorder.² Numerous literature empirically revealed that PLEs in early age are robustly associated with later psychotic disorders^{3,4} and non-psychotic disorders.^{5,6} In addition to being an early signal of psychopathology, PLEs have been considered as a sign of broader psychological vulnerability and a transdiagnostic clinical marker of clinically significant mental disorders.^{7,8} Given that PLEs are a psychosis risk marker, it is essential to take PLEs into great consideration and adopt a targeted approach to clinical screening, early detection, and the targeting of interventions.

799

© 2024 Xu et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress.com/terms.php you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission form Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (https://www.dovepress.com/terms.php).

PLEs are prevalent among the general population, especially adolescents. However, in previous studies, the prevalence of PLEs varies due to samples, instruments, and screening criteria. For example, Mamah et al explored the 12month prevalence of PLEs among 9564 Kenyan youth (46.7% females, mean age 21.2 years) with the Washington Early Recognition Center Affectivity and Psychosis (WERCAP) Screen and found that 72% reported at least one PLE over the past year.9 A study using the Positive Subscale of Community Assessment of Psychic Experiences (CAPE-P) recruited a community sample of 1489 Tunisian college students (64.3% females, mean age 22.0 years), among which 51.4% had at least one PLE "nearly always" in their lifetime.¹⁰ Utilizing the same measure to assess lifetime PLEs with a sample of 5427 Chinese middle school students (48.8% girls, mean age 12.6 years), another study surprisingly found that 95.7% of participants reported having more than one episode of PLEs, while 17.2% had "nearly always" PLEs.¹¹ It shall be noted that despite the prevalence, nearly 75–90% of PLEs gradually disappear over time, with only a small percentage persisting and eventually developing into clinically significant outcomes.¹² Consequently, the assessment of PLEs over a long time frame (eg, lifetime) is limited in utility in interpreting relationships with current distress or of use in longitudinal or interventional studies.¹³ In fact, most psychiatric diagnoses, as well as attenuated psychotic symptoms (APS), are assessed according to the patient's status in the past month. Therefore, assessing current PLEs within the past month can better and accurately reflect their recent mental health status, making it possible to better track individual mental health in the future. Building on this, our team used the 8-item Positive Subscale of the Community Assessment of Psychic Experiences (CAPE-P8) to examine the current PLEs among 67,538 junior and senior high school students (46.3% girls, mean age 14.56 years) in urban China.¹⁴ Specifically speaking, the study demonstrated that nearly half of the students (49.8%) experienced at least one PLE in the past month, while 15.4% "often" or "nearly always" had PLEs. However, few previous studies have evaluated one-month PLEs among college students with a small sample.^{15,16} Undoubtedly, studies regarding PLEs among college students are insufficient and limited. Therefore, there is an urgent need to use larger epidemiological data to explore current PLEs among college students as a population of late adolescence and provide data-driven insights for clinical intervention in the future.

Prior studies have identified numerous influential factors of PLEs, such as sex,^{17,18} socioeconomic status,^{9,15} residency status,¹¹ sleep disturbance,¹⁹ depression,^{20,21} and anxiety.^{20,22} Among the above factors, sleep disturbance, depression, and anxiety have received special attention because they are more likely to contribute to catastrophic consequences, such as suicide behaviors.^{23–25} Alarmingly, suicide is the second leading cause of death among adolescents and young adults worldwide.²⁶ In China, suicide is the leading cause of death among the population aged 15 to 34,²⁷ accounting for 19% of all deaths.²⁸ Although the overall suicide rate in China has significantly decreased over the past decades, the suicide rate among young people has remained unchanged.^{29,30} Therefore, identifying the risk factors for suicide among adolescents and young adults is of crucial significance.

In recent years, prior studies extensively documented the strong association between PLEs and suicidality.^{31,32} Considering the congenerous risk factors between PLEs and suicidality, particularly sleep disturbance and emotional symptoms, it is crucial to delve deeper into the independent association between PLEs and suicidal behavior, while accounting for these factors through appropriate adjustments. This will enable us to elucidate the mechanism of how PLEs can contribute to suicidal behavior. To our knowledge, few studies have considered the effect of confounding factors on the relationship between PLEs and suicidality. Interestingly, in some studies, the significant relationship between PLEs and suicidal behavior persisted even after controlling for depression.³³ However, this result was not replicated in other studies.³⁴ Accordingly, using a large sample of representative college students, the current study intended to examine the relationship between PLEs and suicidal ideation, which was considered to be a precursor to suicide.³⁵ Additionally, this study explored the independent predictors of suicidal ideation among adolescents with PLEs, while accounting for the effects of confounding factors.

Methods

Subjects

A large-scale online survey was conducted, utilizing a convenience sampling method to recruit college students from four colleges/universities in three provinces (Shanxi, Hunan, and Guangdong) in China. The web-based survey was

distributed through the "Survey Star" online platform and the Quick Response (QR) code of the questionnaire was forwarded to participants via WeChat. All Participants were required to complete the questionnaire by scanning a QR code from their cell phones. The platform automatically recognizes the completeness of the questionnaire filling, and the submission can only be completed if the participant completes all the questions. The recruitment period ran from October 17 to 29, 2023.

A total of 19,236 college students participated in our survey and submitted responses during the recruitment period. To improve the quality of the data, certain exclusion criteria were applied for participation, including: (i) completion time less than 5 minutes, (ii) incorrect identity information (ie, incomplete student number), and (iii) inconsistent survey contents. According to the criteria above, 585 participants were excluded. One question-"Have you ever suffered from any kind of mental health problems?" was used to investigate participants' history of mental disorders. Given that PLEs are a precursor to the onset of mental disorders, 73 participants with a history of psychiatric conditions were excluded from the subsequent analyses. Accordingly, our study obtained results from a final sample of 18,578 participants with valid data.

This study employed an anonymous survey methodology, where participants were identified through a unique student number. Prior to the survey, all participants were required to submit an electronic informed consent form that explicitly stated their right to discontinue or withdraw from the survey at any time without penalty. During the survey, a toll-free psychological hotline was established to offer assistance to participants in need. Furthermore, a comprehensive written report about the overall status of participants was provided to the respective schools. The current study was approved by the Ethics Committee of the School of Psychology, South China Normal University (SCNU-PSY-2023-345).

Measures

Social-Demographic Characteristics

The socio-demographic characteristics of the participants included sex, age, grade, ethnicity, parental marital status, residence location, left-behind experiences (eg, living separately from one or both parents for more than 6 months before 16 years old),³⁶ single child status, parents' education, family economic status, chronic physical illness (e.g, having at least one of arthritis, angina, asthma, diabetes, visual impairment, or hearing problems),³⁷ family history of mental disorder, and body mass index (BMI, calculated as weight (kg)/height (m)²).

PLEs

The CAPE-P8 was employed to assess participants' PLEs over the past month.^{38,39} Each item was rated on a 4-point scale (1 = never; 2 = sometimes; 3 = often; and 4 = nearly always). Participants were regarded to have frequent PLEs when they selected "often" or "nearly always" on one or more items of CAPE-P8.^{40,41} In this study, its Cronbach's alpha was 0.93.

Sleep Disturbance

Sleep disturbance over the past month was assessed through four questions: (1) "Have you had trouble falling asleep?", (2) "Have you woken up frequently during the night?", (3) "Have you woken up very early and can't get back to sleep?", and (4) "How would you rate your overall sleep quality".¹⁹ The first three questions examined insomnia symptoms, and responses to questions range from 1 to 5 (1 = never, 2 = < 1 night/week, 3 = 1-2 nights/week, 4 = 3-5 nights/week, 5 = 5-7 nights/week). The last question evaluated subjective sleep quality, and the options were rated on a 5-point Likert scale, from 1(very good) to 5 (very poor). The higher the score, the poorer the quality of sleep. Participants were classified as having sleep disturbance if they reported any of three insomnia symptoms occurring more than three times per week (rated as a response option of 4 or 5), subjectively reported poor (option 4) or very poor (option 5) sleep quality.^{25,42} In this study, the Cronbach's alpha for 4 items was 0.80.

Depression

The 2-item Patient Health Questionnaire (PHQ-2) was used to measure depressive symptoms over the past two weeks.⁴³ Each item was rated on a 4-point scale from 0 (not at all) to 3 (almost every day), with a total score from 0 to 6. The cutoff total score of 3 and above refers to probable depression.⁴⁴ In this study, its Cronbach's alpha was 0.82

Anxiety

The 2-item Generalized Anxiety Disorder (GAD-2) was utilized to assess anxiety symptoms over the past two weeks.⁴⁵ Each item was rated on a 4-point scale from 0 = not at all, 1 = several days, 2 = more than half the days, to 3 = almost every day, with a total score from 0 to 6. A cut-off total score of 3 and above represented the clinical level of anxiety.⁴⁶ In this study, Cronbach's alpha was 0.87.

Suicidal Ideation

Suicidal ideation was evaluated through three dichotomous questions based on the three-step theory of suicide.^{47,48} The first question was "Are you in pain and hopeless?". Participants are considered to have suicidal ideation if they answered "yes". Participants with suicidal ideation went on to answer the second question - "Does your pain exceed your connectedness?". If participants answered "yes", they had a strong suicidal desire. These participants with strong suicidal ideation were required to answer the third question - "Do you have the capacity to commit suicide?" Participants are considered to have suicidal ideation was 9.5% (N = 1773) among the current sample. Of those with suicidal ideation, 620 college students reported strong suicidal ideation, with 391 of them having suicide attempts. Due to the low rate of suicide attempts (2.1%), this study used overall suicidal ideation (9.5%) for logistic regression analysis.

Statistical Analyses

In this study, continuous variables were reported as mean (standard deviation, SD), and categorical variables were described as frequencies (percentage, %). No evidence of multicollinearity was found among all variables, as indicated by variance inflation factor (VIF) values of 1.800 or below.⁴⁹ Due to the non-normal distribution of age, as confirmed by the Kolmogorov–Smirnov one-sample test conducted on the sample, the Mann–Whitney *U*-test was employed to assess age differences among groups. Moreover, the Chi-square test was used to examine differences among groups for other categorical variables. Multivariate logistic regression analyses were conducted to identify factors that were independently correlated with frequent PLEs and suicidal ideation. Only statistically significant factors (p < 0.05) from the univariate analyses were included in the regression models.⁵⁰ All statistical analyses were performed through SPSS 23.0, and the significance level was set at 0.05.

Results

Description of the Sample

Among 18,578 participants, 5931 (31.9%) were male and 12,647 (68.1%) were female. The mean (SD) age was 20.07 (1.63) years. 488 participants (2.6%) had a chronic physical illness, while 226 (1.2%) reported a family history of mental disorder. Table 1 summarized the detailed social-demographic characteristics of the participants.

Prevalence of PLEs and Suicidal Ideation

Results indicated that two-fifths of participants (40.3%, N = 7485) endured at least one PLE in the past month. 7.5% (N = 1398) of students "often" or "nearly always" experienced PLEs over the past month. The frequency of each item of CAPE-P8 was depicted in Table 2. In addition, the prevalence of suicidal ideation was 9.5% (N=1773).

Correlates of PLEs

As shown in Table 3, there were statistically significant differences between participants with and without frequent PLEs in the social-demographic characteristics variables, namely sex, age, grade, ethnicity, parental marital status, residence location, left-behind experiences, single child status, father's education, mother's education, family economic status, chronic physical illness, family history of mental disorder, and body mass index (all p < 0.01). Participants with frequent PLEs were found to have a higher likelihood of sustaining sleep disturbance, depression, and anxiety and having suicidal ideation (all p < 0.001), compared with participants without frequent PLEs.

After controlling for confounding factors in the binary logistic regression analysis, the following variables were still significant (see Table 4): males (OR = 1.44, 95% CI = $1.27 \sim 1.64$), age (OR = 0.88, 95% CI = $0.84 \sim 0.93$), residence location in town (OR = 0.83, 95% CI = $0.71 \sim 0.98$), left-behind experiences (OR = 1.37, 95% CI = $1.17 \sim 1.60$), poor family economic

| Characteristics | | Ν | % |
|---|----------------------------------|--------|------|
| Sex | Female | 12,647 | 68.I |
| | Male | 5931 | 31.9 |
| Age [year, Mean (SD)] | 20.07(1.63) | | |
| Grade | Freshman | 6448 | 34.7 |
| | Sophomore | 4653 | 25.0 |
| | Junior | 3911 | 21.1 |
| | Senior | 3566 | 19.2 |
| Ethnicity | Han ^a | 18,257 | 98.3 |
| | Others | 321 | 1.7 |
| Parental marital status | Married | 16,979 | 91.4 |
| | Not current married ^b | 1599 | 8.6 |
| Residence location | Rural | 9332 | 50.2 |
| | Town | 4790 | 25.8 |
| | Urban | 4456 | 24.0 |
| Left-behind experiences ^c | Yes | 2664 | 14.3 |
| Single child status | Yes | 4018 | 21.6 |
| Father's education | Primary school or below | 3427 | 18.4 |
| | Junior high school | 8690 | 46.8 |
| | Senior high school | 4120 | 22.2 |
| | College or above | 2341 | 12.6 |
| Mother's education | Primary school or below | 4022 | 21.6 |
| | Junior high school | 8904 | 47.9 |
| | Senior high school | 3652 | 19.7 |
| | College or above | 2000 | 10.8 |
| Family economic status | Good | 839 | 4.5 |
| | Fair | 13,605 | 73.2 |
| | Poor | 4134 | 22.3 |
| Chronic physical illness ^d | Yes | 488 | 2.6 |
| Family history of mental disorder | Yes | 226 | 1.2 |
| Body mass index (kg/m ²) ^e | <18.5[underweight] | 3384 | 18.2 |
| | 18.5~23.9[normal weight] | 10,378 | 55.9 |
| | 24~27.9[overweight] | 2552 | 13.7 |
| | ≥28[obesity] | 2264 | 12.2 |

Table I Social-Demographic Characteristics of the Participants (N = 18,578)

Notes: ^aHan is the major ethnic group in China. ^bNot current married included separated, divorced, and widowed. ^cLive separately from one or both parents for more than 6 months before 16 years old. ^dChronic physical conditions referred to having at least one of arthritis, angina, asthma, diabetes, visual impairment, or hearing problems. ^eBody mass index categories were defined refer to the Chinese criteria for adults.

status (OR = 1.42, 95% CI = 1.03 ~ 1.94), chronic physical illness (OR = 1.41, 95% CI = 1.06 ~ 1.88), family history of mental disorder (OR = 1.53, 95% CI = 1.03 ~ 2.27), BMI \ge 28 [obesity] (OR = 1.25, 95% CI = 1.04 ~ 1.49), sleep disturbance (OR = 1.90, 95% CI = 1.64 ~ 2.21), depression (OR = 2.48, 95% CI = 2.09 ~ 2.94), anxiety (OR = 4.28, 95% CI = 3.59 ~ 5.09), and suicidal ideation (OR = 2.05, 95% CI = 1.76 ~ 2.39). The adjusted R square for this model is 26.4%.

Suicidal Ideation in Participants with PLEs

Table 5 presented a comparison of the socio-demographics and clinical characteristics between participants with frequent PLEs who exhibited suicidal behaviors and those who did not. The prevalence of suicidal ideation among frequent PLEs participants was 32.1% (449/1398). Compared with frequent PLEs participants without suicidal ideation, those with suicidal ideation were found to be younger, be in lower grades, have left-behind experiences and chronic physical illness, and suffer from sleep disturbance, depression, and/or anxiety (all p < 0.01).

| | Pre | valence | (%) | ≥ Often (%) | | |
|--------------------------------|---------|---------|--------|-------------|------|--------|
| | Overall | Male | Female | Overall | Male | Female |
| DEs | | | | | | |
| Delusion of reference | 31.5 | 33.2 | 30.9 | 4.2 | 5.7 | 3.5 |
| Delusion of persecution | 10.8 | 15.0 | 8.9 | 2.4 | 3.6 | 1.8 |
| Thought withdraw | 15.5 | 19.3 | 13.7 | 3.2 | 4.7 | 2.4 |
| Thought insertion | 14.7 | 18.7 | 12.8 | 2.8 | 4.2 | 2.2 |
| Thought broadcasting | 21.2 | 25.9 | 19.0 | 3.6 | 5.5 | 2.7 |
| Feeling of being control | 11.4 | 15.9 | 9.3 | 2.4 | 3.8 | 1.7 |
| Any | 39.5 | 41.6 | 38.5 | 7.1 | 9.7 | 5.9 |
| HEs | | | | | | |
| Verbal auditory hallucinations | 12.1 | 16.3 | 10.2 | 2.4 | 4.0 | 1.7 |
| Visual hallucinations | 7.0 | 11.1 | 5.1 | 1.8 | 3.0 | 1.2 |
| Any | 12.6 | 17.0 | 10.6 | 2.8 | 4.7 | 1.9 |
| PLEs | | | | | | |
| Any | 40.3 | 42.4 | 39.3 | 7.5 | 10.3 | 6.2 |

| Table 2 | The | Prevalence | of | Current | PLEs | Among | the | Current | Sample |
|---------|-----|--------------|----|---------|------|-------|-----|---------|--------|
| Table L | THE | i i cvalence | U. | Current | | Among | une | Current | Jampie |

Abbreviations: Des, delusional experiences; HEs, hallucinatory experiences; PLEs, psychotic-like experiences.

Table 3 Demographics and Clinical Characteristics Between Participants with and without PLEs [N(%)]

| Characteristics | | Frequent PLEs (N=1398, 7.5%) | Control (N=17180, 92.5%) | P value | Cramer's V/ Cohen's d |
|-------------------------|-------------------------|---------------------------------|-----------------------------|---------|--------------------------|
| Sex | Females | 786(56.2) | 11,861(69.0) | <0.001 | 0.073 |
| | Males | 612(43.8) | 5319(31.0) | | |
| Age [year, Mean (SD)] | | 19.70(1.67) | 20.10(1.62) | <0.001 | 0.243 |
| Grade | Freshman | 578(41.3) | 5870(34.2) | <0.001 | 0.056 |
| | Sophomore | 391 (28.0) | 4262(24.8) | | |
| | Junior | 231(16.5) | 3680(21.4) | | |
| | Senior | 198(14.2) | 3368(19.6) | | |
| Ethnicity | Han ^a | 1353(96.8) | 16,904(98.4) | <0.001 | 0.033 |
| | Others | 45(3.2) | 276(1.6) | | |
| Parental marital status | Married | 1232(88.1) | 15,747(91.7) | <0.001 | 0.033 |
| | Not current married | 166(11.9) | 1433(8.3) | | |
| Residence location | Rural | 722(51.6) | 8610(50.1) | 0.002 | 0.026 |
| | Town | 307(22.0) | 4483(26.1) | | |
| | Urban | 369(26.4) | 4087(23.8) | | |
| Left-behind experiences | Yes | 357(25.5) | 2307(13.4) | <0.001 | 0.091 |
| Single child status | Yes | 353(25.3) | 3665(21.3) | 0.001 | 0.025 |
| Father's education | Primary school or below | 275(19.7) | 3152(18.3) | <0.001 | 0.036 |
| | Junior high school | 574(41.1) | 8116(47.2) | | |
| | Senior high school | 332(23.7) | 3788(22.0) | | |
| | College or above | 217(15.5) | 2124(12.4) | | |
| Mother's education | Primary school or below | 330(23.6) | 3692(21.5) | <0.001 | 0.038 |
| | Junior high school | 592(42.3) | 8312(48.4) | | |
| | Senior high school | 282(20.2) | 3370(19.6) | | |
| | College or above | 194(13.9) | 1806(10.5) | | |
| Family economic status | Good | 67(4.8) | 772(4.5) | <0.001 | 0.088 |
| | Fair | 842(60.2) | 12,763(74.3) | | |
| | Poor | 489(35.0) | 3645(21.2) | | |

(Continued)

Table 3 (Continued).

| Characteristics | | Frequent PLEs (N=1398, 7.5%) | Control (N=17180, 92.5%) | P value | Cramer's V/ Cohen's d |
|--------------------------------------|--------------------------|---------------------------------|-----------------------------|---------|--------------------------|
| Chronic physical illness | Yes | 82(5.9) | 406(2.4) | <0.001 | 0.058 |
| Family history of mental disorder | Yes | 46(3.3) | 180(1.0) | <0.001 | 0.054 |
| Body mass index (kg/m ²) | <18.5[underweight] | 237(17.0) | 3147(18.3) | 0.003 | 0.027 |
| | 18.5~23.9[normal weight] | 748(53.5) | 9630(56.1) | | |
| | 24~27.9[overweight] | 202(14.4) | 2350(13.7) | | |
| | ≥28[obesity] | 211(15.1) | 2053(11.9) | | |
| Sleep disturbance | Yes | 445(31.8) | 1547(9.0) | <0.001 | 0.195 |
| Depression | Yes | 703(50.3) | 1581(9.2) | <0.001 | 0.330 |
| Anxiety | Yes | 668(49.2) | 1207(7.0) | <0.001 | 0.368 |
| Suicidal ideation | Yes | 449(32.1) | 1324(7.7) | <0.001 | 0.219 |

| Table 4 Multivariate | Analysis for | Variables | Associated | with Fre | auent PLEs |
|----------------------|--------------|------------|------------|----------|-------------|
| | | 141.140.00 | | | -qaoe . ==o |

| Characteristics | | | 95% CI | P value |
|--------------------------------------|------------------------|------|-----------|---------|
| Sex | Female | 1.00 | | |
| | Male | 1.44 | 1.27~1.64 | <0.001 |
| Age [year, Mean (SD)] | | 0.88 | 0.84~0.93 | <0.001 |
| Grade | Freshman | 1.00 | | |
| | Sophomore | 1.09 | 0.93~1.28 | 0.294 |
| | Junior | 0.92 | 0.75~1.12 | 0.411 |
| | Senior | 0.97 | 0.77~1.22 | 0.791 |
| Ethnicity | Han | 1.00 | | |
| | Others | 1.45 | 1.00~2.11 | 0.050 |
| Parental marital status | Married | 1.00 | | |
| | Not current married | 1.00 | 0.82~1.23 | 0.990 |
| Residence location | Rural | 1.00 | | |
| | Town | 0.83 | 0.71~0.98 | 0.025 |
| | Urban | 1.07 | 0.90~1.27 | 0.433 |
| Left-behind experiences | Yes | 1.37 | 1.17~1.60 | <0.001 |
| Single child status | Yes | 1.02 | 0.87~1.19 | 0.756 |
| Father's education | Primary school or less | 1.00 | | |
| | Junior high school | 0.89 | 0.74~1.06 | 0.192 |
| | Senior high school | 1.13 | 0.91~1.39 | 0.268 |
| | College or more | 1.14 | 0.88~1.48 | 0.316 |
| Mother's education | Primary school or less | 1.00 | | |
| | Junior high school | 0.95 | 0.80~1.12 | 0.535 |
| | Senior high school | 1.03 | 0.83~1.27 | 0.792 |
| | College or more | 1.14 | 0.86~1.47 | 0.381 |
| Family economic status | Good | 1.00 | | |
| | Fair | 1.01 | 0.75~1.37 | 0.951 |
| | Poor | 1.42 | 1.03~1.94 | 0.034 |
| Chronic physical illness | Yes | 1.41 | 1.06~1.88 | 0.020 |
| Family history of mental disorder | Yes | 1.53 | 1.03~2.27 | 0.034 |
| Body mass index (kg/m ²) | 18.5~23.9[normal | 1.00 | | |
| | weight] | | | |
| | <18.5[underweight] | 1.05 | 0.88~1.24 | 0.610 |
| | 24~27.9[overweight] | 1.02 | 0.85~1.22 | 0.850 |
| | ≥28[obesity] | 1.25 | 1.04~1.49 | 0.018 |

(Continued)

| Table 4 | Continued |). |
|---------|-----------|----|
|---------|-----------|----|

| Characteristics | | OR | 95% CI | P value |
|-------------------|-----|------|-----------|---------|
| Sleep disturbance | Yes | 1.90 | 1.64~2.21 | <0.001 |
| Depression | Yes | 2.48 | 2.09~2.94 | <0.001 |
| Anxiety | Yes | 4.28 | 3.59~5.09 | <0.001 |
| Suicidal ideation | Yes | 2.05 | 1.76~2.39 | <0.001 |

Note: Bold type indicates a significant odds ratio.

Abbreviations: OR, odds ratio; CI, confidence interval.

A binary logistic regression analysis further illustrated that sleep disturbance (OR = 1.80, 95% CI = $1.40 \sim 2.32$), depression (OR = 1.63, 95% CI = $1.21 \sim 2.21$), and anxiety (OR = 2.26, 95% CI = $1.21 \sim 2.21$) remained significant after controlling for confounding factors (see Table 6). Grade was also negatively associated with suicidal ideation (OR = $0.43 \sim 0.65$). The adjusted R square for this model is 15.9%.

Table 5 Demographics and Clinical Characteristics Between Frequent PLEs Participants with and without Suicidal Ideation

| Characteristics | | With SI (N=449, 32.1%) | Without SI (N=949, 67.9%) | P value | Cramer's V/ Cohen's d |
|--------------------------------------|--------------------------|---------------------------|------------------------------|---------|--------------------------|
| Sex | Female | 264(58.8) | 522(55.0) | 0.185 | - |
| | Male | 185(41.2) | 427(45.0) | | |
| Age [year, Mean (SD)] | | 19.52(1.65) | 19.79(1.68) | 0.005 | 0.162 |
| Grade | Freshman | 217(48.3) | 361(38.0) | 0.001 | 0.109 |
| | Sophomore | 119(26.5) | 272(28.7) | | |
| | Junior | 67(14.9) | 164(17.3) | | |
| | Senior | 46(10.2) | 152(16.0) | | |
| Ethnicity | Han | 434(96.7) | 919(96.8) | 0.872 | - |
| | Others | 15(3.3) | 30(3.2) | | |
| Parental marital status | Married | 389(86.6) | 843(88.8) | 0.250 | - |
| | Not current married | 60(13.4) | 106(11.2) | | |
| Residence location | Rural | 233(51.9) | 489(51.5) | 0.992 | - |
| | Town | 98(21.8) | 209(22.0) | | |
| | Urban | 118(26.3) | 251(26.4) | | |
| Left-behind experiences | Yes | 139(31.0) | 218(23.0) | 0.002 | 0.086 |
| Single child status | Yes | 113(25.2) | 240(25.3) | 1.000 | - |
| Father's education | Primary school or below | 99(22.0) | 176(18.5) | 0.210 | - |
| | Junior high school | 186(41.4) | 388(40.9) | | |
| | Senior high school | 93(20.7) | 239(25.2) | | |
| | College or above | 71(15.8) | 146(15.4) | | |
| Mother's education | Primary school or below | 105(23.4) | 225(23.7) | 0.999 | - |
| | Junior high school | 190(42.3) | 402(42.4) | | |
| | Senior high school | 91(20.3) | 191(20.1) | | |
| | College or above | 63(14.0) | 3 (3.8) | | |
| Family economic status | Good | 22(4.9) | 45(4.7) | 0.396 | - |
| | Fair | 259(57.7) | 583(61.4) | | |
| | Poor | 168(37.4) | 321(33.8) | | |
| Chronic physical illness | Yes | 39(8.7) | 43(4.5) | 0.003 | 0.083 |
| Family history of mental disorder | Yes | 20(4.5) | 26(2.7) | 0.108 | - |
| Body mass index (kg/m ²) | <18.5[underweight] | 74(16.5) | 163(17.2) | 0.585 | - |
| | 18.5~23.9[normal weight] | 251(55.9) | 497(52.4) | | |
| | 24~27.9[overweight] | 58(12.9) | 144(15.2) | | |
| | ≥28[obesity] | 66(14.7) | 145(15.3) | | |
| Sleep disturbance | Yes | 203(45.2) | 242(25.5) | <0.001 | 0.198 |
| Depression | Yes | 302(67.3) | 401(42.3) | <0.001 | 0.234 |
| Anxiety | Yes | 307(68.4) | 381(40.1) | <0.001 | 0.264 |

| Characteristics | OR | 95% CI | P value | |
|--------------------------|-----------|--------|-----------|--------|
| Age | | 0.99 | 0.90~1.08 | 0.751 |
| Grade | Freshman | 1.00 | | |
| | Sophomore | 0.65 | 0.48~0.89 | 0.007 |
| | Junior | 0.61 | 0.41~0.91 | 0.015 |
| | Senior | 0.43 | 0.27~0.70 | 0.001 |
| Left-behind experiences | Yes | 1.28 | 0.98~1.68 | 0.074 |
| Chronic physical illness | Yes | 1.59 | 0.99~2.58 | 0.057 |
| Sleep disturbance | Yes | 1.80 | 1.40~2.32 | <0.001 |
| Depression | Yes | 1.63 | 1.21~2.21 | 0.001 |
| Anxiety | Yes | 2.26 | 1.67~3.07 | <0.001 |

Table 6 Multivariate Analysis for Variables Associated with SuicidalIdeation Among Participants with Frequent PLEs

Note: Bold type indicates a significant odds ratio.

Abbreviations: OR, odds ratio; Cl, confidence interval.

Discussion

This study intended to examine the prevalence of PLEs and suicidal ideation among Chinese college students, as well as to investigate the demographic and clinical factors associated with these experiences. Our study indicated that 40.3% of college students had PLEs over the past month, and 7.5% experienced frequent current PLEs. The prevalence rates observed in our study are higher compared to the findings of a previous survey conducted with a small sample of college students (N=2231, 36.3% and 3.3%).¹⁶ Nevertheless, the rates are lower than the results from a large-scale survey of junior and senior high school students (N=67538, 49.3% and 15.4%)¹⁴ with the same scale and screening criteria. The variation in prevalence rates can be attributed to the difference in the sampled populations. In our study, 9.5% of participants reported suicidal ideation in the preceding month. However, the prevalence rate of suicidal ideation also varies greatly in different measures, samples, and sampling time. For instance, one study using the Epidemiological version of the Kiddie Schedule for Affective Disorders and Schizophrenia (Kiddie-SADS-E) investigated 9510 Taiwanese adolescents (48.3% boys, mean age 14.69 years) and displayed that 18.03% of participants have suicidal ideation over the past month.⁵¹ Zarrouq et al employed the Mini International Neuropsychiatric Interview (MINI) to assess one-month suicidal behaviors among 3020 Moroccan school students (53.0% boys, mean age 16 years) and found that the prevalence rate of suicidal ideation during the last month was 15.7%.⁵²

In this study, male students are more likely to suffer frequent PLEs, which is consistent with prior findings.⁵³ However, the finding is opposed to some studies that argue female adolescents had a higher prevalence of PLEs.^{17,54} One possible reason for the results of this study is that male college students are more prone to adopting unhealthy lifestyles, such as Internet addiction⁵⁵ and smoking,⁵⁶ which may contribute to PLEs.^{57,58} Meanwhile, age served as a protective factor against frequent PLEs. In fact, PLEs are transient in nature, and the majority of PLEs (75 \sim 90%) during early adolescence gradually diminish with age.¹² Our observation of students who have experienced being left-behind revealed a significant increase in current PLEs, in line with previous research findings.^{14,41} Left-behind adolescents, as a consequence of their parents' migration and inadequate care, experience greater exposure to traumatic events (eg, emotional neglect and exploitation, school bullying),⁵⁹ and heightened feelings of loneliness.⁶⁰ These factors make them more vulnerable to mental health issues. Moreover, extensive existing studies have adequately proven that familial poverty is a vital risk factor for increased mental health issues among adolescents,⁶¹ including PLEs.⁹ This argument coincides with our finding that poor family economic status was significantly associated with a higher occurrence of frequent PLEs. This result may provide a partial explanation for the lower prevalence of PLEs reported by participants residing in urban areas, as they tend to have better family economic status compared to those from rural areas in this study. According to our findings, participants residing in towns and urban areas reported a higher risk of frequent PLEs compared to those residing in rural areas. Though it is true that families in urban are generally better off economically, it should be acknowledged that urbanization may also have detrimental effects on the psychological development of individuals, including limited access to green spaces, residential segregation, social disorganization, and other concentrated disadvantages (ie, unemployment).^{62,63} Furthermore, our finding further illustrated that students with obesity are more likely to have frequent current PLEs. Obesity might involve hypothalamic-pituitary-adrenal axis (HPA axis) dysregulation,^{64,65} which is well known to be involved in psychosis.⁶⁶ We also speculated that obese individuals are less inclined to engage in physical exercise, which may result in an increased risk of PLEs.^{14,42} Coinciding with previous work,¹⁴ it has been validated that chronic physical illness and family history of mental disorder are independently related to frequent PLEs. Thus, these factors should also be taken into consideration for implementing psychosocial interventions among college students.

We demonstrated that college students with frequent PLEs had a higher risk of sleep disturbance, depression, and anxiety than those without frequent PLEs. Lee et al investigated the relationship between PLEs and sleep disturbance among adolescents and found that sleep disturbance was independently correlated with the increased likelihood of PLEs.⁶⁷ Other studies conducted among adolescents also affirmed the association between PLEs and depressive and anxiety.^{20,22} This association could be explained by their congenerous risk factors, such as negative life events.^{68–70} Controlled for other confounding factors, our results revealed that participants with frequent PLEs were more than twice as likely to have suicidal ideation (OR=2.05). The strong relationship between PLEs and suicidal ideation was consistent with previous studies.^{16,71} This relationship could be attributed to increased affective reactivity and inadequate coping skills to negative stressors in individuals with PLEs,^{72,73} which in turn contribute to an increase risk of suicidal ideation.

To our knowledge, scant attention has been paid to independent predictors of suicidal ideation among adolescents with PLEs. In this study, we found that students with frequent PLEs were at risk for sleep disturbance, depression, and anxiety, with proportions of 31.8%, 50.3%, and 49.2%, respectively. This prevalence rate increased significantly among frequent PLEs participants who had suicidal ideation (sleep disturbance, 45.2%; depression, 67.3%; anxiety, 68.4%). Several literatures, including this study, have identified that PLEs are independently associated with suicidal ideation.^{16,33,71} Therefore, sleep disturbance and emotional symptoms should be directly related to both PLEs and suicide behavior. Additionally, higher grades were associated with a lower risk of suicidal ideation among participants who reported PLEs frequently. This finding contradicts the general findings on suicidal ideation among college students.⁷⁴ One possible explanation for this is that among college students with frequent PLEs, students in higher grades may have greater access to social support and are easier to seek help when they are in trouble.

The present study makes notable contributions in terms of methodology, such as a large sample size and comprehensive measurement of clinical characteristics. This study also has several limitations that may be reflection points for future research. Firstly, PLEs and other clinical characteristics in this study were assessed through self-report. Especially suicidal ideation was measured with only a single item, which may lead to certain reporting biases. Also, to minimize participant burden, we opted to measure depressive and anxiety symptoms with the PHQ-2 and GAD-2. However, it should be noted that PHQ-2 and GAD-2 may not be as comprehensive or accurate as the PHQ-9 and GAD-7 in accessing the symptoms.⁷⁵ Secondly, this study employed a cross-sectional design and no further follow-up was conducted with the high-risk group, which may limit our ability to infer causality. Therefore, it is necessary to further conduct longitudinal research to explore the relationship between PLEs and other clinical symptoms. Finally, there are several factors related to the mental health of college students that have not been taken into account, such as the study field⁷⁶ and life events.⁶⁸ Future studies can attempt to unearth these influential factors.

Conclusion

This study demonstrated that PLEs was prevalent among Chinese college students. The independent relationship between PLEs and suicidal ideation were also proven. Our findings highlighted the importance of screening college students with PLEs. We also identified several factors associated with frequent PLEs, which broadened the theoretical and empirical basis concerning PLEs and provided new insights into the pathomechanisms underlying PLEs. This study also provided valuable data for healthcare professionals and school administrators to refer to when assessing the risks of suicide among individuals with frequent PLEs. Further longitudinal studies in the future are necessary to gain a deeper understanding of the relationship between these factors and PLEs.

Data Sharing Statement

The data presented in this study are available on request from the corresponding authors (Dr. Dongfang Wang).

Ethics Statement

The investigation was carried out in accordance with the Helsinki Declaration as revised 1989 and approved by the Ethics Committee of the School of Psychology, South China Normal University. Informed consent was obtained from all participants involved in the study.

Funding

This research was funded by the youth project of the Social Science Fund of Fujian Province (FJ2023C018).

Disclosure

The authors declare that they have no conflicts of interest in this work.

References

- 1. Polanczyk GV, Salum GA, Sugaya LS, Caye A, Rohde LA. Annual research review: a meta-analysis of the worldwide prevalence of mental disorders in children and adolescents. *J Child Psychol Psychiatr*. 2015;56(3):345–365. doi:10.1111/jcpp.12381
- 2. Kelleher I, Cannon M. Psychotic-like experiences in the general population: characterizing a high-risk group for psychosis. *Psychol Med.* 2011;41 (1):1–6. doi:10.1017/S0033291710001005
- 3. Poulton R, Caspi A, Moffitt TE, Cannon M, Murray R, Harrington H. Children's self-reported psychotic symptoms and adult schizophreniform disorder: a 15-year longitudinal study. *Arch Gen Psychiatry*. 2000;57(11):1053–1058. doi:10.1001/archpsyc.57.11.1053
- Dominguez MD, Wichers M, Lieb R, Wittchen HU, van Os J. Evidence that onset of clinical psychosis is an outcome of progressively more persistent subclinical psychotic experiences: an 8-year cohort study. Schizophr Bull. 2011;37(1):84–93. doi:10.1093/schbul/sbp022
- 5. Healy C, Brannigan R, Dooley N, et al. Childhood and adolescent psychotic experiences and risk of mental disorder: a systematic review and meta-analysis. *Psychol Med.* 2019;49(10):1589–1599. doi:10.1017/S0033291719000485
- Rimvall MK, van Os J, Verhulst F, et al. Mental Health Service Use and Psychopharmacological Treatment Following Psychotic Experiences in Preadolescence. Am J Psychiatry. 2020;177(4):318–326. doi:10.1176/appi.ajp.2019.19070724
- 7. Kelleher I, Cannon M. A neural efficiency-threshold model to understand psychotic experiences. *Psychol Med.* 2021;51(11):1777–1782. doi:10.1017/S0033291721001495
- 8. Kelleher I, Cannon M. Putting psychosis in its place. Am J Psychiatry. 2016;173(10):951-952. doi:10.1176/appi.ajp.2016.16070810
- 9. Mamah D, Mutiso VN, Ndetei DM. Psychotic-like experiences among 9564 Kenyan adolescents and young adults. *Psychiatry Res.* 2021;302:113994. doi:10.1016/j.psychres.2021.113994
- Fekih-Romdhane F, Sassi H, Ennaifer S, Tira S, Cheour M. Prevalence and correlates of psychotic like experiences in a large community sample of young adults in Tunisia. *Commu Ment Health J.* 2020;56(6):991–1003. doi:10.1007/s10597-019-00542-1
- 11. Sun M, Hu X, Zhang W, et al. Psychotic-like experiences and associated socio-demographic factors among adolescents in China. *Schizophr Res.* 2015;166(1–3):49–54. doi:10.1016/j.schres.2015.05.031
- 12. van Os J, Linscott RJ, Myin-Germeys I, Delespaul P, Krabbendam L. A systematic review and meta-analysis of the psychosis continuum: evidence for a psychosis proneness-persistence-impairment model of psychotic disorder. *Psychol Med.* 2009;39(2):179–195. doi:10.1017/S0033291708003814
- 13. Capra C, Kavanagh DJ, Hides L, Scott J. Current CAPE-15: a measure of recent psychotic-like experiences and associated distress. *Early Interv Psychiatry*. 2017;11(5):411–417. doi:10.1111/eip.12245
- 14. Wang D, Chen H, Chen Z, et al. Current psychotic-like experiences among adolescents in China: identifying risk and protective factors. *Schizophr Res.* 2022;244:111–117. doi:10.1016/j.schres.2022.05.024
- Sun M, Wang D, Jing L, Zhou L. Changes in psychotic-like experiences and related influential factors in technical secondary school and college students during COVID-19. Schizophr Res. 2021;231:3–9. doi:10.1016/j.schres.2021.02.015
- Luo X, Yu T, Yang Z, Wang D. Psychotic-like experiences and suicidal ideation among adolescents: the chain mediating role of insomnia symptoms and resilience. *Psychol Res Behav Manag.* 2023;16:3519–3530. doi:10.2147/PRBM.S426363
- 17. Wang D, Zhou L, Chen C, Sun M. Psychotic-like experiences during COVID-19 lockdown among adolescents: prevalence, risk and protective factors. *Schizophr Res.* 2023;252:309–316. doi:10.1016/j.schres.2023.01.027
- 18. Wu Z, Liu D, Zhang J, et al. Sex difference in the prevalence of psychotic-like experiences in adolescents: results from a pooled study of 21,248 Chinese participants. *Psychiatry Res.* 2022;317:114894. doi:10.1016/j.psychres.2022.114894
- 19. Wang D, Ma Z, Scherffius A, et al. Sleep disturbance is predictive of psychotic-like experiences among adolescents: a two-wave longitudinal survey. *Sleep Med.* 2023;101:296–304. doi:10.1016/j.sleep.2022.11.011
- 20. Wu Z, Liu Z, Zou Z, et al. Changes of psychotic-like experiences and their association with anxiety/depression among young adolescents before COVID-19 and after the lockdown in China. *Schizophr Res.* 2021;237:40–46. doi:10.1016/j.schres.2021.08.020
- 21. Ered A, Cooper S, Ellman LM. Sleep quality, psychological symptoms, and psychotic-like experiences. J Psychiatr Res. 2018;98:95–98. doi:10.1016/j.jpsychires.2017.12.016
- 22. Yamasaki S, Usami S, Sasaki R, et al. The association between changes in depression/anxiety and trajectories of psychotic-like experiences over a year in adolescence. *Schizophr Res.* 2018;195:149–153. doi:10.1016/j.schres.2017.10.019
- 23. Ahmed HU, Hossain MD, Aftab A, et al. Suicide and depression in the World Health Organization South-East Asia Region: a systematic review. WHO South East Asia J Public Health. 2017;6(1):60–66. doi:10.4103/2224-3151.206167

- 24. Bentley KH, Franklin JC, Ribeiro JD, Kleiman EM, Fox KR, Nock MK. Anxiety and its disorders as risk factors for suicidal thoughts and behaviors: a meta-analytic review. *Clin Psychol Rev.* 2016;43:30–46. doi:10.1016/j.cpr.2015.11.008
- 25. Wang D, Ross B, Zhou X, et al. Sleep disturbance predicts suicidal ideation during COVID-19 pandemic: a two-wave longitudinal survey. *J Psychiatr Res.* 2021;143:350–356. doi:10.1016/j.jpsychires.2021.09.038
- 26. Patton GC, Coffey C, Sawyer SM, et al. Global patterns of mortality in young people: a systematic analysis of population health data. *Lancet*. 2009;374(9693):881-892. doi:10.1016/S0140-6736(09)60741-8
- 27. Centers for Disease Control and Prevention (CDC). Suicide and attempted suicide–China, 1990–2002. MMWR Morb Mortal Wkly Rep. 2004;53 (22):481–484.
- 28. Phillips MR, Li X, Zhang Y. Suicide rates in China, 1995–99. Lancet. 2002;359:9309):835–840. doi:10.1016/S0140-6736(02)07954-0
- 29. Wang CW, Chan CL, Yip PS. Suicide rates in China from 2002 to 2011: an update. Soc Psychiatry Psychiatr Epidemiol. 2014;49(6):929–941. doi:10.1007/s00127-013-0789-5
- Wang SY, Li YH, Chi GB, et al. Injury-related fatalities in China: an under-recognised public-health problem. *Lancet*. 2008;372(9651):1765–1773. doi:10.1016/S0140-6736(08)61367-7
- 31. Yates K, Lang U, Cederlof M, et al. Association of psychotic experiences with subsequent risk of suicidal ideation, suicide attempts, and suicide deaths: a systematic review and meta-analysis of longitudinal population studies. JAMA Psychiatry. 2019;76(2):180–189. doi:10.1001/jamapsychiatry.2018.3514
- 32. Hielscher E, DeVylder JE, Saha S, Connell M, Scott JG. Why are psychotic experiences associated with self-injurious thoughts and behaviours? A systematic review and critical appraisal of potential confounding and mediating factors. *Psychol Med.* 2018;48(9):1410–1426. doi:10.1017/S0033291717002677
- 33. Grattan RE, Karcher NR, Maguire AM, Hatch B, Barch DM, Niendam TA. Psychotic like experiences are associated with suicide ideation and behavior in 9 to 10 year old children in the United States. *Res Child Adolesc Psychopathol.* 2021;49(2):255–265. doi:10.1007/s10802-020-00721-9
- Zhou R, Foo JC, Nishida A, Ogawa S, Togo F, Sasaki T. Longitudinal relationships of psychotic-like experiences with suicidal ideation and self-harm in adolescents. *Eur Child Adolesc Psychiatry*. 2023. doi:10.1007/s00787-023-02299-1
- 35. Klonsky ED, Pachkowski MC, Shahnaz A, May AM. The three-step theory of suicide: description, evidence, and some useful points of clarification. *Prev Med.* 2021;152(1):106549. doi:10.1016/j.ypmed.2021.106549
- 36. UNICEF. Population status of children in China in 2015: facts and figures; 2018. Available from: https://www.unicef.cn/atlas-2018-cn/. Accessed February 24, 2020.
- 37. Koyanagi A, Stubbs B, Lara E, et al. Psychotic experiences and subjective cognitive complaints among 224 842 people in 48 low- and middle-income countries. *Epidemiol Psychiatr Sci.* 2018;29:e11. doi:10.1017/S2045796018000744
- Wang D, Sun M, Ouyang X, Fan F. Validity and reliability of the simplified version of community assessment of psychic experiences in college students. *Chin Mental Health J.* 2022;36(2):172–178.
- 39. Wang D, Sun M, Xi C, et al. Gender and longitudinal measurement invariance of the community assessment of psychic experiences-positive subscale. *Chin J Clin Psych*. 2020;28(1):41–45. doi:10.16128/j.cnki.1005-3611.2020.01.010
- 40. Wang D, Ma Z, Fan Y, Chen H, Sun M, Fan F. Tobacco smoking, second-hand smoking exposure in relation to psychotic-like experiences in adolescents. *Early Interv Psychiatry*. 2023. doi:10.1111/eip.13439
- 41. Sun M, Zhang W, Guo R, et al. Psychotic-like experiences and correlation with childhood trauma and other socio-demographic factors: a cross-sectional survey in adolescence and early adulthood in China. *Psychiatry Res.* 2017;255:272–277. doi:10.1016/j.psychres.2017.03.059
- 42. Wang D, Ma Z, Zhai S, Sun M, Fan F. Sleep disturbance and psychotic-like experiences among urban adolescents with and without parental migration. *Front Public Health*. 2022;10:1037963. doi:10.3389/fpubh.2022.1037963
- 43. Kroenke K, Spitzer RL, Williams JB. The patient health questionnaire-2: validity of a two-item depression screener. *Med Care*. 2003;41 (11):1284–1292. doi:10.1097/01.MLR.0000093487.78664.3C
- 44. Arrieta J, Aguerrebere M, Raviola G, et al. Validity and Utility of the Patient Health Questionnaire (PHQ)-2 and PHQ-9 for screening and diagnosis of depression in rural chiapas, Mexico: a cross-sectional study. J Clin Psychol. 2017;73(9):1076–1090. doi:10.1002/jclp.22390
- 45. Hughes AJ, Dunn KM, Chaffee T, Bhattarai JJ, Beier M. Diagnostic and Clinical Utility of the GAD-2 for screening anxiety symptoms in individuals with multiple sclerosis. Arch Phys Med Rehabil. 2018;99(10):2045–2049. doi:10.1016/j.apmr.2018.05.029
- 46. Plummer F, Manea L, Trepel D, McMillan D. Screening for anxiety disorders with the GAD-7 and GAD-2: a systematic review and diagnostic metaanalysis. *Gen Hosp Psychiatry*. 2016;39:24–31. doi:10.1016/j.genhosppsych.2015.11.005
- 47. Klonsky ED, May AM. The Three-Step Theory (3ST): a New Theory of Suicide Rooted in the "Ideation-to-Action" framework. *Int J Cogn Ther*. 2015;8(2):114–129. doi:10.1521/ijct.2015.8.2.114
- Klonsky ED, May AM, Saffer BY. Suicide, Suicide Attempts, and Suicidal Ideation. Annu Rev Clin Psychol. 2016;12:307–330. doi:10.1146/ annurev-clinpsy-021815-093204
- 49. O'Brien R. A caution regarding rules of thumb for variance inflation factors. Qual Quantity. 2007;41(5):673-690. doi:10.1007/s11135-006-9018-6
- Wang D, Chen H, Chen J, et al. A comparative study of mental health status among left-behind, migrant, and local adolescents in China. J Affect Disord. 2023;324:521-528. doi:10.1016/j.jad.2022.12.134
- 51. Lin IH, Ko CH, Chang YP, et al. The association between suicidality and Internet addiction and activities in Taiwanese adolescents. *Compr Psychiatry*. 2014;55(3):504–510. doi:10.1016/j.comppsych.2013.11.012
- 52. Zarrouq B, Bendaou B, Elkinany S, et al. Suicidal behaviors among Moroccan school students: prevalence and association with socio-demographic characteristics and psychoactive substances use: a cross-sectional study. *BMC Psychiatry*. 2015;15:284. doi:10.1186/s12888-015-0680-x
- 53. Laurens KR, Hodgins S, Maughan B, Murray RM, Rutter ML, Taylor EA. Community screening for psychotic-like experiences and other putative antecedents of schizophrenia in children aged 9–12 years. *Schizophr Res.* 2007;90:130–146. doi:10.1016/j.schres.2006.11.006
- 54. Stainton A, Chisholm K, Woodall T, et al. Gender differences in the experience of psychotic-like experiences and their associated factors: a study of adolescents from the general population. Schizophr Res. 2021;228:410–416. doi:10.1016/j.schres.2021.01.008
- 55. Lin X, Gu JY, Guo WJ, et al. The gender-sensitive social risk factors for internet addiction in college undergraduate students. *Psychiatry Invest*. 2021;18(7):636–644. doi:10.30773/pi.2020.0277
- 56. Li X, Tan Y, Li S, Wang X. Psychological distress and smoking behaviors of Chinese college students: mediating effects of the dimensions of learning burnout. *BMC Psychol*. 2022;10(1):125. doi:10.1186/s40359-022-00840-6

- Mittal VA, Dean DJ, Pelletier A. Internet addiction, reality substitution and longitudinal changes in psychotic-like experiences in young adults. Early Interv Psychiatry. 2013;7(3):261–269. doi:10.1111/j.1751-7893.2012.00390.x
- Mallet J, Mazer N, Dubertret C, Le Strat Y. Tobacco smoking and psychotic-like experiences in a general population sample. J Clin Psychiatry. 2018;79(6):17m11994. doi:10.4088/JCP.17m11994
- 59. Givaudan M, Pick S. Children left behind: how to mitigate the effects and facilitate emotional and psychosocial development: supportive community networks can diminish the negative effects of parental migration. *Child Abuse Negl.* 2013;37(12):1080–1090. doi:10.1016/j. chiabu.2013.10.019
- 60. Jia Z, Tian W. Loneliness of left-behind children: a cross-sectional survey in a sample of rural China. *Child Care Health Dev.* 2010;36(6):812–817. doi:10.1111/j.1365-2214.2010.01110.x
- 61. Reiss F. Socioeconomic inequalities and mental health problems in children and adolescents: a systematic review. Soc Sci Med. 2013;90:24–31. doi:10.1016/j.socscimed.2013.04.026
- McKenzie K, Murray A, Booth T. Do urban environments increase the risk of anxiety, depression and psychosis? An epidemiological study. J Affect Disord. 2013;150(3):1019–1024. doi:10.1016/j.jad.2013.05.032
- 63. Galea S, Uddin M, Koenen K. The urban environment and mental disorders: epigenetic links. *Epigenetics*. 2011;6(4):400-404. doi:10.4161/ epi.6.4.14944
- 64. Pasquali R, Vicennati V. Activity of the hypothalamic-pituitary-adrenal axis in different obesity phenotypes. Int J Obes Relat Metab Disord. 2000;24:2.
- 65. Walker BR. Activation of the hypothalamic-pituitary-adrenal axis in obesity: cause or consequence? *Growth Horm IGF Res.* 2001;11(Suppl A): S91–S95. doi:10.1016/s1096-6374(01)80015-0
- 66. Shah JL, Malla AK. Much ado about much: stress, dynamic biomarkers and HPA axis dysregulation along the trajectory to psychosis. Schizophr Res. 2015;162(1-3):253–260. doi:10.1016/j.schres.2015.01.010
- 67. Lee YJ, Cho SJ, Cho IH, Jang JH, Kim SJ. The relationship between psychotic-like experiences and sleep disturbances in adolescents. *Sleep Med*. 2012;13(8):1021–1027. doi:10.1016/j.sleep.2012.06.002
- Yang S, Huang P, Li B, Gan T, Lin W, Liu Y. The relationship of negative life events, trait-anxiety and depression among Chinese university students: a moderated effect of self-esteem. J Affect Disord. 2023;339:384–391. doi:10.1016/j.jad.2023.07.010
- 69. Lee KH, Lee HY, Park I, et al. Life stress, sleep disturbance and depressive symptoms: the moderating role of prefrontal activation during emotion regulation. Aust N Z J Psychiatry. 2022;56(6):709–720. doi:10.1177/00048674211025729
- Gibson LE, Reeves LE, Cooper S, Olino TM, Ellman LM. Traumatic life event exposure and psychotic-like experiences: a multiple mediation model of cognitive-based mechanisms. *Schizophr Res.* 2019;205:15–22. doi:10.1016/j.schres.2018.02.005
- Jang JH, Lee YJ, Cho SJ, Cho IH, Shin NY, Kim SJ. Psychotic-like experiences and their relationship to suicidal ideation in adolescents. *Psychiatry* Res. 2014;215(3):641–645. doi:10.1016/j.psychres.2013.12.046
- 72. Lataster T, Wichers M, Jacobs N, et al. Does reactivity to stress cosegregate with subclinical psychosis? A general population twin study. *Acta Psychiatr Scand*. 2009;119(1):45–53. doi:10.1111/j.1600-0447.2008.01263.x
- 73. Lin A, Wigman JT, Nelson B, et al. The relationship between coping and subclinical psychotic experiences in adolescents from the general population--a longitudinal study. *Psychol Med.* 2011;41:2535–2546. doi:10.1017/S0033291711000560
- 74. Wang M, Kou C, Bai W, et al. Prevalence and correlates of suicidal ideation among college students: a mental health survey in Jilin Province, China. J Affect Disord. 2019;246:166–173. doi:10.1016/j.jad.2018.12.055
- Anand P, Bhurji N, Williams N, Desai N. Comparison of PHQ-9 and PHQ-2 as screening tools for depression and school related stress in inner City adolescents. J Prim Care Commu Health. 2021;12:627227978. doi:10.1177/21501327211053750
- Frajerman A, Chevance A, Chaumette B, Morvan Y. Prevalence and factors associated with depression and suicidal ideation among French students in 2016: a national study. *Psychiatry Res.* 2023;326:115263. doi:10.1016/j.psychres.2023.115263

Psychology Research and Behavior Management

Dovepress

DovePress

Publish your work in this journal

Psychology Research and Behavior Management is an international, peer-reviewed, open access journal focusing on the science of psychology and its application in behavior management to develop improved outcomes in the clinical, educational, sports and business arenas. Specific topics covered in the journal include: Neuroscience, memory and decision making; Behavior modification and management; Clinical applications; Business and sports performance management; Social and developmental studies; Animal studies. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/psychology-research-and-behavior-management-journal

fi 🔰 in 🖪

811