

Prevalence of Depression and Its Associated Factors among Hemodialysis Patients in Hodeida City, Yemen [Response to Letter]

Sameer A Alkubati ^{1,2}, Khaled M Al-Sayaghi ^{3,4}, Basma Salameh ⁵, Abdulsalam M Halboup ^{6,7}, Waled A Ahmed ⁸, Mohannad J Alkuwaisi ¹, Mohamed A Zoromba ^{9,10}

¹Department of Medical Surgical Nursing, University of Hail, Hail, Saudi Arabia; ²Department of Nursing, Hodeida University, Hodeida, Yemen; ³Department of Medical Surgical Nursing, Taibah University, Al-Madinah Al-Munawarah, Saudi Arabia; ⁴Nursing Division, Sana'a University, Sana'a, Yemen; ⁵Department of Nursing, Arab American University, Jenin, Palestine; ⁶Department of Clinical Pharmacy and Pharmacy Practice, University of Science and Technology, Sana'a, Yemen; ⁷Discipline of Clinical Pharmacy, Universiti Sains Malaysia, Penang, Malaysia; ⁸Community Health Nursing Department, Al-Baha University, Al-Baha, Saudi Arabia; ⁹Department of Nursing, Prince Sattam Bin Abdulaziz University, Al Kharij, Saudi Arabia; ¹⁰Department of Psychiatric and Mental Health Nursing, Mansoura University, Mansoura, Egypt

Correspondence: Abdulsalam M Halboup, Departments of Clinical Pharmacy and Pharmacy Practice, University of Science and Technology, Sana'a, Yemen, Tel +967774960247, Email a.halboup@ust.edu.ye

Dear editor

We sincerely appreciate your interest in our study, "Prevalence of depression and its associated factors among hemodialysis patients in Hodeida city, Yemen". We would like to express our gratitude to all the authors for recognizing the study's importance in shedding light on an often overlooked population amid war-related crises. We also acknowledge the positive feedback on employing the OpenEpi website formula for sample size calculation and conducting a comparative analysis with international studies.¹

We acknowledge the insightful feedback from Wang et al,² particularly regarding the suggestion to use G*Power version 3.1.9.7 software for sample size calculation. We will explore this alternative for future research. Regarding the areas that need correction, the authors rightly pointed out two corrections. The female count in Table 2 needs correction to 51 instead of 52, and the odds ratio and confidence interval values on page 693 should be as follows: (odds ratio (OR) = 3.352; 95% confidence interval (CI) = 1.484–7.570; p = 0.004).

Regarding the limited variables included in the multivariate logistic regression, we recognize the potential benefits and risks associated with incorporating variables with P-values less than 0.1 or 0.2 from univariate analysis.³ In our study, we excluded the marital status variable due to one cell containing zero, which could impact interpretation.⁴ We also considered other factors based on the significant association in the chi-square test, as outlined in the limitations of the published study.

In the recommendation for future research is well-taken. We acknowledge the need for a large-scale, multicenter study to provide more robust conclusions. The suggestions for further investigation, including qualitative research on the subjective experiences of hemodialysis patients with depressive symptoms during the ongoing war in Yemen, establishing a control group, and utilizing structural equation models or mediating effects models, are valuable and will be considered in our future research endeavors.

In summary, we appreciate both the acknowledgment of the study's strengths and the constructive feedback for improvement provided in the letter to the editor.

Disclosure

The authors declare that they have no conflicts of interest in this communication.

References

1. Alkubati SA, Al-Sayaghi KM, Salameh B, et al. Prevalence of depression and its associated factors among hemodialysis patients in Hodeida City, Yemen. *J Multidiscip Healthc.* 2024;17:689–699. doi:10.2147/JMDH.S452935
2. Wang S, Song J, Jiang X. Prevalence of depression and its associated factors among hemodialysis patients in Hodeida City, Yemen [Letter]. *J Multidiscip Healthc.* 2024;17:1041–1042. doi:10.2147/JMDH.S466645
3. Hawkins DM. The problem of overfitting. *J Chem Inf Comput Sci.* 2004;44(1):1–12. doi:10.1021/ci0342472
4. Stoltzfus JC. Logistic regression: a brief primer. *Acad Emerg Med.* 2011;18(10):1099–1104. doi:10.1111/j.1553-2712.2011.01185.x

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Journal of Multidisciplinary Healthcare 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Journal of Multidisciplinary Healthcare editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Journal of Multidisciplinary Healthcare

Dovepress

Publish your work in this journal

The Journal of Multidisciplinary Healthcare is an international, peer-reviewed open-access journal that aims to represent and publish research in healthcare areas delivered by practitioners of different disciplines. This includes studies and reviews conducted by multidisciplinary teams as well as research which evaluates the results or conduct of such teams or healthcare processes in general. The journal covers a very wide range of areas and welcomes submissions from practitioners at all levels, from all over the world. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-multidisciplinary-healthcare-journal>

<https://doi.org/10.2147/JMDH.S467906>