

Thoughts about “Assessment of Quality of Diabetic Care in Teaching Hospitals in Ethiopia: In Comparison to International Guidelines” [Letter]

Sabrina Shahid , Mahdi Sadeghi , Fathima Simla Mohamed Sajahan

Medical Education, King's College London, London, UK

Correspondence: Mahdi Sadeghi, King's College London, Guy's Campus, Great Maze Pond, London, SE1 1UL, UK, Email Mahdi.Sadeghi@kcl.ac.uk

Dear editor

We thank Akale et al¹ for their insightful study assessing the quality of diabetic care within Ethiopia at the Yekatit 12 Hospital Medical College, where they compared the local practices against international guidelines. We commend them for their invaluable contribution to diabetic care in Ethiopia. As medical students in the UK who have interests in public health and improving patient outcomes, this study has provided insight into the current suboptimal level of care and the prospects of future studies to implement mediums to spread awareness.

The study conducted a cross-sectional study well and directly applied established international standards, the American Diabetes Association and the International Diabetic Federation, as validation tools.¹ However, the results could have been more generalisable if they had adopted a multi-centre study, allowing an increased diverse pool of participants and resulting in a more representative population within Ethiopia as approximately over 2.5 million individuals had a diabetes diagnosis.¹ Given the study's¹ relatively small sample size (n=250), predominantly with type 2 diabetes, it may not be representative of the population of individuals in Ethiopia with diabetes. Therefore, future studies could entail a larger participant pool and implement stratified sampling to ensure that subgroups (such as type 1 vs type 2 diabetes, varying socioeconomic status, urban vs rural residence) are adequately represented. Also, adopting a longitudinal approach or extending the data collection period can provide greater insight into the progression of care quality or provide more generalisable results by mitigating factors such as seasonal variation in glycaemic control.²

We commend Akale et al¹ for conducting interviews to collect quantitative data. However, the interviews could create social desirability bias, as the interviewees might have altered their responses due to not wanting to be judged or felt like it was a test of what they understood about their condition.³ Henceforth, we recommend adopting a mixed-method study – for example, using an anonymous questionnaire to collect intimate information. Furthermore, within Tables 1 and 2,¹ smaller ranges (eg age intervals) would provide more significant information, identifying more specific risk factors of variable relationships with diabetic care outcomes. Furthermore, we appreciate that comorbidities and lifestyle risk factors such as smoking or alcohol intake were recorded.¹ However, having more specific variables, such as the hypertension stages⁴ or smoking pack years,⁵ could be more beneficial. This data would also aid in providing increased information about the relationship between each identified risk factor and type 2 diabetes in Ethiopia.

As data was collected in 2022,¹ future research can look at current advancements in diabetic care within Ethiopia. Additionally, research can investigate the impact of targeted interventions in providing practical solutions to improve diabetic care. Overall, in hindsight, we thank Akale et al¹ for providing insight into the current level of diabetic care and, consequently, providing the rationale for the need for interventions to improve patient outcomes.

Disclosure

The author(s) report no conflicts of interest in this communication.

References

1. Akale M, Tadesse T, Arega B. Assessment of quality of diabetic care in teaching hospitals in Ethiopia: in comparison to international guidelines. *Diabet Metab Syndr Obes*. 2024;17:691–700. doi:10.2147/dmso.s441764
2. Ishii H, Suzuki H, Baba T, Nakamura K, Watanabe T. Seasonal variation of glycemic control in type 2 diabetic patients. *Diabetes Care*. 2001;24(8):1503. doi:10.2337/diacare.24.8.1503
3. Teh WL, Abidin E, Puv A, et al. Measuring social desirability bias in a multi-ethnic cohort sample: its relationship with self-reported physical activity, dietary habits, and factor structure. *BMC Public Health*. 2023;23(1). doi:10.1186/s12889-023-15309-3
4. Iqbal AM. Essential hypertension. StatPearls; 2023. <https://www.ncbi.nlm.nih.gov/books/NBK539859/>. Accessed April 02, 2024
5. De Vita MJ, Maisto SA, Ansell EB, Zale EL, Ditre JW. Pack-years of tobacco cigarette smoking as a predictor of spontaneous pain reporting and experimental pain reactivity. *Exp Clin Psychopharmacol*. 2019;27(6):552–560. doi:10.1037/pha0000258

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Diabetes, Metabolic Syndrome and Obesity 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Diabetes, Metabolic Syndrome and Obesity editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Diabetes, Metabolic Syndrome and Obesity

Dovepress

Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert opinion and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-journal>

<https://doi.org/10.2147/DMSO.S470330>